
4.

FURTHER DIFFERENTIATION

In this chapter, you will review differentiation and learn how to differentiate trigonometric, exponential and logarithmic functions, and inverse functions, including inverse trigonometric functions. You will also look at higher derivatives and anti-derivatives.

CHAPTER OUTLINE

4.01 Differentiation review
4.02 Derivative of exponential functions
4.03 Derivative of logarithmic functions
4.04 Derivative of trigonometric functions
4.05 Second derivatives
4.06 Anti-derivative graphs
4.07 Anti-derivatives
4.08 Further anti-derivatives



IN THIS CHAPTER YOU WILL:

- review differentiation
- differentiate trigonometric functions
- find the derivative of exponential and logarithmic functions
- understand the notation and find second and further derivatives
- identify and find anti-derivatives

Shutterstock.com/Doug James

TERMINOLOGY

anti-derivative: A function $F(x)$ whose derivative is $f(x)$, that is, $F'(x) = f(x)$. Also called the **primitive** or **integral** function.

anti-differentiation: The process of finding the original function given its derivative.

second derivative: The derivative $f''(x)$ or $\frac{d^2y}{dx^2}$; the derivative of the derivative $f'(x)$ or $\frac{dy}{dx}$.

4.01 Differentiation review

Chain rule

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} \qquad \frac{d}{dx}[f(x)]^n = f'(x)n[f(x)]^{n-1}$$

Product rule

If $y = uv$, then $\frac{dy}{dx} = v\frac{du}{dx} + u\frac{dv}{dx}$ or $y' = u'v + v'u$.

Quotient rule

If $y = \frac{u}{v}$, then $\frac{dy}{dx} = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$ or $y' = \frac{u'v - v'u}{v^2}$.

Rates of change

The **average rate of change** between 2 points (x_1, y_1) and (x_2, y_2) is the gradient:

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

The **instantaneous rate of change** at point (x, y) is the derivative $f'(x)$ or $\frac{dy}{dx}$.

EXAMPLE 1

Water is pumped into a dam according to the formula $Q = 3t^3 + 2t^2 + 270$ where Q is the amount of water in kL and t is time in hours. Find:

a the amount of water in the dam after 6 hours

b the average rate at which the water is pumped into the dam between 3 and 6 hours

c the rate of change after 6 hours

ISBN 9780170413220

Solution

a $Q = 3t^3 + 2t^2 + 270$

When $t = 6$

$Q = 3(6)^3 + 2(6)^2 + 270$

$= 990$

So there is 990 kL of water in the dam after 6 hours.

b When $t = 3$

$Q = 3(3)^3 + 2(3)^2 + 270$

$= 369$

$$\text{Average rate of change} = \frac{Q_2 - Q_1}{t_2 - t_1}$$

$$= \frac{990 - 369}{6 - 3}$$

$$= \frac{621}{3}$$

$$= 207$$

So average rate of change is 207 kL h^{-1}.

c $\frac{dQ}{dt} = 9t^2 + 4t$

When $t = 6$

$\frac{dQ}{dt} = 9(6)^2 + 4(6)$

$= 348$

So the rate of increase after 6 hours is 348 kL h^{-1}.

Exercise 4.01 Differentiation review

1 Differentiate each function.

a $3x^4 - 2x^3 + 7x - 4$

b $2x + 5$

c $6x^2 - 3x - 2$

2 Find the derivative $f'(x)$ given $f(x) = 4x^5 + 9x^2$.

3 Find $\frac{dx}{dt}$ if $x = 2\pi t^3 - 3t^2 + 1$.

4 Find $f'(-2)$ when $f(x) = 8x^3 + 5x - 2$.

5 Differentiate:

a x^{-5} **b** $x^{\frac{2}{3}}$ **c** $\frac{1}{x^2}$

d $\sqrt[4]{x}$ **e** $-\frac{5}{x^4}$

6 Find the derivative of $y = \sqrt[3]{x}$ at the point where $x = 8$.

ISBN 9780170413220

7 Differentiate:

a $(3x-1)^7$ **b** $(x^2-x+2)^3$ **c** $\sqrt{7x-2}$

d $\dfrac{1}{3x-2}$ **e** $\sqrt[3]{x^2-3}$

8 Find the derivative of:

a $x^2(x+4)$ **b** $(2x-1)(6x+5)$ **c** $4x(x^2+1)$

d $(4x+3)(x^2-1)^2$ **e** $2x^3\sqrt{x+1}$

9 Differentiate:

a $\dfrac{2x+3}{x-5}$ **b** $\dfrac{x^3}{4x-7}$ **c** $\dfrac{x^2+3}{2x-3}$

d $\dfrac{3x+1}{(2x+9)^2}$ **e** $\dfrac{3x+4}{\sqrt{2x-1}}$

10 Find the gradient of the tangent to the curve:

a $y=x^2-2x+5$ at the point where $x=-2$

b $f(x)=x^3-3$ at the point $(-1,-4)$

11 Find the gradient of the normal to the curve:

a $f(x)=3x^4+x^2-2$ at the point where $x=-1$

b $y=x^2+x-3$ at the point $(-3,3)$

12 Find the equation of the tangent to the curve:

a $y=2x^2-5x-6$ at the point $(3,-3)$

b $y=5x^3-2x^2-x$ at the point where $x=2$

13 Find the equation of the normal to the curve:

a $f(x)=x^3+2x^2-3x-5$ at the point $(-1,-1)$

b $y=x^2-3x+1$ at the point where $x=3$

14 For the curve $y=x^2-8x+15$, find any values of x for which $\dfrac{dy}{dx}=0$.

15 Find the coordinates of the points at which the curve $y=x^3-2$ has a tangent with gradient 12.

16 Function $f(x)=x^2+x-4$ has a tangent parallel to the line $3x+y-4=0$ at point P. Find the equation of the tangent at P.

17 Find the coordinates of P if the gradient of the tangent to $y=\sqrt{x}$ is $\dfrac{1}{4}$ at point P.

ISBN 9780170413220

18 For the curve $y = \dfrac{5x-3}{4x+1}$ at the point where $x = 0$, find the equation of:

a the tangent **b** the normal

19 Find a formula for the rate of change $\dfrac{dQ}{dt}$ given:

a $Q = 3t^2 + 8$ **b** $Q = \dfrac{2}{t-3}$ **c** $Q = \sqrt[3]{2x+3}$

20 The mass M in kg of a snowball as it rolls down a hill over time t seconds is given by $M = t^2 + 3t + 4$.

a Find the average rate at which the mass changes between:

i 2 and 5 seconds **ii** 6 and 8 seconds

b Find the rate at which the mass is changing after:

i 5 seconds **ii** a minute

21 According to Boyle's Law, the pressure of a gas in pascals (Pa) is given by the formula $P = \dfrac{k}{V}$, where k is a constant and V is the volume of the gas in m^3. If $k = 250$ for a certain gas, find the rate of change in the pressure when $V = 10.7$.

22 The height of a ball in metres is given by $h = 4t - 2t^2$ where t is time in seconds.

a Find the height after:

i 1 s **ii** 1.5 s

b How long does it take for the ball to reach the ground?

c Find the velocity of the ball after:

i 0.5 s **ii** 1 s **iii** 2 s

4.02 Derivative of exponential functions

WS Derivatives of exponential functions

You learned how to differentiate $y = e^x$ in Year 11, in Chapter 8, *Exponential and logarithmic functions*.

Differentiation rules for e^x

$$\frac{d}{dx}e^x = e^x$$

$$\text{If } y = e^{f(x)} \text{ then } \frac{dy}{dx} = f'(x)\,e^{f(x)}$$

EXAMPLE 2

a If $f(x) = 3e^x$, find the equation of the tangent to the curve at $(2, 3e^2)$.

b Differentiate :

i x^2e^x **ii** e^{8x} **iii** e^{5x-2}

Solution

a $f(x) = 3e^x$

$f'(x) = 3e^x$

At $(2, 3e^2)$

$f'(2) = 3e^2$

So $m = 3e^2$

Equation:

$$y - y_1 = m(x - x_1)$$
$$y - 3e^2 = 3e^2(x - 2)$$
$$= 3e^2x - 6e^2$$
$$y = 3e^2x - 3e^2$$

(or $3e^2x - y - 3e^2 = 0$)

b **i** $y' = u'v + v'u$

where $u = x^2$ and $v = e^x$

$u' = 2x \qquad v' = e^x$

$y' = 2xe^x + e^xx^2$

$= xe^x(2 + x)$

ii $\dfrac{dy}{dx} = ae^{ax}$

$= 8e^{8x}$

iii $\dfrac{dy}{dx} = f'(x)e^{f(x)}$

$= 5e^{5x-2}$

We can differentiate other exponential functions.

EXAMPLE 3

Differentiate 2^x.

Solution

$2 = e^{\ln 2}$

$2^x = (e^{\ln 2})^x$

$= e^{x \ln 2}$

$\dfrac{dy}{dx} = \ln 2 \; e^{x \ln 2}$ ← ln 2 is a constant

$= \ln 2 \times 2^x$

$= 2^x \ln 2$

ISBN 9780170413220

Derivative of a^x

$$\text{If } y = a^x, \text{ then } \frac{dy}{dx} = a^x \ln a$$

The proof of this has the same steps as in the previous example.

Exercise 4.02 Derivative of exponential functions

1 Differentiate:

a	e^{7x}	**b**	e^{-x}	**c**	e^{6x-2}	**d**	e^{x^2+1}
e	e^{x^3+5x+7}	**f**	e^{5x}	**g**	e^{-2x}	**h**	e^{10x}
i	$e^{2x}+x$	**j**	x^2+2x+e^{1-x}	**k**	$(x+e^{4x})^5$	**l**	xe^{2x}
m	$\dfrac{e^{3x}}{x^2}$	**n**	x^3e^{5x}	**o**	$\dfrac{e^{2x+1}}{2x+5}$		

2 If $f(x) = e^{3x-2}$ find the exact value of $f'(1)$.

3 Find the derivative of:

a 3^x **b** 10^x **c** 2^{3x-4}

4 Find the gradient of the tangent to the curve $y = e^{5x}$ at the point where $x = 0$.

5 Find the equation of the tangent to the curve $y = e^{2x} - 3x$ at the point (0, 1).

6 For the curve $y = e^{3x}$ at the point where $x = 1$, find the exact gradient of:

a the tangent **b** the normal

7 For the curve $y = e^{x^2}$ at the point $(1, e)$, find the equation of:

a the tangent **b** the normal

8 Find the equation of the tangent to the curve $y = 4^{x+1}$ at the point (0, 4).

9 The population of a city is given by $P = 24\,500e^{0.038t}$ where t is time in years.

a Find the population after:

i 5 years **ii** 10 years

b Find the average rate of change in population between:

i the 1st and 5th years **ii** the 5th and 10th years

c Find the rate of change in population after:

i 5 years **ii** 10 years

10 The displacement of a particle is given by $s = 10e^{2t} - 5t$ cm after t minutes.

a Find the average rate of change in displacement between 1 and 5 minutes.

b Find the rate of change in displacement after:

i 1 minute **ii** 2 minutes **iii** 8 minutes

ISBN 9780170413220

11 A radioactive substance has a mass of $M = 20e^{-0.021t}$ in grams over time t years.

a Find the initial mass.

b Find the mass after 50 years.

c Find the average rate of change in mass between 50 and 100 years.

d Find the rate of change in mass after:

i 50 years **ii** 100 years **iii** 200 years

12 An object moves according to the formula $x = 3e^{2t}$ where x is displacement in cm and t is time in s.

a Find the displacement at 5 s.

b Find the velocity at 5 s.

INVESTIGATION

DERIVATIVE OF A LOGARITHMIC FUNCTION

Draw the derivative (gradient) function of a logarithm function.

What is the shape of the derivative function?

WS Derivatives of logarithmic functions

WS Exponential and logarithmic functions

4.03 Derivative of logarithmic functions

Logarithm rules

$$\text{If } y = a^x \text{ then } \log_a y = x$$

$$\log_a xy = \log_a x + \log_a y$$

$$\log_a \frac{x}{y} = \log_a x - \log_a y$$

$$\log_a x^n = n \log_a x$$


$$\log_a x = \frac{\log_b x}{\log_b a}$$




iStock.com/Anna_Om

ISBN 9780170413220

To find the derivative of a logarithmic function, notice that the gradient of the function is always positive but is decreasing.



The derivative function of a logarithmic function is a hyperbola.



There is a special rule for $y = \ln x$.

Derivative of $y = \ln x$

If $y = \ln x$, then $\frac{dy}{dx} = \frac{1}{x}$ where $x > 0$.

Proof

$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$

Given $y = \ln x = \log_e x$

Then $x = e^y$

$\frac{dx}{dy} = e^y$

$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$

$= \frac{1}{e^y}$

$= \frac{1}{x}$



Shutterstock.com/Gary Le Feuvre

ISBN 9780170413220

EXAMPLE 4

a Differentiate $(\ln x + 1)^3$.

b Find the equation of the tangent to the curve $y = \ln x$ at the point $(3, \ln 3)$.

Solution

a $(\ln x + 1)^3$ is a composite function in the form $y = [f(x)]^n$.

$$\frac{dy}{dx} = f'(x)\, nf(x)^{n-1}$$

$$= \frac{1}{x} \times 3(\ln x + 1)^2$$

$$= \frac{3(\ln x + 1)^2}{x}$$

b $\frac{dy}{dx} = \frac{1}{x}$

At $(3, \ln 3)$

$$\frac{dy}{dx} = \frac{1}{3}$$

So $m = \frac{1}{3}$

Equation:

$$y - y_1 = m(x - x_1)$$

$$y - \ln 3 = \frac{1}{3}(x - 3)$$

$$3y - 3\ln 3 = x - 3$$

$$0 = x - 3y - 3 + 3\ln 3$$

Chain rule

If $y = \ln f(x)$, then $\frac{dy}{dx} = \frac{f'(x)}{f(x)}$ where $f(x) > 0$

Proof

$y = \ln f(x)$ is a composite function.

Let $y = \ln u$ and $u = f(x)$

$$\frac{dy}{du} = \frac{1}{u} \quad \text{and} \quad \frac{du}{dx} = f'(x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$= \frac{1}{u} \times f'(x)$$

$$= \frac{1}{f(x)} \times f'(x)$$

$$= \frac{f'(x)}{f(x)}$$

ISBN 9780170413220

EXAMPLE 5

a Differentiate:

i $\ln(x^2 - 3x + 1)$ **ii** $\ln\left(\dfrac{x+1}{3x-4}\right)$

b Find the gradient of the normal to the curve $y = \ln(x^3 - 5)$ at the point where $x = 2$.

Solution

a **i** $\dfrac{dy}{dx} = \dfrac{f'(x)}{f(x)}$

$= \dfrac{2x-3}{x^2-3x+1}$

ii It is easier to simplify first using log laws.

$y = \ln\left(\dfrac{x+1}{3x-4}\right)$

$= \ln(x+1) - \ln(3x-4)$

$\dfrac{dy}{dx} = \dfrac{f'(x)}{f(x)}$

$= \dfrac{1}{x+1} - \dfrac{3}{3x-4}$

$= \dfrac{1(3x-4)}{(x+1)(3x-4)} - \dfrac{3(x+1)}{(3x-4)(x+1)}$

$= \dfrac{3x-4-3(x+1)}{(x+1)(3x-4)}$

$= \dfrac{3x-4-3x-3}{(x+1)(3x-4)}$

$= \dfrac{-7}{(x+1)(3x-4)}$

b $\dfrac{dy}{dx} = \dfrac{3x^2}{x^3-5}$

When $x = 2$

$\dfrac{dy}{dx} = \dfrac{3(2)^2}{2^3-5}$

$= 4$

$m_1 = 4$

The normal is perpendicular to the tangent:

$m_1 m_2 = -1$

$4m_2 = -1$

$m_2 = -\dfrac{1}{4}$

ISBN 9780170413220

We can differentiate logarithmic functions with a different base, a.

EXAMPLE 6

Differentiate $y = \log_2 x$.

Solution

$y = \log_2 x$

$= \dfrac{\ln x}{\ln 2}$ using the change of base law

$= \dfrac{1}{\ln 2} \ln x$

$\dfrac{dy}{dx} = \dfrac{1}{\ln 2} \times \dfrac{1}{x}$ ← ln 2 is a constant

$= \dfrac{1}{x \ln 2}$

Derivative of $\log_a x$

If $y = \log_a x$, then $\dfrac{dy}{dx} = \dfrac{1}{x \ln a}$

The proof of this has the same steps as in the above example.

Exercise 4.03 Derivative of logarithmic functions

1 Differentiate:

a	$x + \ln x$	**b**	$1 - \ln 3x$	**c**	$\ln (3x + 1)$
d	$\ln (x^2 - 4)$	**e**	$\ln (5x^3 + 3x - 9)$	**f**	$\ln (5x + 1) + x^2$
g	$3x^2 + 5x - 5 + \ln 4x$	**h**	$\ln (8x - 9) + 2$	**i**	$\ln (2x + 4)(3x - 1)$
j	$\ln\left(\dfrac{4x+1}{2x-7}\right)$	**k**	$(1 + \ln x)^5$	**l**	$(\ln x - x)^9$
m	$(\ln x)^4$	**n**	$(x^2 + \ln x)^6$	**o**	$x \ln x$
p	$\dfrac{\ln x}{x}$	**q**	$(2x + 1)\ln x$	**r**	$x^3 \ln (x + 1)$
s	$\ln (\ln x)$	**t**	$\dfrac{\ln x}{x - 2}$	**u**	$\dfrac{e^{2x}}{\ln x}$
v	$e^x \ln x$	**w**	$5(\ln x)^2$		

2 Find $f'(1)$ if $f(x) = \ln \sqrt{2 - x}$.

3 Find the derivative of $\log_{10} x$.

4 Find the equation of the tangent to the curve $y = \ln x$ at the point $(2, \ln 2)$.

ISBN 9780170413220

5 Find the equation of the tangent to the curve $y = \ln(x - 1)$ at the point where $x = 2$.

6 Find the gradient of the normal to the curve $y = \ln(x^4 + x)$ at the point $(1, \ln 2)$.

7 Find the exact equation of the normal to the curve $y = \ln x$ at the point where $x = 5$.

8 Find the equation of the tangent to the curve $y = \ln(5x + 4)$ at the point where $x = 3$.

9 Find the derivative of $\log_3(2x + 5)$.

10 Find the equation of the normal to the curve $y = \log_2 x$ at the point where $x = 2$.

11 The formula for the time t in years for kangaroo population growth on Kangaroo Island is given by $t = \dfrac{\ln\left(\dfrac{P}{20\,000}\right)}{0.021}$.

a What is the initial population?

b Find correct to one decimal place the time it takes for the population to grow to:

i 25 000

ii 50 000

c Change the subject of the equation to P.

d Find correct to the nearest whole number the average rate of change in population between 2 and 5 years.

e Find correct to the nearest whole number the rate at which the population is growing after:

i 3 years

ii 5 years

iii 10 years

CLASS INVESTIGATION

DERIVATIVE OF TRIGONOMETRIC FUNCTIONS

1 Draw the derivative (gradient) function of sine, cosine and tangent functions.

What is the shape of the derivative function of each graph?

2 By substituting values of x in radians close to 0, find approximations to $\lim\limits_{x \to 0} \dfrac{\sin x}{x}$, $\lim\limits_{x \to 0} \dfrac{\tan x}{x}$ and $\lim\limits_{x \to 0} \dfrac{\cos x}{x}$.

3 Differentiate by first principles to find the derivative of each trigonometric function using the above limits. The sine function will use the EXT1 trigonometric identity $\sin(A + B) = \sin A \cos B + \cos A \sin B$.

ISBN 9780170413220

WS Derivatives of trigonometric functions

WS Trigonometric functions and gradient


WS Further trigonometric equations

WS Differentiating trigonometric functions


4.04 Derivative of trigonometric functions

Derivative of sin x

We can sketch the derivative (gradient) function of $y = \sin x$.



The sketch of the gradient function is $y = \cos x$.



Derivative of sin x

$$\text{If } y = \sin x, \text{ then } \frac{dy}{dx} = \cos x$$

Proof

This proof uses trigonometric results from the investigation on the previous page.

$$f'(x) = \lim_{h\to 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h\to 0} \frac{\sin(x+h) - \sin x}{h}$$

$$= \lim_{h\to 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h}$$

$$= \lim_{h\to 0} \frac{\sin x(\cos h - 1) + \cos x \sin h}{h}$$

$$= \lim_{h\to 0} \frac{\sin x(\cos h - 1)}{h} + \lim_{h\to 0} \frac{\cos x \sin h}{h}$$

$$= \sin x \lim_{h\to 0} \frac{(\cos h - 1)}{h} + \cos x \lim_{h\to 0} \frac{\sin h}{h}$$

$$= \sin x \lim_{h\to 0} \frac{(\cos h - 1)}{h} + \cos x \lim_{h\to 0} \frac{\sin h}{h}$$

$$= \sin x \times 0 + \cos x \times 1$$

$$= \cos x$$

ISBN 9780170413220

EXAMPLE 7

a Differentiate $y = x \sin x$.

b Find the equation of the tangent to the curve $y = \sin x$ at the point $(\pi, 0)$.

Solution

a $y = x \sin x$ is in the form $y = uv$

where $u = x$ and $v = \sin x$

$u' = 1$ and $v' = \cos x$

$y' = u'v + v'u$

$= 1 \times \sin x + \cos x \times x$

$= \sin x + x \cos x$

b $\dfrac{dy}{dx} = \cos x$

At $(\pi, 0)$

$\dfrac{dy}{dx} = \cos \pi$

$= -1$

So $m = -1$

Equation:

$y - y_1 = m(x - x_1)$


$y - 0 = -1(x - \pi)$

$y = -x + \pi$


or $x + y - \pi = 0$

Derivative of cos *x*

We can sketch the derivative (gradient) function of $y = \cos x$.



The sketch of the gradient function below is $y = -\sin x$.



Derivative of cos *x*

If $y = \cos x$, then $\dfrac{dy}{dx} = -\sin x$

ISBN 9780170413220

You can prove this in a similar way to the derivative of $y = \sin x$. A simpler proof involves changing $\cos x$ into $\sin\left(\frac{\pi}{2} - x\right)$ and using the derivative of $y = \sin x$.

EXAMPLE 8

a Find the derivative of $y = \cos x$ at the point where $x = \frac{\pi}{3}$.

b Find the equation of the tangent to $y = \cos x$ at this point.


Solution

a $\frac{dy}{dx} = -\sin x$

When $x = \frac{\pi}{3}$

$\frac{dy}{dx} = -\sin\frac{\pi}{3}$

$= -\frac{\sqrt{3}}{2}$



b When $x = \frac{\pi}{3}$

$y = \cos\frac{\pi}{3}$

$= \frac{1}{2}$

Equation:

$y - y_1 = m(x - x_1)$

$y - \frac{1}{2} = -\frac{\sqrt{3}}{2}\left(x - \frac{\pi}{3}\right)$

$2y - 1 = -\sqrt{3}\left(x - \frac{\pi}{3}\right)$ (multiplying both sides by 2)

$= -\sqrt{3}x + \frac{\pi\sqrt{3}}{3}$


$6y - 3 = -3\sqrt{3}x + \pi\sqrt{3}$ (multiplying both sides by 3)

$3\sqrt{3}x + 6y - 3 - \pi\sqrt{3} = 0$


ISBN 9780170413220

Derivative of tan *x*

We can sketch the derivative (gradient) function of $y = \tan x$. Notice that the gradient function will have asymptotes in the same place as the original graph, because this is where the tangent is vertical and the gradient is undefined.



The gradient function is $y = \sec^2 x$, where $\sec x = \dfrac{1}{\cos x}$.



Derivative of tan *x*

$$\text{If } y = \tan x, \text{ then } \frac{dy}{dx} = \sec^2 x$$

You can prove this in a similar way to the derivative of $y = \sin x$. A simpler proof involves changing $\tan x$ into $\dfrac{\sin x}{\cos x}$ and using the quotient rule.

ISBN 9780170413220

EXAMPLE 9

a Differentiate $y = \dfrac{\tan x}{3x^2}$.

b Find the gradient of the tangent to the curve $f(x) = \tan x$ at the point where $x = \dfrac{\pi}{4}$.

Solution

a $y = \dfrac{\tan x}{3x^2}$ is in the form $y = \dfrac{u}{v}$.

$u = \tan x \quad$ and $\quad v = 3x^2$

$u' = \sec^2 x \qquad v' = 6x$

$$y' = \frac{u'v - v'u}{u^2}$$

$$= \frac{\sec^2 x \times 3x^2 - 6x \times \tan x}{(3x^2)^2}$$

$$= \frac{3x(x \sec^2 x - 2\tan x)}{9x^4}$$

$$= \frac{x \sec^2 x - 2\tan x}{3x^3}$$

b $\dfrac{dy}{dx} = \sec^2 x$

At $x = \dfrac{\pi}{4}$

$$\frac{dy}{dx} = \sec^2 \frac{\pi}{4}$$

$$= (\sqrt{2})^2$$

$$= 2$$

Chain rule

If $y = \sin f(x)$, then $\dfrac{dy}{dx} = f'(x) \cos f(x)$

If $y = \cos f(x)$, then $\dfrac{dy}{dx} = -f'(x) \sin f(x)$

If $y = \tan f(x)$, then $\dfrac{dy}{dx} = f'(x) \sec^2 f(x)$

ISBN 9780170413220

Here is the proof for $y = \sin f(x)$. The others are similar.

Proof

$y = \sin f(x)$ is a composite function

where $y = \sin u$ and $u = f(x)$

$$\frac{dy}{du} = \cos u \text{ and } \frac{du}{dx} = f'(x)$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$= \cos u \times f'(x)$$

$$= f'(x) \cos u$$

$$= f'(x) \cos f(x)$$

EXAMPLE 10

a Differentiate each function.

i $y = \sin 7x$ **ii** $y = \cos\left(4x^3 + \frac{\pi}{3}\right)$ **iii** $y = \tan(5x - \pi)$

b Find the gradient of the normal to the curve $f(x) = \cos\frac{x}{2}$ at the point where $x = \pi$.

Solution

a **i** $\frac{dy}{dx} = f'(x) \cos f(x)$

$= 7 \cos 7x$

ii $\frac{dy}{dx} = -f'(x) \sin f(x)$

$= -12x^2 \sin\left(4x^3 + \frac{\pi}{3}\right)$

iii $\frac{dy}{dx} = f'(x) \sec^2 f(x)$

$= 5 \sec^2(5x - \pi)$

b $\frac{dy}{dx} = -f'(x) \sin f(x)$

$= -\frac{1}{2} \sin\frac{x}{2}$

At $x = \pi$:

$\frac{dy}{dx} = -\frac{1}{2} \sin\frac{\pi}{2}$

$= -\frac{1}{2} \times 1$

$= -\frac{1}{2}$

So $m_1 = -\frac{1}{2}$

Normal is perpendicular to the tangent:

$m_1 m_2 = -1$

$-\frac{1}{2} m_2 = -1$

$m_2 = 2$

While trigonometric functions are usually expressed in radians, we can differentiate angles *in degrees* by using the conversion $\pi = 180°$.

EXAMPLE 11

Differentiate $y = \sin x°$.

Solution

$180° = \pi$ radians

$$1° = \frac{\pi}{180}$$

$$x° = \frac{\pi x}{180}$$

So $y = \sin x°$ becomes

$$y = \sin \frac{\pi x}{180}$$

$$\frac{dy}{dx} = f'(x) \cos f(x)$$

$$= \frac{\pi}{180} \cos \frac{\pi x}{180}$$

$$= \frac{\pi}{180} \cos x°$$

We can also differentiate composite functions involving trigonometric functions.

EXAMPLE 12

Differentiate:

a $\tan (e^x)$

b $\ln (\cos x)$

Solution

a
$$\frac{dy}{dx} = f'(x) \sec^2 f(x)$$

$$= e^x \sec^2 (e^x)$$

b
$$\frac{dy}{dx} = \frac{f'(x)}{f(x)}$$

$$= \frac{-\sin x}{\cos x}$$

$$= -\tan x$$

Exercise 4.04 Derivative of trigonometric functions

1 Differentiate:

a $\sin 4x$ **b** $\cos 3x$ **c** $\tan 5x$

d $\tan (3x + 1)$ **e** $\cos (-x)$ **f** $3 \sin x$

g $4 \cos (5x - 3)$ **h** $2 \cos (x^3)$ **i** $7 \tan (x^2 + 5)$

j $\sin 3x + \cos 8x$ **k** $\tan (\pi + x) + x^2$ **l** $x \tan x$

m $\sin 2x \tan 3x$ **n** $\dfrac{\sin x}{2x}$ **o** $\dfrac{3x + 4}{\sin 5x}$

ISBN 9780170413220

p $(2x + \tan 7x)^9$ **q** $\sin^2 x$ **r** $3\cos^3 5x$

s $e^x - \cos 2x$ **t** $\sin(1 - \ln x)$ **u** $\sin(e^x + x)$

v $\ln(\sin x)$ **w** $e^{3x}\cos 2x$ **x** $\dfrac{e^{2x}}{\tan 7x}$

2 Find the gradient of the tangent to the curve $y = \tan 3x$ at the point where $x = \dfrac{\pi}{9}$.

3 Find the equation of the tangent to the curve $y = \sin(\pi - x)$ at the point $\left(\dfrac{\pi}{6}, \dfrac{1}{2}\right)$ in exact form.

4 Differentiate $\ln(\cos x)$.

5 Find the exact gradient of the normal to $y = \sin 3x$ at the point where $x = \dfrac{\pi}{18}$.

6 Differentiate $e^{\tan x}$.

7 Find the equation of the normal to the curve $y = 3\sin 2x$ at the point where $x = \dfrac{\pi}{8}$ in exact form.

8 Show that $\dfrac{d}{dx}[\ln(\tan x)] = \tan x + \cot x$.

9 Differentiate each function.

a $y = \tan x°$ **b** $y = 3\cos x°$ **c** $y = \dfrac{\sin x°}{5}$

10 Find the derivative of $\cos x \sin^4 x$.

11 The population of salmon in a salmon farm grows and reduces as fish are born and sold.

The population is given by $P = 225\cos\dfrac{2\pi t}{9} + 750$ where t is time in days.

a What is the centre of the population?

b What is the minimum number of salmon in the farm at any one time?

c What is the maximum population?

d At what times is the population 700?

e At what rate is the population changing after:

i 3 days? **ii** a week? **iii** 10 days? **iv** 18 days?

f At what times is the population growing at the rate of 25 fish per day?

12 The tide was measured over time at a beach at Merimbula and given the formula $D = 8\sin\dfrac{\pi t}{6} + 9$ where D is depth of water in metres and t is time in hours.

a How deep was the water:

i initially? **ii** after 5 hours?

b When was the water 10 m deep?

c At what rate was the depth changing after:

i 3 hours? **ii** 11 hours? **iii** 12 hours?

d At what times was the depth of water decreasing by 3 m h^{-1}?

ISBN 9780170413220

WS The second derivative

WS First and second derivatives

4.05 Second derivatives

Second derivative

Differentiating $f(x)$ gives $f'(x)$, the first derivative.

Differentiating $f'(x)$ gives $f''(x)$, the **second derivative**.

It is also possible to differentiate further.

Using function notation, differentiating several times gives $f'(x), f''(x), f'''(x)$ and so on.

Using $\frac{dy}{dx}$ notation, differentiating several times gives $\frac{d^2y}{dx^2}, \frac{d^3y}{dx^3}$ and so on.

The notation $\frac{d^2y}{dx^2}$ comes from $\frac{d^2}{dx^2}(y)$.

EXAMPLE 13

a Find the first 4 derivatives of $f(x) = x^3 - 4x^2 + 3x - 2$.

b Find the second derivative of $y = (2x + 5)^7$.

c If $f(x) = 4 \cos 3x$, show that $f''(x) = -9f(x)$

Solution

a $f'(x) = 3x^2 - 8x + 3$

$f''(x) = 6x - 8$

$f'''(x) = 6$

$f''''(x) = 0$

b $\frac{dy}{dx} = f'(x) \times nf(x)^{n-1}$

$= 2 \times 7(2x + 5)^6$

$= 14(2x + 5)^6$

$\frac{d^2y}{dx^2} = f'(x) \times nf(x)^{n-1}$

$= 2 \times 6 \times 14(2x + 5)^5$

$= 168(2x + 5)^5$

c $f'(x) = -f'(x) \times \sin f(x)$

$= -3 \times 4 \sin 3x$

$= -12 \sin 3x$

$f''(x) = f'(x) \times \cos f(x)$

$= 3 \times (-12 \cos 3x)$

$= -36 \cos 3x$

$= -9(4 \cos 3x)$

$= -9f(x)$ since $f(x) = 4 \cos 3x$

ISBN 9780170413220

Exercise 4.05 Second derivatives

1 Find the first 4 derivatives of $x^7 - 2x^5 + x^4 - x - 3$.

2 If $f(x) = x^9 - 5$, find $f''(x)$.

3 Find $f'(x)$ and $f''(x)$ if $f(x) = 2x^5 - x^3 + 1$.

4 Find $f'(1)$ and $f''(-2)$, given $f(t) = 3t^4 - 2t^3 + 5t - 4$.

5 Find the first 3 derivatives of $x^7 - 2x^6 + 4x^4 - 7$.

6 Find the first and second derivatives of $y = 2x^2 - 3x + 3$.

7 If $f(x) = x^4 - x^3 + 2x^2 - 5x - 1$, find $f'(-1)$ and $f''(2)$.

8 Find the first and second derivatives of x^{-4}.

9 If $g(x) = \sqrt{x}$, find $g''(4)$.

10 Given $h = 5t^3 - 2t^2 + t + 5$, find $\dfrac{d^2h}{dt^2}$ when $t = 1$.

11 Find any values of x for which $\dfrac{d^2y}{dx^2} = 3$, given $y = 3x^3 - 2x^2 + 5x$.

12 Find all values of x for which $f''(x) > 0$ given that $f(x) = x^3 - x^2 + x + 9$.

13 Find the first and second derivatives of $(4x - 3)^5$.

14 Find $f'(x)$ and $f''(x)$ if $f(x) = \sqrt{2 - x}$.

15 Find the first and second derivatives of $f(x) = \dfrac{x + 5}{3x - 1}$.

16 Find $\dfrac{d^2v}{dt^2}$ if $v = (t + 3)(2t - 1)^2$.

17 Find the value of b in $y = bx^3 - 2x^2 + 5x + 4$ if $\dfrac{d^2y}{dx^2} = -2$ when $x = \dfrac{1}{2}$.

18 Find $f''(1)$ if $f(t) = t(2t - 1)^7$.

19 Find the value of b if $f(x) = 5bx^2 - 4x^3$ and $f''(-1) = -3$.

20 If $y = e^{4x} + e^{-4x}$, show that $\dfrac{d^2y}{dx^2} = 16y$.

21 Prove that $\dfrac{d^2y}{dx^2} - 3\dfrac{dy}{dx} + 2y = 0$ given $y = 3e^{2x}$.

ISBN 9780170413220

22 Show that $\frac{d^2y}{dx^2} = b^2y$ for $y = ae^{bx}$.

23 Find the value of n if $y = e^{3x}$ satisfies the equation $\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + ny = 0$.

24 Show that $\frac{d^2y}{dx^2} = -25y$ if $y = 2\cos 5x$.

25 Given $f(x) = -2\sin x$, show that $f''(x) = -f(x)$.

26 If $y = 2\sin 3x - 5\cos 3x$, show that $\frac{d^2y}{dx^2} = -9y$.

27 Find values of a and b if $\frac{d^2y}{dx^2} = ae^{3x}\cos 4x + be^{3x}\sin 4x$, given $y = e^{3x}\cos 4x$.

28 Find the exact value of $f''(2)$ if $f(x) = x\sqrt{3x-4}$.

29 The displacement of a particle moving in a straight line is given by $x = 2t^3 - 5t^2 + 7t + 8$, where x is in metres and t is in seconds.

- **a** Find the initial displacement.
- **b** Find the displacement after 3 seconds.
- **c** Find the velocity after 3 seconds.
- **d** Find the acceleration after 3 seconds.

30 The height in cm of a pendulum as it swings is given by $h = 8\cos \pi t + 12$ where t is time in seconds.

- **a** What is the height of the pendulum after 3 s?
- **b** What is the maximum and minimum height of the pendulum?
- **c** What is the velocity of the pendulum after:
  - **i** 1 s?
  - **ii** 1.5 s?
- **d** What is the acceleration of the pendulum:
  - **i** initially?
  - **ii** after 1 s?
  - **iii** after 1.5 s?

ISBN 9780170413220

4.06 Anti-derivative graphs




Shapes of curves

The process of finding the original function $y = f(x)$ given the derivative $y = f'(x)$ is called **anti-differentiation**, and the original function is called the **anti-derivative** function, also called the **primitive** or **integral function**.


Anti-derivative graphs

EXAMPLE 14

Sketch the graph of the anti-derivative (primitive function) given the graph of the derivative function below and an initial condition, or starting point, of (0, 2).



Solution


Remember that when you sketch a derivative function, the x-intercepts are where the original function has zero gradient, or stationary (turning) points.

On this graph the stationary points are at $x = x_1$ and $x = x_2$.

Above the x-axis shows where the original function has a positive gradient (it is increasing). On this graph, this is where $x < x_1$ and $x > x_2$.

Below the x-axis shows where the original function has a negative gradient (it is decreasing). On this graph, this is where $x_1 < x < x_2$.


We can sketch this information together with the point (0, 2):



ISBN 9780170413220

We are not given enough information to sketch a unique graph. There is no way of knowing what the y values of the stationary points are or the stretch or compression of the graph. Also, if we are not given a fixed point on the function, we could sketch many graphs that satisfy the information from the derivative function.


The anti-derivative gives a **family** of curves.




Exercise 4.06 Anti-derivative graphs

1 For each function graphed, sketch the graph of the anti-derivative function given it passes through:


a (0, –1)




b (1, 2)




c (0, 3)



d (–1, –1)




e (0, 1)




ISBN 9780170413220

2 Sketch a family of graphs that could represent the anti-derivative function of each graph.


a




b




c




d




e




3 The anti-derivative function of the graph below passes through (0, –1). Sketch its graph.



4 Sketch the graph of the anti-derivative function of $y = \cos x$ given that it passes through (0, 0).



5 Sketch a family of anti-derivative functions for the graph.



ISBN 9780170413220

INVESTIGATION

ANTI-DERIVATIVE OF $y = x^n$

1 Differentiate:

a x^2 **b** $x^2 + 5$ **c** $x^2 - 7$ **d** $x^2 + 4$ **e** $x^2 - 75$

What would be the anti-derivative of $2x$?

2 Differentiate:

a x^3 **b** $x^3 + 1$ **c** $x^3 + 6$ **d** $x^3 - 2$ **e** $x^3 - 14$

What would be the anti-derivative of $3x^2$?

3 Differentiate:

a x^4 **b** $x^4 - 3$ **c** $x^4 + 2$ **d** $x^4 + 10$ **e** $x^4 - 1$

What would be the anti-derivative of $4x^3$?

4 Differentiate:

a x^n **b** $x^n + 7$ **c** $x^n + 9$ **d** $x^n - 5$ **e** $x^n - 2$

What would be the anti-derivative of nx^{n-1}?

Can you find a general rule for anti-derivatives that would work for these examples?

WS Anti-derivatives

WS Antiderivatives

WS Antidifferentiation

4.07 Anti-derivatives

Since anti-differentiation is the reverse of differentiation, we can find the equation of an anti-derivative function.

Anti-derivative of x^n

$$\text{If } \frac{dy}{dx} = x^n, \text{ then } y = \frac{1}{n+1}x^{n+1} + C \text{ where } C \text{ is a constant.}$$

Proof

$$\frac{d}{dx}\left(\frac{1}{n+1}x^{n+1} + C\right) = \frac{(n+1)x^n}{n+1}$$
$$= x^n$$

ISBN 9780170413220

We can apply the same rules to anti-derivatives as we use for derivatives. Here are some of the main ones we use.

Anti-derivative rules

If $\frac{dy}{dx} = k$ then $y = kx$.

If $\frac{dy}{dx} = kx^n$ then $y = \frac{1}{n+1} kx^{n+1} + C$.

If $\frac{dy}{dx} = f(x) + g(x)$ then $y = F(x) + G(x) + C$ where $F(x)$ and $G(x)$ are the anti-derivatives of $f(x)$ and $g(x)$ respectively.



iStock.com/hmICA

EXAMPLE 15

Find the anti-derivative of $x^4 - 4x^3 + 9x^2 - 6x + 5$.

Solution

If $f(x) = x^4 - 4x^3 + 9x^2 - 6x + 5$

$$F(x) = \frac{1}{5}x^5 - 4 \times \frac{1}{4}x^4 + 9 \times \frac{1}{3}x^3 - 6 \times \frac{1}{2}x^2 + 5x + C$$

Anti-derivative of 5 is 5x.

$$= \frac{x^5}{5} - x^4 + 3x^3 - 3x^2 + 5x + C$$

If we have some information about the anti-derivative function, we can use this to evaluate the constant C.

The anti-derivative function

EXAMPLE 16

a The gradient of a curve is given by $\frac{dy}{dx} = 6x^2 + 8x$. If the curve passes through the point $(1, -3)$, find its equation.

b If $f''(x) = 6x + 2$ and $f'(1) = f(-2) = 0$, find $f(3)$.

Solution

a $\frac{dy}{dx} = 6x^2 + 8x$

So $y = 6 \times \frac{1}{3}x^3 + 8 \times \frac{1}{2}x^2 + C$

$= 2x^3 + 4x^2 + C$

Substitute $(1, -3)$:

$-3 = 2(1)^3 + 4(1)^2 + C$

$= 6 + C$

$-9 = C$

Equation is $y = 2x^3 + 4x^2 - 9$.

b $f''(x) = 6x + 2$

$f'(x) = 6 \times \frac{1}{2}x^2 + 2 \times \frac{1}{1}x^1 + C$

$= 3x^2 + 2x + C$

Since $f'(1) = 0$:

$0 = 3(1)^2 + 2(1) + C$

$= 5 + C$

$-5 = C$

So $f'(x) = 3x^2 + 2x - 5$

$f(x) = 3 \times \frac{1}{3}x^3 + 2 \times \frac{1}{2}x^2 - 5 \times \frac{1}{1}x^1 + D$

$= x^3 + x^2 - 5x + D$

Since $f(-2) = 0$:

$0 = (-2)^3 + (-2)^2 - 5(-2) + D$

$= -8 + 4 + 10 + D$

$= 6 + D$

$-6 = D$

Equation is $f(x) = x^3 + x^2 - 5x - 6$

$f(3) = 3^3 + 3^2 - 5(3) - 6$

$= 27 + 9 - 15 - 6$

$= 15$

Chain rule

If $\frac{dy}{dx} = (ax + b)^n$, then $y = \frac{1}{a(n+1)}(ax + b)^{n+1} + C$ where C is a constant, $a \neq 0$ and $n \neq -1$.

ISBN 9780170413220

Proof

$$\frac{d}{dx}\left(\frac{1}{a(n+1)}(ax+b)^{n+1}+C\right)=\frac{a(n+1)(ax+b)^{n}}{a(n+1)}$$

$$=(ax+b)^{n}$$

EXAMPLE 17

a Find the anti-derivative of $(3x+7)^8$.

b The gradient of a curve is given by $\frac{dy}{dx}=(2x-3)^4$. If the curve passes through the point $(2,-7)$, find its equation.

Solution

a $\frac{dy}{dx}=(3x+7)^8$

$$y=\frac{1}{a(n+1)}(ax+b)^{n+1}+C$$

$$=\frac{1}{3(8+1)}(3x+7)^{8+1}+C$$

$$=\frac{1}{27}(3x+7)^{9}+C$$

$$=\frac{(3x+7)^{9}}{27}+C$$

b $\frac{dy}{dx}=(2x-3)^4$

$$y=\frac{1}{a(n+1)}(ax+b)^{n+1}+C$$

$$=\frac{1}{2(4+1)}(2x-3)^{4+1}+C$$

$$=\frac{1}{10}(2x-3)^{5}+C$$

Substitute $(2,-7)$:

$$-7=\frac{1}{10}(2\times 2-3)^{5}+C$$

$$=\frac{1}{10}(1)^{5}+C$$

$$=\frac{1}{10}+C$$

$$-7\frac{1}{10}=C$$

So the equation is $y=\frac{1}{10}(2x-3)^{5}-7\frac{1}{10}$

$$=\frac{(2x-3)^{5}-71}{10}$$

ISBN 9780170413220

General chain rule

If $\frac{dy}{dx} = f'(x)[f(x)]^n$ then $y = \frac{1}{n+1}[f(x)]^{n+1} + C$ where C is a constant and $n \neq -1$

Proof

$$\frac{d}{dx}\left(\frac{1}{n+1}[f(x)]^{n+1} + C\right) = \frac{1}{n+1} f'(x)(n+1)[f(x)]^{n+1-1}$$
$$= f'(x)[f(x)]^n$$

EXAMPLE 18

Find the anti-derivative of:

a $8x^3(2x^4 - 1)^5$ **b** $x^2(x^3 + 2)^7$

Solution

a Given $f(x) = 2x^4 - 1$

$$f'(x) = 8x^3$$
$$\frac{dy}{dx} = 8x^3(2x^4 - 1)^5$$
$$= f'(x)[f(x)]^n$$
$$y = \frac{1}{n+1} f(x)^{n+1} + C$$
$$= \frac{1}{5+1}(2x^4 - 1)^{5+1} + C$$
$$= \frac{1}{6}(2x^4 - 1)^6 + C$$
$$= \frac{(2x^4 - 1)^6}{6} + C$$

b Given $f(x) = x^3 + 2$

$$f'(x) = 3x^2$$
$$\frac{dy}{dx} = x^2(x^3 + 2)^7$$
$$= \frac{1}{3} \times 3x^2(x^3 + 2)^7$$
$$= \frac{1}{3} f'(x)[f(x)]^n$$

$$y = \frac{1}{3} \times \frac{1}{n+1} f(x)^{n+1} + C$$
$$= \frac{1}{3} \times \frac{1}{7+1}(x^3 + 2)^{7+1} + C$$
$$= \frac{1}{24}(x^3 + 2)^8 + C$$
$$= \frac{(x^3 + 2)^8}{24} + C$$

ISBN 9780170413220

Exercise 4.07 Anti-derivatives

1 Find the anti-derivative of:

a $2x - 3$ **b** $x^2 + 8x + 1$ **c** $x^5 - 4x^3$

d $(x - 1)^2$ **e** 6 **f** $(3x + 2)^5$

g $8(2x - 7)^4$

2 Find $f(x)$ if:

a $f'(x) = 6x^2 - x$ **b** $f'(x) = x^4 - 3x^2 + 7$ **c** $f'(x) = x - 2$

d $f'(x) = (x + 1)(x - 3)$ **e** $f'(x) = x^{\frac{1}{2}}$

3 Express y in terms of x if:

a $\frac{dy}{dx} = 5x^4 - 9$ **b** $\frac{dy}{dx} = x^{-4} - 2x^{-2}$ **c** $\frac{dy}{dx} = \frac{x^3}{5} - x^2$

d $\frac{dy}{dx} = \frac{2}{x^2}$ **e** $\frac{dy}{dx} = x^3 - \frac{2x}{3} + 1$

4 Find the anti-derivative of:

a $\sqrt{x}$ **b** x^{-3} **c** $\frac{1}{x^8}$

d $x^{-\frac{1}{2}} + 2x^{-\frac{2}{3}}$ **e** $x^{-7} - 2x^{-2}$

5 Find the anti-derivative of:

a $2x(x^2 + 5)^4$ **b** $3x^2(x^3 - 1)^9$ **c** $8x(2x^2 + 3)^3$

d $15x^4(x^5 + 1)^6$ **e** $x(x^2 - 4)^7$ **f** $x^5(2x^6 - 7)^8$

g $(2x - 1)(x^2 - x + 3)^4$ **h** $(3x^2 + 4x - 7)(x^3 + 2x^2 - 7x)^{10}$

i $(x - 3)(x^2 - 6x - 1)^5$

6 If $\frac{dy}{dx} = x^3 - 3x^2 + 5$ and $y = 4$ when $x = 1$, find an equation for y in terms of x.

7 If $f'(x) = 4x - 7$ and $f(2) = 5$, find an equation for $y = f(x)$.

8 Given $f'(x) = 3x^2 + 4x - 2$ and $f(-3) = 4$, find the value of $f(1)$.

9 Given that the gradient of the tangent to a curve is given by $\frac{dy}{dx} = 2 - 6x$ and the curve passes through $(-2, 3)$, find the equation of the curve.

10 If $\frac{dx}{dt} = (t - 3)^2$ and $x = 7$ when $t = 0$, find x when $t = 4$.

11 Given $\frac{d^2y}{dx^2} = 8$, and $\frac{dy}{dx} = 0$ and $y = 3$ when $x = 1$, find the equation of y in terms of x.

12 If $\frac{d^2y}{dx^2} = 12x + 6$ and $\frac{dy}{dx} = 1$ at the point $(-1, -2)$, find the equation of the curve.

ISBN 9780170413220

13 If $f''(x) = 6x - 2$ and $f'(2) = f(2) = 7$, find the equation of the function $y = f(x)$.

14 Given $f''(x) = 5x^4$, $f'(0) = 3$ and $f(-1) = 1$, find $f(2)$.

15 A curve has $\frac{d^2y}{dx^2} = 8x$ and the tangent at $(-2, 5)$ has an angle of inclination of 45° with the x-axis. Find the equation of the curve.

16 The tangent to a curve with $\frac{d^2y}{dx^2} = 2x - 4$ makes an angle of inclination of 135° with the x-axis at the point $(2, -4)$. Find its equation.

17 A function has a tangent parallel to the line $4x - y - 2 = 0$ at the point $(0, -2)$, and $f''(x) = 12x^2 - 6x + 4$. Find the equation of the function.

18 A curve has $\frac{d^2y}{dx^2} = 6$ and the tangent at $(-1, 3)$ is perpendicular to the line $2x + 4y - 3 = 0$. Find the equation of the curve.

19 A function has $f'(1) = 3$ and $f(1) = 5$. Evaluate $f(-2)$ given $f''(x) = 6x + 18$.

20 The velocity of an object is given by $\frac{dx}{dt} = 6t - 5$. If the object has initial displacement of -2, find the equation for the displacement.

21 The acceleration of a particle is given by $\frac{d^2x}{dt^2} = 24t^2 - 12t + 6$ m s^{-2}. Its velocity $\frac{dx}{dt} = 0$ when $t = 1$ and its displacement $x = -3$ when $t = 0$. Find the equation for its displacement.

4.08 Further anti-derivatives

Anti-derivative of exponential functions

$$\text{If } \frac{dy}{dx} = e^x, \text{ then } y = e^x + C$$

Chain rule

$$\text{If } \frac{dy}{dx} = e^{ax+b}, \text{ then } y = \frac{1}{a}e^{ax+b} + C$$

$$\text{If } \frac{dy}{dx} = f'(x)e^{f(x)}, \text{ then } y = e^{f(x)} + C$$

Proof (by differentiation)

$$\frac{d}{dx}\left(\frac{1}{a}e^{ax+b} + C\right) = \frac{1}{a} \times ae^{ax+b}$$
$$= e^{ax+b}$$

$$\frac{d}{dx}[e^{f(x)} + C] = f'(x)e^{f(x)}$$

ISBN 9780170413220

EXAMPLE 19

a Find the anti-derivative of $e^{4x} + 1$.

b Find the equation of the function $y = f(x)$ given $f'(x) = 6e^{3x}$ and $f(2) = 2e^6$.

Solution

a $\frac{1}{a} e^{ax+b} + C = \frac{1}{4} e^{4x} + C$

b $f'(x) = 6e^{3x}$

$f(x) = 6 \times \frac{1}{3} e^{3x} + C$

$= 2e^{3x} + C$

If $f(2) = 2e^6$:

$2e^6 = 2e^{3 \times 2} + C$

$= 2e^6 + C$

$0 = C$

So $f(x) = 2e^{3x}$

Anti-derivative of $\frac{1}{x}$

$$\text{If } \frac{dy}{dx} = \frac{1}{x}, \text{ then } y = \ln|x| + C$$

Chain rule

$$\text{If } \frac{dy}{dx} = \frac{f'(x)}{f(x)}, \text{ then } y = \ln|f(x)| + C$$

Proof

$\frac{d}{dx}(\ln x) = \frac{1}{x}$ for $x > 0$, because $\ln x$ is defined only for $x > 0$.

So the anti-derivative of $\frac{1}{x}$ when $x > 0$ is $\ln x$.

Suppose $x < 0$.

Then $\ln(-x)$ is defined because $-x$ is positive.

ISBN 9780170413220

$$\frac{d}{dx}\left[\ln(-x)\right] = \frac{f'(x)}{f(x)}$$

$$= \frac{-1}{-x}$$

$$= \frac{1}{x}, \qquad x < 0$$

So if $\frac{dy}{dx} = \frac{1}{x}$, then $y = \begin{cases} \ln x + C & \text{if } x > 0 \\ \ln(-x) + C & \text{if } x < 0 \end{cases}$,

or more simply, $y = \ln|x| + C$

EXAMPLE 20

a Find the anti-derivative of $\frac{3}{x}$.

b Find the equation of the function that has $\frac{dy}{dx} = \frac{6x}{x^2 - 5}$ and passes through $(3, 3 \ln 4)$.

Solution

a $\frac{dy}{dx} = \frac{3}{x}$

$= 3 \times \frac{1}{x}$

$y = 3 \ln|x|$

b $\frac{dy}{dx} = \frac{6x}{x^2 - 5}$

$= 3 \times \frac{2x}{x^2 - 5}$

$= 3 \times \frac{f'(x)}{f(x)}$ where $f(x) = x^2 - 5$

$y = 3 \ln f|x| + C$

$= 3 \ln|x^2 - 5| + C$

Substitute $(3, 3 \ln 4)$:

$3 \ln 4 = 3 \ln|3^2 - 5| + C$

$= 3 \ln 4 + C$

$0 = C$

So $y = 3 \ln|x^2 - 5|$

ISBN 9780170413220

Anti-derivatives of trigonometric functions

If $\frac{dy}{dx} = \cos x$, then $y = \sin x + C$ since $\frac{d}{dx}(\sin x) = \cos x$

If $\frac{dy}{dx} = \sin x$, then $y = -\cos x + C$ since $\frac{d}{dx}(\cos x) = -\sin x$ so $\frac{d}{dx}(-\cos x) = \sin x$

If $\frac{dy}{dx} = \sec^2 x$, then $y = \tan x + C$ since $\frac{d}{dx}(\tan x) = \sec^2 x$

Chain rule

If $\frac{dy}{dx} = \cos(ax + b)$, then $y = \frac{1}{a}\sin(ax + b) + C$

If $\frac{dy}{dx} = \sin(ax + b)$, then $y = -\frac{1}{a}\cos(ax + b) + C$

If $\frac{dy}{dx} = \sec^2(ax + b)$, then $y = \frac{1}{a}\tan(ax + b) + C$

If $\frac{dy}{dx} = f'(x)\cos f(x)$, then $y = \sin f(x) + C$

If $\frac{dy}{dx} = f'(x)\sin f(x)$, then $y = -\cos f(x) + C$

If $\frac{dy}{dx} = f'(x)\sec^2 f(x)$, then $y = \tan f(x) + C$

Proof

$$\frac{d}{dx}\left[\frac{1}{a}\sin(ax+b)+C\right] = \frac{1}{a} \times a\cos(ax+b)$$
$$= \cos(ax + b)$$

The other results can be proved similarly.

EXAMPLE 21

a Find the anti-derivative of $\cos 3x$.

b Find the equation of the curve that passes through $\left(\frac{\pi}{4}, 3\right)$ and has $\frac{dy}{dx} = \sec^2 x$.

Solution

a $y = \frac{1}{a}\sin(ax + b) + C$

$= \frac{1}{3}\sin 3x + C$

ISBN 9780170413220

b $y = \tan x + C$

Substitute $\left(\frac{\pi}{4}, 3\right)$:

$3 = \tan\frac{\pi}{4} + C$

$= 1 + C$

$2 = C$

So $y = \tan x + 2$

Exercise 4.08 Further anti-derivatives

1 Find the anti-derivative of:

a $\sin x$ **b** $\sec^2 x$ **c** $\cos x$

d $\sec^2 7x$ **e** $\sin(2x - \pi)$

2 Anti-differentiate:

a e^x **b** e^{6x} **c** $\frac{1}{x}$

d $\frac{3}{3x-1}$ **e** $\frac{x}{x^2+5}$

3 Find the anti-derivative of:

a $e^x + 5$ **b** $\cos x + 4x$ **c** $x + \frac{1}{x}$

d $8x^3 - 3x^2 + 6x - 3 + x^{-1}$ **e** $\sin 5x - \sec^2 9x$

4 Find the equation of a function with $\frac{dy}{dx} = \cos x$ and passing through $\left(\frac{\pi}{2}, -4\right)$.

5 Find the equation of the function that has $f'(x) = \frac{5}{x}$ and $f(1) = 3$.

6 A function has $\frac{dy}{dx} = 4\cos 2x$ and passes through the point $\left(\frac{\pi}{6}, 2\sqrt{3}\right)$.
Find the exact equation of the function.

7 A curve has $f''(x) = 27e^{3x}$ and has $f(2) = f'(2) = e^6$. Find the equation of the curve.

8 The rate of change of a population over time t years is given by $\frac{dP}{dt} = 1350e^{0.054t}$.
If the initial population is 35 000, find:

a the equation for population **b** the population after 10 years

9 The velocity of a particle is given by $\frac{dx}{dt} = 3e^{3t}$ and the particle has an initial displacement of 5 metres. Find the equation for displacement of the particle.

ISBN 9780170413220

10 A pendulum has acceleration given by $\frac{d^2x}{dt^2} = -9 \sin 3t$, initial displacement 0 cm and initial velocity 3 cm s^{-1}.

a Find the equation for its velocity.

b Find the displacement after 2 seconds.

c Find the times when the pendulum has displacement 0 cm.

Summary of differentiation rules

Rule	Chain rule
$\frac{d}{dx}(x^n) = nx^{n-1}$	$\frac{d}{dx}[f(x)]^n = f'(x)n[f(x)]^{n-1}$
$\frac{d}{dx}(e^x) = e^x$	$\frac{d}{dx}[e^{f(x)}] = f'(x)e^{f(x)}$
$\frac{d}{dx}(\ln x) = \frac{1}{x}$	$\frac{d}{dx}[\ln f(x)] = \frac{f'(x)}{f(x)}$
$\frac{d}{dx}(\sin x) = \cos x$	$\frac{d}{dx}[\sin f(x)] = f'(x)\cos f(x)$
$\frac{d}{dx}(\cos x) = -\sin x$	$\frac{d}{dx}[\cos f(x)] = -f'(x)\sin f(x)$
$\frac{d}{dx}(\tan x) = \sec^2 x$	$\frac{d}{dx}[\tan f(x)] = f'(x)\sec^2 f(x)$

Product rule: $\frac{d}{dx}(uv) = u'v + v'u$

Quotient rule: $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{u'v - v'u}{v^2}$



Shutterstock.com/Shairith

ISBN 9780170413220

4. TEST YOURSELF

Practice quiz

For Questions **1** to **4**, choose the correct answer **A**, **B**, **C** or **D**.

1 The anti-derivative of $\sin 6x$ is:

A $\frac{1}{6}\cos 6x$ **B** $6\cos 6x$ **C** $-6\cos 6x$ **D** $-\frac{1}{6}\cos 6x$


2 The gradient of the tangent to the curve $y = e^{2x} + x$ at $(0, 1)$ is:

A 2 **B** 3 **C** e **D** 1


3 If $y = \cos 2x$, then $\frac{d^2y}{dx^2}$ is :

A $-4y$ **B** $-2y$ **C** $4y$ **D** y


4 The graph of the derivative $y = f'(x)$ is shown. Which of the following graphs could be the graph of $y = f(x)$?




A




B



C



D



5 Differentiate:

a e^{5x} **b** $2e^{1-x}$ **c** $\ln 4x$ **d** $\ln(4x + 5)$

e xe^x **f** $\frac{\ln x}{x}$ **g** $(e^x + 1)^{10}$

6 Differentiate:

a $\cos x$ **b** $2\sin x$ **c** $\tan x + 1$ **d** $x\sin x$

e $\frac{\tan x}{x}$ **f** $\cos 3x$ **g** $\tan 5x$

ISBN 9780170413220

7 Find the equation of the tangent to the curve $y = 2 + e^{3x}$ at the point where $x = 0$.

8 Find the equation of the tangent to the curve $y = \sin 3x$ at the point $\left(\frac{\pi}{4}, \frac{1}{\sqrt{2}}\right)$.

9 If $x = \cos 2t$, show that $\frac{d^2x}{dt^2} = -4x$.

10 Find the exact gradient of the normal to the curve $y = x - e^{-x}$ at the point where $x = 2$.

11 Find the anti-derivative of:


a $10x^4 - 4x^3 + 6x - 3$ **b** e^{5x} **c** $\sec^2 9x$

d $\frac{1}{x+5}$ **e** $\cos 2x$ **f** $\sin\left(\frac{x}{4}\right)$

12 Find the gradient of the tangent to the curve $y = 3 \cos 2x$ at the point where $x = \frac{\pi}{6}$.

13 A curve has $\frac{dy}{dx} = 6x^2 + 12x - 5$. If the curve passes through the point $(2, -3)$, find the equation of the curve.

14 Sketch the graph of the anti-derivative of the following function, given that the anti-derivative passes through $(0, 4)$.



15 Find the equation of the normal to the curve $y = \ln x$ at the point $(2, \ln 2)$.

16 Find the equation of the normal to the curve $y = \tan x$ at the point $\left(\frac{\pi}{4}, 1\right)$.

17 Differentiate:

a $(5x^2 + 7)^4$ **b** $4x(2x - 3)^7$ **c** $\frac{5x-1}{3x+4}$

d $2x^3e^x$ **e** $\frac{\tan 3x}{x+1}$

18 If $f''(x) = 15x + 12$ and $f(2) = f'(2) = 5$, find the equation of $y = f(x)$.


19 If $f(x) = 3x^5 - 2x^4 + x^3 - 2$, find:

a $f(-1)$ **b** $f'(-1)$ **c** $f''(-1)$


ISBN 9780170413220

20 Sketch an example of the graph of an anti-derivative function for each graph.


a



b



c



21 A function has $f'(3) = 5$ and $f(3) = 2$. If $f''(x) = 12x - 6$, find the equation of the function.

22 Find the anti-derivative of:

a $x^3(3x^4 - 5)^6$ **b** $3x(x^2 + 1)^9$

ISBN 9780170413220

4. CHALLENGE EXERCISE

1 Find the exact gradient of the tangent to the curve $y = e^{x + \ln x}$ at the point where $x = 1$.

2 Find the first and second derivatives of $\dfrac{5 - x}{(4x^2 + 1)^3}$.

3 Find the anti-derivative of:

a $2xe^{x^2}$ **b** $x^2 \sin(x^3)$

4 Differentiate $e^{x \sin 2x}$.

5 A curve passes through the point $(0, -1)$ and the gradient at any point is given by $(x + 3)(x - 5)$. Find the equation of the curve.

6 The rate of change of V with respect to t is given by $\dfrac{dV}{dt} = (2t - 1)^2$.

If $V = 5$ when $t = \dfrac{1}{2}$, find V when $t = 3$.

7 Find the derivative of $y = \dfrac{x \log_e x}{e^x}$.

8 **a** Differentiate $\ln(\tan x)$.

b Find the anti-derivative of $\tan x$.

9 Find the anti-derivative of:

a $x^2 \sin(x^3 - \pi)$ **b** xe^{x^2}

ISBN 9780170413220