Solutions to Exercise 10A

1a Initially, $x = y = 0$, and, $\dot{x} = 30\sqrt{2}\cos 45°$ $\dot{y} = 30\sqrt{2}\sin 45°$

$\dot{x} = 30$ $\dot{y} = 30$

To begin, $\ddot{x} = 0$ (1) To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{x} = C_1$ Integrating, $\dot{y} = -10t + C_2$

When $t = 0$, $\dot{x} = 30$ When $t = 0$, $\dot{y} = 30$

$30 = C_1$ $30 = C_2$

so $\dot{x} = 30$ (2) so $\dot{y} = -10t + 30$ (4)

1b To begin, $\dot{x} = 30$ To begin, $\dot{y} = -10t + 30$

Integrating, $x = 30t + C_3$ Integrating, $y = -5t^2 + 30t + C_4$

When $t = 0$, $x = 0$ When $t = 0$, $y = 0$

$0 = C_3$ $0 = C_4$

so $x = 30t$ (5) So $y = -5t^2 + 30t$ (6)

1c When $y = 0$,

From (6):

$0 = -5t^2 + 30t$

$5t^2 - 30t = 0$

$5t(t - 6) = 0$

$t = 0$ or $t = 6$

Particle returns to the ground at $t > 0$ seconds, so at $t = 6$ seconds.

1d When $t = 6$,

From (5):

$x = 30 \times 6$

$= 180$ m

The horizontal distance travelled by the particle is 180 m.

© Cambridge University Press 2019

1e When $\dot{y} = 0$,

From (4):

$0 = -10t + 30$

$10t = 30$

$t = 3$ seconds

The particle reaches its greatest height after 3 seconds.

1f When $t = 3$,

From (6):

$y = -45 + 90$

$= 45$ m

The greatest height of the particle is 45 m.

2a Initially, $x = y = 0$, and, $\dot{x} = 40\cos 30°$ $\dot{y} = 40\sin 30°$

$\dot{x} = 20\sqrt{3}$ $\dot{y} = 20$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = 20\sqrt{3}$

$20\sqrt{3} = C_1$

so $\dot{x} = 20\sqrt{3}$ (2)

To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{y} = -10t + C_2$

When $t = 0$, $\dot{y} = 20$

$20 = C_2$

so $\dot{y} = -10t + 20$ (4)

2b To begin, $\dot{x} = 20\sqrt{3}$

Integrating, $x = 20t\sqrt{3} + C_3$

When $t = 0$, $x = 0$

$0 = C_3$

so $x = 20t\sqrt{3}$ (5)

To begin, $\dot{y} = -10t + 20$

Integrating, $y = -5t^2 + 20t + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + 20t$ (6)

© Cambridge University Press 2019

2c i The particle returns to the ground when $y = 0$.

Substituting into (6):

$0 = -5t^2 + 20t$

$5t^2 - 20t = 0$

$5t(t - 4) = 0$

$t = 0$ or $t = 4$

Since $t > 0$, particle returns to the ground at $t = 4$ seconds.

2c ii From (5):

When $t = 4$,

$x = 20 \times 4 \times \sqrt{3}$

$= 80\sqrt{3}$ m

The horizontal distance travelled by the particle is $80\sqrt{3}$ m.

2c iii Particle reaches greatest height when $\dot{y} = 0$.

Substituting into (4):

$0 = -10t + 20$

$10t = 20$

$t = 2$

Substituting into (6):

$y = -20 + 40$

$= 20$ m

The greatest height reached above the ground is 20 m.

3a Initially, $x = y = 0$, and, $\dot{x} = 20\cos 60°$ $\dot{y} = 20\sin 60°$

$\dot{x} = 10$ $\dot{y} = 10\sqrt{3}$

To begin, $\ddot{x} = 0$ (1) To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{x} = C_1$ Integrating, $\dot{y} = -10t + C_2$

© Cambridge University Press 2019

When $t = 0$, $\dot{x} = 10$

$10 = C_1$

so $\dot{x} = 10$ (2)

When $t = 0$, $\dot{y} = 10\sqrt{3}$

$10\sqrt{3} = C_2$

so $\dot{y} = -10t + 10\sqrt{3}$ (4)

3b To begin, $\dot{x} = 10$

Integrating, $x = 10t + C_3$

When $t = 0$, $x = 0$

$0 = C_3$

so $x = 10t$ (5)

To begin, $\dot{y} = -10t + 10\sqrt{3}$

Integrating, $y = -5t^2 + 10t\sqrt{3} + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + 10t\sqrt{3}$ (6)

3c i Since the particle starts at the origin and moves to a point with horizontal component x and vertical component y, we can consider a right-angled triangle whose hypotenuse equals the distance travelled.

Pythagoras' theorem can be used to calculate this hypotenuse.

$r^2 = x^2 + y^2$

$r = \sqrt{x^2 + y^2}$

From (5) and (6):

$$r = \sqrt{(10t)^2 + \left(-5t^2 + 10t\sqrt{3}\right)^2}$$

When $t = 1$,

$$r = \sqrt{10^2 + \left(-5 + 10\sqrt{3}\right)^2}$$

$$= \sqrt{100 + 25 - 100\sqrt{3} + 300}$$

$$= \sqrt{425 - 100\sqrt{3}}$$

$\div 15.9$ m

So the distance of the particle from the origin after one second is about 15.9 m.

© Cambridge University Press 2019

3c ii The particle's speed can be calculated by applying Pythagoras' theorem to the particle's velocity vector $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$.

$v^2 = \dot{x}^2 + \dot{y}^2$

$v = \sqrt{\dot{x}^2 + \dot{y}^2}$

From (2) and (4):

$v = \sqrt{10^2 + \left(-10t + 10\sqrt{3}\right)^2}$

When $t = 1$,

$v = \sqrt{10^2 + \left(-10 + 10\sqrt{3}\right)^2}$

$= \sqrt{100 + 100 - 200\sqrt{3} + 300}$

$= \sqrt{500 - 200\sqrt{3}}$

$\div 12.4$ m/s

The speed of the particle after one second is about 12.4 m/s.

4a Initially, $x = y = 0$, and, $\dot{x} = 60\cos\left(\tan^{-1}\frac{4}{3}\right)$ $\dot{y} = 60\sin\left(\tan^{-1}\frac{4}{3}\right)$

$\dot{x} = 36$ $\dot{y} = 48$

To begin,	$\ddot{x} = 0$	(1)		To begin,	$\ddot{y} = -10$	(3)
Integrating,	$\dot{x} = C_1$			Integrating,	$\dot{y} = -10t + C_2$	
When $t = 0$,	$\dot{x} = 36$			When $t = 0$,	$\dot{y} = 48$	
	$36 = C_1$				$48 = C_2$	
so	$\dot{x} = 36$	(2)		so 3	$\dot{y} = -10t + 48$	(4)

4b

To begin,	$\dot{x} = 36$			To begin,	$\dot{y} = -10t + 48$	
Integrating,	$x = 36t + C_3$			Integrating,	$y = -5t^2 + 48t + C_4$	
When $t = 0$,	$x = 0$			When $t = 0$,	$y = 0$	
	$0 = C_3$				$0 = C_4$	
so	$x = 36t$	(5)		so	$y = -5t^2 + 48t$	(6)

© Cambridge University Press 2019

4c $d^2 = x^2 + y^2$

$d = \sqrt{x^2 + y^2}$

From (5) and (6):

$d = \sqrt{(36t)^2 + (-5t^2 + 48t)^2}$

When $t = 3$,

$d = \sqrt{108^2 + (-45 + 144)^2}$

$= \sqrt{21\,465}$

$\div 146.5$ m

The distance of the particle, from the point of projection, after 3 seconds is about 146.5 m.

4d $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

From (2) and (4):

$\underline{v} = 36\underline{i} + (-10t + 48)\underline{j}$

When $t = 3$,

$\underline{v} = 36\underline{i} + 18\underline{j}$

$v^2 = 36^2 + 18^2$

$v = \sqrt{36^2 + 18^2}$

$= \sqrt{1296 + 324}$

$= \sqrt{1620}$

$= \sqrt{324 \times 5}$

$= 18\sqrt{5}$ m/s

$\tan\theta = \frac{18}{36} = \frac{1}{2}$

$\theta = \tan^{-1}\frac{1}{2}$

© Cambridge University Press 2019

So, the velocity of the particle after three seconds is $18\sqrt{5}$ m/s at an angle of $\tan^{-1}\frac{1}{2}$ above the horizontal.

5a Initially, $x = y = 0$, and, $\dot{x} = 8$ $\dot{y} = 6$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = 8$

$8 = C_1$

so $\dot{x} = 8$ (2)

To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{y} = -10t + C_2$

When $t = 0$, $\dot{y} = 6$

$6 = C_2$

so $\dot{y} = -10t + 6$ (4)

Now $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

so $\underline{v} = 8\underline{i} + (-10t + 6)\underline{j}$

$\underline{v} = 8\underline{i} + (6 - 10t)\underline{j}$

5b To begin, $\dot{x} = 8$

Integrating, $x = 8t + C_3$

When $t = 0$, $x = 0$

$0 = C_3$

so $x = 8t$ (5)

To begin, $\dot{y} = -10t + 6$

Integrating, $y = -5t^2 + 6t + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + 6t$ (6)

$\underline{r} = x\underline{i} + y\underline{j}$

so $\underline{r} = 8t\underline{i} + (-5t^2 + 6t)\underline{j}$

$\underline{r} = (8t)\underline{i} + (6t - 5t^2)\underline{j}$

5c i $\underline{v} = 8\underline{i} + 6\underline{j}$

$v^2 = 8^2 + 6^2$

$v = \sqrt{8^2 + 6^2}$

$= \sqrt{64 + 36}$

© Cambridge University Press 2019

$= \sqrt{100}$

$= 10$ m/s

The initial speed of the particle is 10 m/s.

5c ii $\underline{r} = (8t)\underline{i} + (6t - 5t^2)\underline{j}$

When $t = 2$,

$\underline{r} = 16\underline{i} + (12 - 20)\underline{j}$

$= 16\underline{i} - 8\underline{j}$

5c iii From (4):

When $\dot{y} = 0$,

$0 = -10t + 6$

$10t = 6$

$t = 0.6$ seconds

When $t = 0.6$,

$\underline{r} = 4.8\underline{i} + (3.6 - 1.8)\underline{j}$

$= 4.8\underline{i} + 1.8\underline{j}$

6a To begin, $x = 40t$ (1)

Differentiate, $\dot{x} = 40$ (2)

When $t = 0$, $\dot{x} = 40$

To begin, $y = -5t^2 + 25t$ (3)

Differentiate, $\dot{y} = -10t + 25$ (4)

When $t = 0$, $\dot{y} = 25$

6b From (2) and (4):

$\underline{v} = 40\underline{i} + 25\underline{j}$

$v^2 = 40^2 + 25^2$

$v = \sqrt{40^2 + 25^2}$

$= \sqrt{1600 + 625}$

© Cambridge University Press 2019

$= \sqrt{2225}$

$= 5\sqrt{89}$ m/s

$\tan\theta = \frac{25}{40} = \frac{5}{8}$

$\theta = \tan^{-1}\frac{5}{8}$

$\theta \div 32°$

7a Initially, $x = y = 0$, and, $\dot{x} = V\cos\alpha \quad \dot{y} = V\sin\alpha$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = V\cos\alpha$

$V\cos\alpha = C_1$

so $\dot{x} = V\cos\alpha$ (2a)

Integrating, $x = Vt\cos\alpha + C_2$

When $t = 0$, $x = 0$

$0 = C_2$

so $x = Vt\cos\alpha$ (3a)

When $t = 2$, $8 = 2V\cos\alpha$

so $V\cos\alpha = 4$

Hence, $\underline{v} = 4\underline{i} + 4\underline{j}$

From (2a): $\dot{x} = 4$ (2b)

From (3a): $x = 4t$ (3b)

To begin, $\ddot{y} = -10$ (4)

Integrating, $\dot{y} = -10t + C_3$

When $t = 0$, $\dot{y} = V\sin\alpha$

$V\sin\alpha = C_3$

so $\dot{y} = -10t + V\sin\alpha$ (5a)

Integrating, $y = -5t^2 + Vt\sin\alpha + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + Vt\sin\alpha$ (6a)

When $t = 2$, $-12 = -20 + 2V\sin\alpha$

so $V\sin\alpha = 4$

From (5a): $\dot{y} = -10t + 4$ (5b)

From (6a): $y = -5t^2 + 4t$ (6b)

7b $\underline{r} = x\underline{i} + y\underline{j}$

From (3b) and (6b):

$\underline{r} = 4t\underline{i} + (-5t^2 + 4t)\underline{j}$

When $t = 0.5$,

© Cambridge University Press 2019

$\underline{r} = 2\underline{i} + (-1.25 + 2)\underline{j}$

$= 2\underline{i} + 0.75\underline{j}$

8a Initially, $x = y = 0$, and, $\dot{x} = V\cos\theta \quad \dot{y} = V\sin\theta$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = V\cos\theta$		When $t = 0$,	$\dot{y} = V\sin\theta$	
	$V\cos\theta = C_1$			$V\sin\theta = C_3$	
so	$\dot{x} = V\cos\theta$	(2a)	so	$\dot{y} = -10t + V\sin\theta$	(5a)
Integrating,	$x = Vt\cos\theta + C_2$		Integrating,	$y = -5t^2 + Vt\sin\theta + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x = Vt\cos\theta$	(3a)	so	$y = -5t^2 + Vt\sin\theta$	(6a)
When $t = 2$,	$30 = 2V\cos\theta$		When $t = 2$,	$30 = -20 + 2V\sin\theta$	
so	$V\cos\theta = 15$		so	$V\sin\theta = 25$	
From (2a):	$\dot{x} = 15$	(2b)	From (5a):	$\dot{y} = -10t + 25$	(5b)
From (3a):	$x = 15t$	(3b)	From (6a):	$y = -5t^2 + 25t$	(6b)

8b $\underline{V} = V\cos\theta\,\underline{i} + V\sin\theta\,\underline{j}$

So $\underline{V} = 15\underline{i} + 25\underline{j}$

$V^2 = 15^2 + 25^2$

$V = \sqrt{15^2 + 25^2}$

$= \sqrt{225 + 625}$

$= \sqrt{850}$

$= 5\sqrt{34}$ m/s

$\tan\theta = \frac{25}{15} = \frac{5}{3}$

© Cambridge University Press 2019

$\theta = \tan^{-1}\frac{5}{3}$

9a Initially, $x = y = 0$, and, $\dot{x} = 12\cos(-30°)$ $\dot{y} = 12\sin(-30°)$
$\dot{x} = 6\sqrt{3}$ $\dot{y} = -6$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = 6\sqrt{3}$

$6\sqrt{3} = C_1$

so $\dot{x} = 6\sqrt{3}$ (2)

To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{y} = -10t + C_2$

When $t = 0$, $\dot{y} = -6$

$-6 = C_2$

so $\dot{y} = -10t - 6$ (4)

9b To begin, $\dot{x} = 6\sqrt{3}$

Integrating, $x = 6t\sqrt{3} + C_3$

When $t = 0$, $x = 0$

$0 = C_3$

so $x = 6t\sqrt{3}$ (5)

To begin, $\dot{y} = -10t - 6$

Integrating, $y = -5t^2 - 6t + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 - 6t$ (6)

9c The stone will hit the ground when $y = -11$.

Substituting into (6):

$-11 = -5t^2 - 6t$

$5t^2 + 6t - 11 = 0$

$(5t + 11)(t - 1) = 0$

$t = -\frac{11}{5}$ or $t = 1$

But $t \geq 0$, so the stone hits the ground when $t = 1$ second.

9d From (5):

$x = 6t\sqrt{3}$

When $t = 1$,

© Cambridge University Press 2019

$x = 6\sqrt{3}$

$\div 10.4$ m

The stone hits the ground about 10.4 m from the base of the tower.

9e $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

From (2) and (4):

$\underline{v} = 6t\sqrt{3}\underline{i} + (-10t - 6)\underline{j}$

When $t = 1$,

$\underline{v} = 6\sqrt{3}\underline{i} - 16\underline{j}$

$\tan\theta = -\dfrac{16}{6\sqrt{3}}$

$= -\dfrac{8\sqrt{3}}{9}$

$\theta = \tan^{-1}\left(-\dfrac{8\sqrt{3}}{9}\right)$

So the stone will hit the ground at an angle of $\tan^{-1}\frac{8\sqrt{3}}{9}$ below the horizontal.

10a Initially, $x = y = 0$, and, $\dot{x} = 6\cos 45°$ $\dot{y} = 6\sin 45°$

$\dot{x} = 3\sqrt{2}$ $\dot{y} = 3\sqrt{2}$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = 3\sqrt{2}$

$3\sqrt{2} = C_1$

so $\dot{x} = 3\sqrt{2}$ (2)

To begin, $\dot{x} = 3\sqrt{2}$

Integrating, $x = 3t\sqrt{2} + C_2$

When $t = 0$, $x = 0$

$0 = C_2$

To begin, $\ddot{y} = -10$ (4)

Integrating, $\dot{y} = -10t + C_3$

When $t = 0$, $\dot{y} = 3\sqrt{2}$

$3\sqrt{2} = C_3$

so $\dot{y} = -10t + 3\sqrt{2}$ (5)

To begin, $\dot{y} = -10t + 3\sqrt{2}$

Integrating, $y = -5t^2 + 3t\sqrt{2} + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

© Cambridge University Press 2019

so $x = 3t\sqrt{2}$ (3) so $y = -5t^2 + 3t\sqrt{2}$ (6)

The ball will hit the ground when $y = -2$.

Substituting into (6):

$-2 = -5t^2 + 3t\sqrt{2}$

$5t^2 - 3t\sqrt{2} - 2 = 0$

Using the quadratic formula:

$$t = \frac{3\sqrt{2} \pm \sqrt{\left(-3\sqrt{2}\right)^2 - 4 \times 5 \times (-2)}}{2 \times 5}$$

$$= \frac{3\sqrt{2} \pm \sqrt{18 + 40}}{10}$$

$$= \frac{3\sqrt{2} \pm \sqrt{58}}{10}$$

$t \div -0.34$ or $t \div 1.19$

But $t \geq 0$, so the stone hits the ground when $t \div 1.19$ seconds.

10b From (3):

$x = 3t\sqrt{2}$

When $t = \frac{3\sqrt{2} + \sqrt{58}}{10}$,

$$x = 3 \times \frac{3\sqrt{2} + \sqrt{58}}{10} \times \sqrt{2}$$

$\div 5.03$ m

The horizontal distance travelled by the ball is about 5.03 m.

11 Let V be the initial speed of the projectile

Initially, $x = y = 0$, and, $\dot{x} = V\cos 36°$ $\dot{y} = V\sin 36°$

To begin, $\ddot{x} = 0$ (1) To begin, $\ddot{y} = -10$ (4)

Integrating, $\dot{x} = C_1$ Integrating, $\dot{y} = -10t + C_3$

© Cambridge University Press 2019

When $t = 0$, $\dot{x} = V\cos 36°$

$V\cos 36° = C_1$

so $\dot{x} = V\cos 36°$ (2)

To begin, $\dot{x} = V\cos 36°$

Integrating, $x = Vt\cos 36° + C_2$

When $t = 0$, $x = 0$

$0 = C_2$

so $x = Vt\cos 36°$ (3)

When $t = 0$, $\dot{y} = V\sin 36°$

$V\sin 36° = C_3$

so $\dot{y} = -10t + V\sin 36°$ (5)

To begin, $\dot{y} = -10t + V\sin 36°$

Integrating, $y = -5t^2 + Vt\sin 36° + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + Vt\sin 36°$ (6)

From (3):

$$t = \frac{x}{V\cos 36°}$$

Substituting for t in (6):

$$y = -5\left(\frac{x}{V\cos 36°}\right)^2 + \left(\frac{Vx}{V\cos 36°}\right)\sin 36°$$

$$= \frac{-5x^2}{V^2\cos^2 36°} + x\tan 36°$$

If the projectile just clears the wall, it will reach a point where $x = 20$ and $y = 10$

$$10 = \frac{-5(20)^2}{V^2\cos^2 36°} + 20\tan 36°$$

$$10 = \frac{-2000}{V^2\cos^2 36°} + 20\tan 36°$$

$$10 - 20\tan 36° = \frac{-2000}{V^2\cos^2 36°}$$

$$V^2\cos^2 36° = \frac{-2000}{10 - 20\tan 36°}$$

$$V^2\cos^2 36° = \frac{2000}{20\tan 36° - 10}$$

$$V^2 = \frac{2000}{\cos^2 36°\,(20\tan 36° - 10)}$$

$$V = \sqrt{\frac{2000}{\cos^2 36°\,(20\tan 36° - 10)}}$$

© Cambridge University Press 2019

$V \div 26$ m/s

The initial speed of the projectile is about 26 m/s.

12 Initially, $x = y = 0$, and, $\dot{x} = 20\cos\alpha$ $\dot{y} = 20\sin\alpha$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = 20\cos\alpha$		When $t = 0$,	$\dot{y} = 20\sin\alpha$	
	$20\cos\alpha = C_1$			$20\sin\alpha = C_3$	
so	$\dot{x} = 20\cos\alpha$	(2)	so	$\dot{y} = -10t + 20\sin\alpha$	(5)
To begin,	$\dot{x} = 20\cos\alpha$		To begin,	$\dot{y} = -10t + 20\sin\alpha$	
Integrating,	$x = 20t\cos\alpha + C_2$		Integrating,	$y = -5t^2 + 20t\sin\alpha + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x = 20t\cos\alpha$	(3)	so	$y = -5t^2 + 20t\sin\alpha$	(6)

As the ball reaches its highest point, $\dot{y} = 0$.

Substituting into (5):

$0 = -10t + 20\sin\alpha$

$10t = 20\sin\alpha$

$t = 2\sin\alpha$

Substituting for t in (6):

$y = -5(2\sin\alpha)^2 + 40\sin^2\alpha$

$= -20\sin^2\alpha + 40\sin^2\alpha$

$= 20\sin^2\alpha$

In order to clear the wall at its highest point, $y = 3$ when $\dot{y} = 0$.

$3 = 20\sin^2\alpha$

$\sin^2\alpha = \frac{3}{20}$

© Cambridge University Press 2019

$$\sin\alpha = \sqrt{\frac{3}{20}}$$

$$= \sqrt{\frac{3}{20}} \times \frac{\sqrt{5}}{\sqrt{5}}$$

$$= \frac{\sqrt{15}}{\sqrt{100}}$$

$$= \frac{\sqrt{15}}{10}$$

$$\alpha = \sin^{-1}\frac{\sqrt{15}}{10}$$

13 For P_1

Initially, $x_1 = y_1 = 0$, and, $\dot{x}_1 = 20$ $\dot{y}_1 = 30$

To begin, $\ddot{x}_1 = 0$ (1a)

Integrating, $\dot{x}_1 = C_1$

When $t = 0$, $\dot{x}_1 = 20$

$20 = C_1$

so $\dot{x}_1 = 20$ (2a)

To begin, $\dot{x}_1 = 20$

Integrating, $x_1 = 20t + C_2$

When $t = 0$, $x_1 = 0$

$0 = C_2$

so $x_1 = 20t$ (3a)

To begin, $\ddot{y}_1 = -10$ (4a)

Integrating, $\dot{y}_1 = -10t + C_3$

When $t = 0$, $\dot{y}_1 = 30$

$30 = C_3$

so $\dot{y}_1 = -10t + 30$ (5a)

To begin, $\dot{y}_1 = -10t + 30$

Integrating, $y_1 = -5t^2 + 30t + C_4$

When $t = 0$, $y_1 = 0$

$0 = C_4$

so $y_1 = -5t^2 + 30t$ (6a)

For P_2

Initially, $x_2 = y_2 = 0$, and, $\dot{x}_2 = 60$ $\dot{y}_2 = 50$

To begin, $\ddot{x}_2 = 0$ (1b)

Integrating, $\dot{x}_2 = C_1$

When $t = 0$, $\dot{x}_2 = 60$

$60 = C_1$

To begin, $\ddot{y}_2 = -10$ (4b)

Integrating, $\dot{y}_2 = -10t + C_3$

When $t = 0$, $\dot{y}_2 = 50$

$50 = C_3$

© Cambridge University Press 2019

so	$\dot{x}_2 = 60$	(2b)	so	$\dot{y}_2 = -10t + 50$	(5b)
To begin,	$\dot{x}_2 = 60$		To begin,	$\dot{y}_2 = -10t + 50$	
Integrating,	$x_2 = 60t + C_2$		Integrating,	$y_2 = -5t^2 + 50t + C_4$	
When $t = 0$,	$x_2 = 0$		When $t = 0$,	$y_2 = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x_2 = 60t$	(3b)	so	$y_2 = -5t^2 + 50t$	(6b)

Since P_2 is projected two seconds after P_1,

then $x_2 = 60(t - 2)$

and $y_2 = -5(t - 2)^2 + 50(t - 2)$

If the particles collide, the same value of t must allow $x_1 = x_2$ and $y_1 = y_2$.

From (3a) and (3b):

$20t = 60(t - 2)$

$20t = 60t - 120$

$40t = 120$

$t = 3$

From (6a) and (6b):

$-5t^2 + 30t = -5(t - 2)^2 + 50(t - 2)$

Check when $t = 3$,

LHS $= -45 + 90 = 45$

RHS $= -5 + 50 = 45$

Hence LHS = RHS.

P_1 and P_2 collide at 3 seconds after P_1 is projected (2 seconds after P_2 is projected).

© Cambridge University Press 2019

14a Initially, $x = y = 0$, and, $\dot{x} = 20\sqrt{2}\cos\theta$ $\dot{y} = 20\sqrt{2}\sin\theta$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = 20\sqrt{2}\cos\theta$

$20\sqrt{2}\cos\theta = C_1$

so $\dot{x} = 20\sqrt{2}\cos\theta$ (2)

To begin, $\dot{x} = 20\sqrt{2}\cos\theta$

Integrating, $x = 20t\sqrt{2}\cos\theta + C_2$

When $t = 0$, $x = 0$

$0 = C_2$

so $x = 20t\sqrt{2}\cos\theta$ (3)

To begin, $\ddot{y} = -10$ (4)

Integrating, $\dot{y} = -10t + C_3$

When $t = 0$, $\dot{y} = 20\sqrt{2}\sin\theta$

$20\sqrt{2}\sin\theta = C_3$

so $\dot{y} = -10t + 20\sqrt{2}\sin\theta$ (5)

To begin, $\dot{y} = -10t + 20\sqrt{2}\sin\theta$

Integrating, $y = -5t^2 + 20t\sqrt{2}\sin\theta + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + 20t\sqrt{2}\sin\theta$ (6)

At time t, $x = 20$ and $y = 15$.

Substituting $x = 20$ into (3):

$20 = 20t\sqrt{2}\cos\theta$

so $\sqrt{2}\,t\cos\theta = 1$ (7)

Substituting $y = 15$ into (6):

$15 = -5t^2 + 20t\sqrt{2}\sin\theta$

so $4\sqrt{2}\,t\sin\theta - t^2 = 3$ (8)

14b From (7):

$$t = \frac{1}{\sqrt{2}\cos\theta}$$

Substituting for t in (8):

$$\frac{4\sqrt{2}}{\sqrt{2}\cos\theta}\sin\theta - \left(\frac{1}{\sqrt{2}\cos\theta}\right)^2 = 3$$

$$4\tan\theta - \frac{1}{2\cos^2\theta} = 3$$

© Cambridge University Press 2019

$$4\tan\theta - \frac{1}{2\cos^2\theta} - 3 = 0$$

$$\frac{1}{\cos^2\theta} - 8\tan\theta + 6 = 0$$

$$\sec^2\theta - 8\tan\theta + 6 = 0$$

But $\sec^2\theta = \tan^2\theta + 1$

so $\tan^2\theta - 8\tan\theta + 7 = 0$

14c $\tan^2\theta - 8\tan\theta + 7 = 0$

$(\tan\theta - 1)(\tan\theta - 7) = 0$

$\tan\theta = 1$ or 7

$\theta = 45°$ or $\tan^{-1}7$

$\theta = 45°$ or $\theta \div 81°52'$

15a Initially, $x = y = 0$, and, $\dot{x} = 50\cos\theta$ $\dot{y} = 50\sin\theta$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = 50\cos\theta$		When $t = 0$,	$\dot{y} = 50\sin\theta$	
$50\cos\theta = C_1$			$50\sin\theta = C_3$		
so	$\dot{x} = 50\cos\theta$	(2)	so	$\dot{y} = -10t + 50\sin\theta$	(5)
To begin,	$\dot{x} = 50\cos\theta$		To begin,	$\dot{y} = -10t + 50\sin\theta$	
Integrating,	$x = 50t\cos\theta + C_2$		Integrating,	$y = -5t^2 + 50t\sin\theta + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
$0 = C_2$			$0 = C_4$		
so	$x = 50t\cos\theta$	(3)	so	$y = -5t^2 + 50t\sin\theta$	(6)

At point P, $x = 100$ and $y = 25$.

Substituting $x = 100$ into (3):

$100 = 50t\cos\theta$

© Cambridge University Press 2019

$$t = \frac{2}{\cos\theta} \qquad (7)$$

Substituting $y = 25$ into (6):

$$25 = -5t^2 + 50t\sin\theta$$

$$t^2 - 10t\sin\theta + 5 = 0 \qquad (8)$$

Substituting (7) into (8):

$$\frac{4}{\cos^2\theta} - \frac{20\sin\theta}{\cos\theta} + 5 = 0$$

$$4\sec^2\theta - 20\tan\theta + 5 = 0$$

$$\sec^2\theta - 5\tan\theta + 1.25 = 0$$

But $\sec^2\theta = \tan^2\theta + 1$

$$\tan^2\theta - 5\tan\theta + 2.25 = 0$$

$$\left(\tan\theta - \frac{1}{2}\right)\left(\tan\theta - \frac{9}{2}\right) = 0$$

$$\tan\theta = \frac{1}{2} \text{ or } \frac{9}{2}$$

$$\theta = \tan^{-1}\frac{1}{2} \text{ or } \tan^{-1}\frac{9}{2}$$

15b i

For $\theta = \tan^{-1}\frac{1}{2}$,

consider a right-angled triangle with angle θ and opposite side 1 unit, adjacent side 2 units and hypotenuse $\sqrt{1^2 + 2^2} = \sqrt{5}$ units.

Hence, $\sin\theta = \frac{1}{\sqrt{5}} = \frac{\sqrt{5}}{5}$ and $\cos\theta = \frac{2}{\sqrt{5}} = \frac{2\sqrt{5}}{5}$

Substituting for $\cos\theta$ in (7):

$$t = \frac{2}{\cos\theta}$$

$$= \frac{2}{\left(\frac{2\sqrt{5}}{5}\right)}$$

© Cambridge University Press 2019

$= \dfrac{5}{\sqrt{5}}$

$= \sqrt{5}$ seconds

For $\theta = \tan^{-1}\dfrac{9}{2}$,

consider a right-angled triangle with angle θ and opposite side 9 units, adjacent side 2 units and hypotenuse $\sqrt{9^2 + 2^2} = \sqrt{85}$ units.

Hence, $\sin\theta = \dfrac{9}{\sqrt{85}} = \dfrac{9\sqrt{85}}{85}$ and $\cos\theta = \dfrac{2}{\sqrt{85}} = \dfrac{2\sqrt{85}}{85}$

Substituting for $\cos\theta$ in (7):

$t = \dfrac{2}{\cos\theta}$

$= \dfrac{2}{\left(\dfrac{2\sqrt{85}}{85}\right)}$

$= \dfrac{85}{\sqrt{85}}$

$= \sqrt{85}$ seconds

15b ii To begin, $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

From (2) and (5):

$\underline{v} = 50\cos\theta\,\underline{i} + (-10t + 50\sin\theta)\underline{j}$

When $\theta = \tan^{-1}\dfrac{1}{2}$, $\sin\theta = \dfrac{1}{\sqrt{5}} = \dfrac{\sqrt{5}}{5}$ and $\cos\theta = \dfrac{2}{\sqrt{5}} = \dfrac{2\sqrt{5}}{5}$,

$v^2 = (50\cos\theta)^2 + \left(-10\sqrt{5} + 50\sin\theta\right)^2$

$= \left(50 \times \dfrac{2\sqrt{5}}{5}\right)^2 + \left(-10\sqrt{5} + 50 \times \dfrac{\sqrt{5}}{5}\right)^2$

$= \left(20\sqrt{5}\right)^2 + \left(-10\sqrt{5} + 10\sqrt{5}\right)^2$

$= \left(20\sqrt{5}\right)^2$

© Cambridge University Press 2019

$v = 20\sqrt{5}$

$\doteqdot 44.7 \text{ m/s}$

$\tan\theta = \dfrac{-10\sqrt{5} + 50\sin\theta}{50\cos\theta}$

$= \dfrac{-10\sqrt{5} + 10\sqrt{5}}{20\sqrt{5}}$

$= 0$

So $\theta = 0°$

When $\theta = \tan^{-1}\dfrac{9}{2}$, $\sin\theta = \dfrac{9}{\sqrt{85}} = \dfrac{9\sqrt{85}}{85}$ and $\cos\theta = \dfrac{2}{\sqrt{85}} = \dfrac{2\sqrt{85}}{85}$

$v^2 = (50\cos\theta)^2 + \left(-10\sqrt{85} + 50\sin\theta\right)^2$

$= \left(\dfrac{50 \times 2\sqrt{85}}{85}\right)^2 + \left(-10\sqrt{85} + \dfrac{50 \times 9\sqrt{85}}{85}\right)^2$

$= \left(\dfrac{20\sqrt{85}}{17}\right)^2 + \left(-10\sqrt{85} + \dfrac{90\sqrt{85}}{17}\right)^2$

$= \left(\dfrac{20 \times \sqrt{17} \times \sqrt{5}}{17}\right)^2 + \left(-10 \times \sqrt{17} \times \sqrt{5} + \dfrac{90 \times \sqrt{17} \times \sqrt{5}}{17}\right)^2$

$= \left(\dfrac{20\sqrt{5}}{\sqrt{17}}\right)^2 + \left(\dfrac{-80\sqrt{5}}{\sqrt{17}}\right)^2$

$= \dfrac{2000}{17} + \dfrac{32\,000}{17}$

$= \dfrac{34\,000}{17}$

$= 2000$

$v = \sqrt{2000}$

$= 20\sqrt{5}$

$\doteqdot 44.7 \text{ m/s}$

$\tan\theta = \dfrac{-10\sqrt{85} + 50\sin\theta}{50\cos\theta}$

© Cambridge University Press 2019

$$= \frac{-10\sqrt{85} + \left(\frac{90\sqrt{85}}{17}\right)}{\left(\frac{20\sqrt{85}}{17}\right)}$$

$$= \frac{\frac{-170\sqrt{85}}{17} + \frac{90\sqrt{85}}{17}}{\left(\frac{20\sqrt{85}}{17}\right)}$$

$$= \frac{\frac{-80\sqrt{85}}{17}}{\left(\frac{20\sqrt{85}}{17}\right)}$$

$$= -4$$

So $\theta \div 76.0°$.

For each possible angle of projection, the velocity of the particle is about 44.7 m/s at angles of 0° and 76.0° to the horizontal.

16a i Initially, $x = y = 0$, and, $\dot{x} = V\cos\theta$ $\dot{y} = V\sin\theta$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = V\cos\theta$

$V\cos\theta = C_1$

so $\dot{x} = V\cos\theta$ (2)

To begin, $\dot{x} = V\cos\theta$

Integrating, $x = Vt\cos\theta + C_2$

When $t = 0$, $x = 0$

$0 = C_2$

so $x = Vt\cos\theta$ (3)

To begin, $\ddot{y} = -g$ (4)

Integrating, $\dot{y} = -g + C_3$

When $t = 0$, $\dot{y} = V\sin\theta$

$V\sin\theta = C_3$

so $\dot{y} = -gt + V\sin\theta$ (5)

To begin, $\dot{y} = -gt + V\sin\theta$

Integrating, $y = -\frac{g}{2}t^2 + Vt\sin\theta + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -\frac{g}{2}t^2 + Vt\sin\theta$ (6)

A particle reaches its greatest height when $\dot{y} = 0$.

Substituting into (5):

$0 = -gt + V\sin\theta$

© Cambridge University Press 2019

$gt = V\sin\theta$

$t = \dfrac{V\sin\theta}{g}$

Substituting for t in (6):

$y = -\dfrac{g}{2}\left(\dfrac{V\sin\theta}{g}\right)^2 + \dfrac{V^2\sin^2\theta}{g}$

$= -\dfrac{V^2\sin^2\theta}{2g} + \dfrac{V^2\sin^2\theta}{g}$

$= \dfrac{V^2\sin^2\theta}{2g}$ metres

Hence the greatest height is $\dfrac{V^2\sin^2\theta}{2g}$ metres.

16a ii The particle returns to the ground when $y = 0$ and $t > 0$.

Substituting for y in (6):

$0 = -\dfrac{g}{2}t^2 + Vt\sin\theta$

$t\left(\dfrac{g}{2}t - V\sin\theta\right) = 0$

$t = 0$ or $t = \dfrac{2V\sin\theta}{g}$

But $t > 0$ so $t = \dfrac{2V\sin\theta}{g}$

Substituting for t in (3):

$x = V\left(\dfrac{2V\sin\theta}{g}\right)\cos\theta$

$x = \dfrac{2V^2\sin\theta\cos\theta}{g}$

But $\sin 2\theta = 2\sin\theta\cos\theta$

so $x = \dfrac{V^2\sin 2\theta}{g}$ metres

Hence the horizontal range is $\dfrac{V^2\sin 2\theta}{g}$ metres.

© Cambridge University Press 2019

16b If the horizontal range is five times the maximum height,

$$\frac{V^2 \sin^2\theta}{2g} \div \frac{V^2 \sin 2\theta}{g} = \frac{1}{5}$$

$$\frac{gV^2 \sin^2\theta}{2gV^2 \sin 2\theta} = \frac{1}{5}$$

$$\frac{\sin^2\theta}{2 \sin 2\theta} = \frac{1}{5}$$

$$5 \sin^2\theta = 2 \sin 2\theta$$

$$5 \sin^2\theta = 2 \times 2 \sin\theta \cos\theta$$

$$5 \sin^2\theta = 4 \sin\theta \cos\theta$$

$$5 \sin\theta = 4 \cos\theta \text{ where } \sin\theta \neq 0$$

$$\frac{\sin\theta}{\cos\theta} = \frac{4}{5}$$

$$\tan\theta = \frac{4}{5}$$

$$\text{so } \theta = \tan^{-1}\frac{4}{5} = \arctan\frac{4}{5}$$

17 The usual equations of motion are:

$\dot{x} = V\cos\theta$ and $\dot{y} = -gt + V\sin\theta$

$x = Vt\cos\theta$ $y = -\frac{g}{2}t^2 + Vt\sin\theta$

The maximum height occurs when $\dot{y} = 0$.

That is, when $t = \frac{V\sin\theta}{g}$.

So when $t = \frac{V\sin\theta}{g}$, $y = 2$

Hence,

$$2 = -\frac{g}{2} \times \frac{V^2 \sin^2\theta}{g^2} + V\sin\theta \times \frac{V\sin\theta}{g}$$

$$2 = \frac{V^2 \sin^2\theta}{2g}$$

© Cambridge University Press 2019

$\sin^2\theta = \frac{4g}{V^2}$

$\sin\theta = \frac{2\sqrt{g}}{V}$ (θ is acute)

The time of flight is $\frac{2V\sin\theta}{g}$.

When $t = \frac{2V\sin\theta}{g}$,

$x = V\cos\theta \times \frac{2V\sin\theta}{g}$

$x = \frac{2V^2\sin\theta\cos\theta}{g}$

If $\sin\theta = \frac{2\sqrt{g}}{V}$,

consider a right-angled triangle with angle θ and opposite side $2\sqrt{g}$ units, hypotenuse V units and adjacent side $\sqrt{V^2 - (2\sqrt{g})^2} = \sqrt{V^2 - 4g}$ units.

Hence, $\cos\theta = \frac{\sqrt{V^2 - 4g}}{V}$

Therefore,

$x = \frac{2V^2}{g} \times \frac{2\sqrt{g}}{V} \times \frac{\sqrt{V^2 - 4g}}{V}$

$x = \frac{4}{\sqrt{g}}\sqrt{V^2 - 4g}$

$x = \sqrt{\frac{16}{g}(V^2 - 4g)}$ as required.

© Cambridge University Press 2019

18a $\dot{x} = V\cos\theta$ and $\dot{y} = -gt + V\sin\theta$

$x = Vt\cos\theta$ $y = -\frac{g}{2}t^2 + Vt\sin\theta$

The greatest height occurs when $\dot{y} = 0$

That is, when $t = \dfrac{V\sin\theta}{g}$.

So when $t = \dfrac{V\sin\theta}{g}$, $y = h$

Hence,

$$h = -\frac{g}{2} \times \frac{V^2\sin^2\theta}{g^2} + V\sin\theta \times \frac{V\sin\theta}{g}$$

$$h = \frac{V^2\sin^2\theta}{2g}$$

$$V^2\sin^2\theta = 2gh \qquad (1)$$

18b

When $y = \dfrac{h}{2}$,

$$\frac{h}{2} = -\frac{gt^2}{2} + Vt\sin\theta$$

$$gt^2 - (2V\sin\theta)t + h = 0$$

Using the quadratic formula:

$$t = \frac{2V\sin\theta \pm \sqrt{4V^2\sin^2\theta - 4gh}}{2g}$$

$$t = \frac{V\sin\theta \pm \sqrt{V^2\sin^2\theta - gh}}{g}$$

From (1), $V^2\sin^2\theta = 2gh$ hence,

$$t = \frac{\sqrt{2gh} \pm \sqrt{2gh - gh}}{g}$$

$$t = \frac{\sqrt{2}\sqrt{gh} \pm \sqrt{gh}}{g}$$

© Cambridge University Press 2019

$$t = \frac{\sqrt{gh}(\sqrt{2} \pm 1)}{g}$$

$$t = \frac{(\sqrt{2} \pm 1)\sqrt{h}}{\sqrt{g}}$$

Hence,

$$t = \frac{(\sqrt{2} + 1)\sqrt{h}}{\sqrt{g}} \text{ or } t = \frac{(\sqrt{2} - 1)\sqrt{h}}{\sqrt{g}} \text{ as required.}$$

18c At $y = h,\ \dot{y} = 0$, so the speed is equal to the horizontal speed, which is $V\cos\theta$.

At $y = \frac{h}{2}$, while the projectile is rising,

$$t = \frac{(\sqrt{2} - 1)\sqrt{h}}{\sqrt{g}}$$

So $V\cos\theta$, and

$$\dot{y} = V\sin\theta - g \times \frac{(\sqrt{2} - 1)\sqrt{h}}{\sqrt{g}}$$

$$\dot{y} = V\sin\theta - (\sqrt{2} - 1)\sqrt{gh}$$

From (1) in part a:

$$V^2\sin^2\theta = 2gh$$

$$gh = \frac{V^2\sin^2\theta}{2}$$

$$\sqrt{gh} = \frac{V\sin\theta}{\sqrt{2}}$$

Hence

$$\dot{y} = V\sin\theta - (\sqrt{2} - 1)\frac{V\sin\theta}{\sqrt{2}}$$

$$\dot{y} = \frac{\sqrt{2}V\sin\theta - \sqrt{2}V\sin\theta + V\sin\theta}{\sqrt{2}}$$

$$\dot{y} = \frac{V\sin\theta}{\sqrt{2}}$$

© Cambridge University Press 2019

At $y = \frac{h}{2}$,

$$\text{speed} = \sqrt{(\dot{x})^2 + (\dot{y})^2}$$

$$= \sqrt{V^2 \cos^2\theta + \frac{V^2 \sin^2\theta}{2}}$$

$$= \frac{V}{\sqrt{2}}\sqrt{2\cos^2\theta + \sin^2\theta}$$

So the ratio of the speed is:

$$\frac{\frac{V}{\sqrt{2}}\sqrt{2\cos^2\theta + \sin^2\theta}}{V\cos\theta} = \frac{\sqrt{5}}{\sqrt{2}}$$

$$\frac{\sqrt{2\cos^2\theta + \sin^2\theta}}{\cos\theta} = \sqrt{5}$$

$$\sqrt{5}\cos\theta = \sqrt{2\cos^2\theta + \sin^2\theta}$$

$$5\cos^2\theta = 2\cos^2\theta + \sin^2\theta$$

$$5\cos^2\theta = 2\cos^2\theta + 1 - \cos^2\theta$$

$$4\cos^2\theta = 1$$

$$\cos^2\theta = \frac{1}{4}$$

$$\cos\theta = \frac{1}{2} \qquad (\text{as } \cos\theta > 0)$$

$$\theta = \frac{\pi}{3} \text{ or } 60°$$

© Cambridge University Press 2019

Solutions to Exercise 10B

1a

$$y = -\frac{5}{324}x^2 + \frac{4}{3}x$$

When $x = 12$,

$$y = -\frac{5}{324} \times 12^2 + \frac{4}{3} \times 12$$

$$= -\frac{720}{324} + 16$$

$$= 13\frac{7}{9} \text{ m}$$

or $y \div 13.8$ m

The height of the particle is about 13.8 m.

1b When $y = 19$,

$$19 = -\frac{5}{324}x^2 + \frac{4}{3}x$$

$$5x^2 - 432x + 6156 = 0$$

Using the quadratic formula:

$$x = \frac{432 \pm \sqrt{432^2 - 4 \times 5 \times 6156}}{2 \times 5}$$

$$= \frac{432 \pm \sqrt{63\,504}}{10}$$

$$= \frac{432 \pm 252}{10}$$

$$= 43.2 - 25.2 \text{ or } 43.2 + 25.2$$

So $x = 18$ m or 68.4. m

The particle has travelled a horizontal distance of 18 m or 68.4 m.

© Cambridge University Press 2019

1c i

$$\frac{dy}{dx} = -\frac{2 \times 5}{324}x + \frac{4}{3}$$

$$= -\frac{5}{162}x + \frac{4}{3}$$

When $x = 0$,

$$\frac{dy}{dx} = -\frac{5}{162} \times 0 + \frac{4}{3}$$

$$= \frac{4}{3}$$

Let $\alpha =$ angle of projection

$$\tan\alpha = \frac{dy}{dx} = \frac{4}{3}$$

$$\alpha = \tan^{-1}\frac{4}{3}$$

So the angle of projection is $\tan^{-1}\frac{4}{3}$ above the horizontal.

1c ii When $x = 18$,

$$\frac{dy}{dx} = -\frac{5}{162} \times 18 + \frac{4}{3}$$

$$= \frac{7}{9}$$

Let $\alpha =$ angle of projection

$$\tan\alpha = \frac{dy}{dx} = \frac{7}{9}$$

$$\alpha = \tan^{-1}\frac{7}{9}$$

So the direction of motion is $\tan^{-1}\frac{7}{9}$ above the horizontal.

1c iii When $x = 54$,

$$\frac{dy}{dx} = -\frac{5}{162} \times 54 + \frac{4}{3}$$

© Cambridge University Press 2019

$= -\frac{1}{3}$

Let $\alpha =$ angle of projection

$\tan\alpha = \frac{dy}{dx} = -\frac{1}{3}$

$\alpha = \tan^{-1}\left(-\frac{1}{3}\right)$

So the direction of motion is $\tan^{-1}\frac{1}{3}$ below the horizontal.

2a Initially, $t = 0 = x = y$.

$x = 48t$

so $t = \frac{x}{48}$ (1)

and

$y = -5t^2 + 20t$ (2)

Substituting (1) into (2):

$y = -5\left(\frac{x}{48}\right)^2 + 20\left(\frac{x}{48}\right)$

$y = -\frac{5}{2304}x^2 + \frac{5}{12}x$ (3)

2b i When $y = 0$,

$0 = -\frac{5}{2304}x^2 + \frac{5}{12}x$

$\frac{5}{2304}x^2 - \frac{5}{12}x = 0$

$5x^2 - 960x = 0$

$5x(x - 192) = 0$

$x = 0$ or 192

Therefore, horizontal range of particle is 192 m.

© Cambridge University Press 2019

2b ii Using symmetry, greatest height is at midpoint of horizontal range.

$$x = \frac{192}{2} = 96$$

Thus, $x = 96$ m at greatest height.

Substituting into (3):

$$y = -\frac{5}{2304} \times 96^2 + \frac{5}{12} \times 96$$

$$= 20 \text{ m}$$

Alternatively, $\frac{dy}{dx} = 0$ for maximum/minimum height.

$$-\frac{5x}{1152} + \frac{5}{12} = 0$$

$$5x - \frac{5 \times 1152}{12} = 0$$

$$5x = 480$$

$$x = 96$$

$\frac{d^2y}{dx^2} = -\frac{5}{1152} < 0$, so $x = 96$ gives maximum height.

Substituting $x = 96$ into (3):

$$y = -\frac{5}{2304} \times 96^2 + \frac{5}{12} \times 96$$

$$= 20 \text{ m}$$

The greatest height of the particle is 20 m.

2b iii Initially, $t = x = 0$

$$\frac{dy}{dx} = -\frac{5x}{1152} + \frac{5}{12}$$

When $x = 0$,

$$\frac{dy}{dx} = -\frac{5 \times 0}{1152} + \frac{5}{12}$$

$$= \frac{5}{12}$$

Let α = angle of projection

© Cambridge University Press 2019

$$\tan\alpha = \frac{dy}{dx} = \frac{5}{12}$$

$$\alpha = \tan^{-1}\frac{5}{12}$$

$$\alpha \div 22.6°$$

The angle of projection is about 22.6°.

2b iv When $x = 120$,

$$\frac{dy}{dx} = -\frac{5 \times 120}{1152} + \frac{5}{12}$$

$$= -\frac{25}{48} + \frac{5}{12}$$

$$= -\frac{5}{48}$$

Let α = angle of projection

$$\tan\alpha = \frac{dy}{dx} = -\frac{5}{48}$$

$$\alpha = \tan^{-1}\left(-\frac{5}{48}\right) \div -6°$$

The direction of the particle will be 6° below the horizontal.

3a $|\underline{v}| = \sqrt{3^2 + 1^2}$

$$= \sqrt{10}$$

The initial speed is $\sqrt{10}$.

3b To begin, $\dot{x} = 3$

Integrating, $x = 3t + C_1$

When $t = 0$, $x = 0$

Thus, $C_1 = 0$

so $x = 3t$ (1)

© Cambridge University Press 2019

To begin, $\dot{y} = 1 - 10t$

Integrating, $y = t - 5t^2 + C_2$

When $t = 0$, $y = 0$

Thus, $C_2 = 0$

so $y = t - 5t^2$ (2)

3c $x = 3t$

$t = \frac{x}{3}$ (3)

Substituting (3) into (2):

$y = \frac{x}{3} - 5\left(\frac{x}{3}\right)^2$

$= \frac{x}{3} - \frac{5x^2}{9}$

$9y = 3x - 5x^2$

3d i

$\frac{dy}{dx} = \frac{1}{3} - \frac{10}{9}x$

When $x = 0.15$,

$\frac{dy}{dx} = \frac{1}{3} - \frac{10}{9} \times 0.15$

$= \frac{6}{18} - \frac{3}{18}$

$= \frac{3}{18}$

$= \frac{1}{6}$

Let α = angle of projection

$\tan\alpha = \frac{dy}{dx} = \frac{1}{6}$

© Cambridge University Press 2019

$\alpha = \tan^{-1}\left(\frac{1}{6}\right) \div 9.5°$

The direction of motion of the particle will be 9.5° above the horizontal.

3d ii

$\frac{dy}{dx} = \frac{1}{3} - \frac{10}{9}x$

When $t = 0.15, x = 3 \times 0.15 = 0.45$ (using $x = 3t$)

$\frac{dy}{dx} = \frac{1}{3} - \frac{10}{9} \times 0.45$

$= \frac{6}{18} - \frac{9}{18}$

$= -\frac{3}{18}$

$= -\frac{1}{6}$

Let α = angle of projection

$\tan\alpha = \frac{dy}{dx} = -\frac{1}{6}$

$\alpha = \tan^{-1}\left(-\frac{1}{6}\right) \div -9.5°$

The direction of motion of the particle will be 9.5° below the horizontal.

4a To begin, $\dot{x} = 5$

Integrating, $x = 5t + C_1$

When $t = 0$, $x = 0$

Thus, $C_1 = 0$

so $x = 5t$ (1)

To begin, $\dot{y} = -10t$

Integrating, $y = -5t^2 + C_2$

When $t = 0$, $y = 0$

© Cambridge University Press 2019

Thus, $C_2 = 0$

so $y = -5t^2$ (2)

From (1):

$t = \frac{x}{5}$

Substituting $t = \frac{x}{5}$ into (2):

$y = -5\left(\frac{x}{5}\right)^2$

$y = -\frac{1}{5}x^2$

4b Object lands at $y = -20$ m

$-20 = -\frac{1}{5}x^2$

$100 = x^2$

$x = 10$ m (as $x > 0$)

The object hits the ground 10 m from the base of the tower.

4c Object hits the ground when $x = 10$ m.

$t = \frac{10}{5}$

$= 2$ s

$\frac{dy}{dx} = -\frac{2}{5}x$

When $x = 10$,

$\frac{dy}{dx} = -\frac{2}{5} \times 10$

$= -4$

Let α = angle of projection

$\tan\alpha = \frac{dy}{dx} = -4$

$\alpha = \tan^{-1}(-4) \div -76°$

© Cambridge University Press 2019

The direction of the particle will be 76° below the horizontal.

5a To begin, $\dot{x} = 24\cos 30° = 12\sqrt{3}$

Integrating, $x = 12\sqrt{3}t + C_1$

When $t = 0$, $x = 0$

Thus, $C_1 = 0$

so $x = 12\sqrt{3}t$ (1)

To begin, $\dot{y} = 24\sin 30° - 10t = 12 - 10t$

Integrating, $y = 12t - 5t^2 + C_2$

When $t = 0$, $y = 0$

Thus, $C_2 = 0$

so $y = 12t - 5t^2$ (2)

5b From (1):

$$t = \frac{x}{12\sqrt{3}}$$

Substituting $t = \frac{x}{12\sqrt{3}}$ into (2):

$$y = 12\left(\frac{x}{12\sqrt{3}}\right) - 5\left(\frac{x}{12\sqrt{3}}\right)^2$$

$$= \frac{1}{\sqrt{3}}x - \frac{5}{432}x^2$$

5c When the ball hits the ground, $x = D$ and $y = -6$.

$$-6 = \frac{1}{\sqrt{3}}D - \frac{5}{432}D^2$$

$$\frac{5}{432}D^2 - \frac{1}{\sqrt{3}}D - 6 = 0$$

$$5D^2 - \frac{432}{\sqrt{3}}D - 2592 = 0$$

© Cambridge University Press 2019

$$5D^2 - \frac{432\sqrt{3}}{3}D - 2592 = 0$$

$$5D^2 - 144\sqrt{3}D - 2592 = 0$$

5d Using the quadratic formula:

$$D = \frac{144\sqrt{3} \pm \sqrt{\left(144\sqrt{3}\right)^2 - 4 \times 5 \times (-2592)}}{2 \times 5}$$

$$D = \frac{144\sqrt{3} \pm \sqrt{62\,208 + 51\,840}}{10}$$

$$D = \frac{144\sqrt{3} \pm \sqrt{114\,048}}{10}$$

$D \div -8.8$ or 58.7

But $D > 0$ so $D \div 58.7$ m.

6 To begin, $\dot{x} = V\cos 27°$

Integrating, $x = Vt\cos 27° + C_1$

When $t = 0$, $x = 0$

Thus, $C_1 = 0$

so $x = Vt\cos 27°$ (1)

To begin, $\dot{y} = -V\sin 27° - 10t$

Integrating, $y = -Vt\sin 27° - 5t^2 + C_2$

When $t = 0$, $y = 0$

Thus, $C_2 = 0$

so $y = -Vt\sin 27° - 5t^2$ (2)

From (1):

$$t = \frac{x}{V\cos 27°} \quad (3)$$

Substituting (3) into (2):

© Cambridge University Press 2019

$$y = -V\left(\frac{x}{V\cos 27°}\right)\sin 27° - 5\left(\frac{x}{V\cos 27°}\right)^2$$

$$y = -\tan 27°\, x - \frac{5x^2}{V^2\cos^2 27°}$$

$$y = -x\tan 27° - \frac{5x^2}{V^2\cos^2 27°} \qquad \text{as required}$$

6b i Since $y = -60$ when $x = 35$,

$$-60 = -35\tan 27° - \frac{5(35)^2}{V^2\cos^2 27°}$$

$$60 - 35\tan 27° = \frac{5(35)^2}{V^2\cos^2 27°}$$

$$V^2\cos^2 27° = \frac{5(35)^2}{60 - 35\tan 27°}$$

$$V^2 = \frac{5(35)^2}{\cos^2 27°\,(60 - 35\tan 27°)}$$

$$V = \sqrt{\frac{5(35)^2}{\cos^2 27°\,(60 - 35\tan 27°)}} \qquad \text{as } V > 0$$

$V \doteqdot 13.5$ m/s

The initial speed of the stone is about 13.5 m/s.

6b ii

$$\frac{dy}{dx} = -\tan 27° - \frac{10x}{V^2\cos^2 27°}$$

When the stone lands in the ocean, $x = 35$ and $V \doteqdot 13.5$.

$$\frac{dy}{dx} \doteqdot -\tan 27° - \frac{10 \times 35}{13.5^2\cos^2 27°}$$

$$= -2.9285\ldots$$

Let α = angle of projection

$$\tan\alpha = \frac{dy}{dx} \doteqdot -2.93$$

$$\alpha = \tan^{-1}(-2.93) \doteqdot -71°$$

© Cambridge University Press 2019

The direction of the stone will be 71° below the horizontal.

7a

$$y = x\tan\theta - \frac{gx^2\sec^2\theta}{2V^2}$$

$y = 0$ when particle lands

$$0 = x\tan\theta - \frac{gx^2}{2V^2\cos^2\theta}$$

$$0 = \frac{x\sin\theta}{\cos\theta} - \frac{gx^2}{2V^2\cos^2\theta}$$

$$0 = \frac{x\sin\theta\,(2V^2\cos\theta)}{2V^2\cos^2\theta} - \frac{gx^2}{2V^2\cos^2\theta}$$

$$0 = \frac{2V^2x\sin\theta\cos\theta}{2V^2\cos^2\theta} - \frac{gx^2}{2V^2\cos^2\theta}$$

$gx^2 - 2V^2x\sin\theta\cos\theta = 0$ (for non-zero range, $\cos\theta \neq 0$ and $V \neq 0$)

$x\,(gx - 2V^2\sin\theta\cos\theta) = 0$

$x = 0$ or $gx = 2V^2\sin\theta\cos\theta$

$x = 0$ is not a feasible solution so

$$x = \frac{2V^2\sin\theta\cos\theta}{g}$$

$$x = \frac{V^2\sin 2\theta}{g}$$ (as $\sin 2\theta = 2\sin\theta\cos\theta$)

So the horizontal range is $\frac{V^2\sin 2\theta}{g}$ as required.

.

7b $V = 30$ m/s and $x = 75$ m. Assume $g = 10$ m/s^2.

Substituting into $x = \frac{V^2\sin 2\theta}{g}$ gives:

$$75 = \frac{30^2 \times \sin 2\theta}{10}$$

$$\sin 2\theta = \frac{75 \times 10}{30^2}$$

© Cambridge University Press 2019

$$\sin 2\theta = \frac{5}{6}$$

$$2\theta \div 56.4°, 123.6°$$

$$\theta \div 28.2°, 61.8°$$

8a

$$y = x\tan\theta - \frac{gx^2\sec^2\theta}{2V^2}$$

$$y = x\tan\theta - \frac{g}{2V^2\cos^2\theta}x^2$$

$$\frac{dy}{dx} = \tan\theta - \frac{g}{V^2\cos^2\theta}x$$

Vertex when $\frac{dy}{dx} = 0$

$$0 = \tan\theta - \frac{g}{V^2\cos^2\theta}x$$

$$\frac{g}{V^2\cos^2\theta}x = \tan\theta$$

$$\frac{g}{V^2\cos^2\theta}x = \frac{\sin\theta}{\cos\theta}$$

$$x = \frac{\sin\theta}{\cos\theta} \times \frac{V^2\cos^2\theta}{g}$$

$$x = \frac{V^2\sin\theta\cos\theta}{g} \qquad \text{where } \cos\theta \neq 0$$

$$x = \frac{V^2 \times 2\sin\theta\cos\theta}{2g}$$

$$x = \frac{V^2\sin 2\theta}{2g} \qquad (\text{since } \sin 2\theta = 2\cos\theta\sin\theta)$$

$$\frac{d^2y}{dx^2} = -\frac{g}{V^2\cos^2\theta} < 0.$$

Hence $x = \frac{V^2\sin 2\theta}{2g}$ gives the maximum turning point of the parabola.

When $x = \frac{V^2\sin 2\theta}{2g}$,

© Cambridge University Press 2019

$$y = \left(\frac{V^2 \sin 2\theta}{2g}\right)\tan\theta - \frac{g}{2V^2\ \cos^2\theta}\left(\frac{V^2 \sin 2\theta}{2g}\right)^2$$

$$= \left(\frac{V^2 \sin 2\theta}{2g}\right)\tan\theta - \frac{g}{2V^2\ \cos^2\theta}\left(\frac{V^2 2\sin\theta\cos\theta}{2g}\right)^2$$

$$= \left(\frac{V^2 \sin\theta\cos\theta}{g}\right) \times \frac{\sin\theta}{\cos\theta} - \frac{g}{2V^2\cos^2\theta} \times \left(\frac{V^4 4\sin^2\theta\cos^2\theta}{4g^2}\right)$$

$$= \frac{V^2\sin^2\theta}{g} - \frac{V^2\sin^2\theta}{2g}$$

$$= \frac{2V^2\sin^2\theta}{2g} - \frac{V^2\sin^2\theta}{2g}$$

$$= \frac{V^2\sin^2\theta}{2g}$$

Thus the coordinates of the vertex are $\left(\frac{V^2 \sin 2\theta}{2g}, \frac{V^2\sin^2\theta}{2g}\right)$.

8b $V = 20$ m/s and $y = 15$ m

At the greatest height,

$$y = \frac{V^2\sin^2\theta}{2g}$$

$$15 = \frac{(20)^2\sin^2\theta}{2 \times 10}$$

$$\sin^2\theta = \frac{300}{400}$$

$$\sin^2\theta = \frac{3}{4}$$

$$\sin\theta = \pm\frac{\sqrt{3}}{2}$$

For angle of projection above the horizontal, $\sin\theta > 0$.

So $\theta = 60°$.

© Cambridge University Press 2019

9a $x = Vt\cos\alpha$ (1)

$y = -\frac{1}{2}gt^2 + Vt\sin\alpha$ (2)

Rearranging (1):

$$t = \frac{x}{V\cos\alpha}$$

Substituting for t in (2):

$$y = -\frac{1}{2}g\left(\frac{x}{V\cos\alpha}\right)^2 + V\left(\frac{x}{V\cos\alpha}\right)\sin\alpha$$

$$= -\frac{gx^2}{2V^2\cos^2\alpha} + \left(\frac{x}{\cos\alpha}\right)\sin\alpha$$

$$= -\frac{gx^2}{2V^2\cos^2\alpha} + x\tan\alpha$$

$$= -\frac{gx^2\sec^2\alpha}{2V^2} + x\tan\alpha$$

$$= -\frac{gx^2(1+\tan^2\alpha)}{2V^2} + x\tan\alpha$$

$$2yV^2 = -gx^2(1+\tan^2\alpha) + 2xV^2\tan\alpha$$

$$2yV^2 + gx^2(1+\tan^2\alpha) - 2xV^2\tan\alpha = 0$$

$$2yV^2 + gx^2 + gx^2\tan^2\alpha - 2xV^2\tan\alpha = 0$$

$$gx^2\tan^2\alpha - 2xV^2\tan\alpha + (2yV^2 + gx^2) = 0$$

9b $V = 200$ m/s

$x = 3000$ m

$y = 500$ m

$$gx^2\tan^2\alpha - 2xV^2\tan\alpha + (2yV^2 + gx^2) = 0$$

$$(10)(3000)^2\tan^2\alpha - 2(3000)(200)^2\tan\alpha + 2(500)(200)^2 + 10(3000)^2 = 0$$

$$90\,000\,000\tan^2\alpha - 240\,000\,000\tan\alpha + 130\,000\,000 = 0$$

$$9\tan^2\alpha - 24\tan\alpha + 13 = 0$$

Using the quadratic formula:

$$\tan\alpha = \frac{24 \pm \sqrt{24^2 - 4\times 9\times 13}}{2\times 9}$$

© Cambridge University Press 2019

$$\tan\alpha = \frac{24 \pm \sqrt{108}}{18}$$

$$\tan\alpha = \frac{24 \pm 6\sqrt{3}}{18}$$

$$\tan\alpha = \frac{4 \pm \sqrt{3}}{3}$$

$$\tan\alpha = \frac{4+\sqrt{3}}{3} \text{ or } \tan\alpha = \frac{4-\sqrt{3}}{3}$$

$\alpha \div 62°22'$ or $37°5'$

10a To begin, $\dot{x} = 34\cos\theta$

Integrating, $x = 34t\cos\theta + C_1$

When $t = 0$, $x = 0$

Thus, $C_1 = 0$

so $x = 34t\cos\theta$

and $t = \frac{x}{34\cos\theta}$ (1)

To begin, $\dot{y} = 34\sin\theta - 10t$

Integrating, $y = 34t\sin\theta - 5t^2 + C_2$

When $t = 0$, $y = 0$

Thus, $C_2 = 0$

so $y = 34t\sin\theta - 5t^2$ (2)

Substituting (1) into (2):

$$y = 34\left(\frac{x}{34\cos\theta}\right)\sin\theta - 5\left(\frac{x}{34\cos\theta}\right)^2$$

$$y = \left(\frac{x}{\cos\theta}\right)\sin\theta - \frac{5x^2}{1156\cos^2\theta}$$

$$y = x\tan\theta - \frac{5x^2\sec^2\theta}{1156}$$

© Cambridge University Press 2019

10b $x = 30$ m

$y = 11$ m

$$11 = 30\tan\theta - \frac{5(30)^2\sec^2\theta}{1156}$$

$$11 = 30\tan\theta - \frac{4500(1+\tan^2\theta)}{1156}$$

$$12\,716 = 34\,680\tan\theta - 4500\,(1+\tan^2\theta)$$

$$12\,716 = 34\,680\tan\theta - 4500 - 4500\tan^2\theta$$

$$4500\ \tan^2\theta - 34\,680\tan\theta + 17\,216 = 0$$

$$1125\ \tan^2\theta - 8670\tan\theta + 4304 = 0$$

Using the quadratic formula:

$$\tan\theta = \frac{8670 \pm \sqrt{(-8670)^2 - 4\times 1125\times 4304}}{2\times 1125}$$

$$\tan\theta = \frac{8670 \pm \sqrt{55\,800\,900}}{2250}$$

$$\tan\theta = \frac{8670 \pm 7470}{2250}$$

$$\tan\theta = \frac{1200}{2250} \text{ or } \frac{16\,140}{2250}$$

$$\tan\theta = \frac{8}{15} \text{ or } \frac{538}{75}$$

$$\theta = \tan^{-1}\frac{8}{15} \text{ or } \tan^{-1}\frac{538}{75}$$

$$\theta = \arctan\frac{8}{15} \text{ or } \arctan\frac{538}{75}$$

11a Initially, $x = y = 0$, and, $\dot{x} = V\cos 45°$ $\dot{y} = V\sin 45°$

To begin, $\ddot{x} = 0$ To begin, $\ddot{y} = -g$

Integrating, $\dot{x} = C_1$ Integrating, $\dot{y} = -gt + C_2$

When $t = 0$, $\dot{x} = V\cos 45° = \frac{V}{\sqrt{2}}$ When $t = 0$, $\dot{y} = V\sin 45° = \frac{V}{\sqrt{2}}$

$\frac{V}{\sqrt{2}} = C_1$ $\frac{V}{\sqrt{2}} = C_2$

© Cambridge University Press 2019

so $\dot{x} = \frac{V}{\sqrt{2}}$ so $\dot{y} = -gt + \frac{V}{\sqrt{2}}$

so $\dot{x} = \frac{V}{\sqrt{2}}$ (2) so $\dot{y} = -gt + \frac{V}{\sqrt{2}}$

Integrating, $x = \frac{Vt}{\sqrt{2}} + C_3$ Integrating, $y = \frac{-gt^2}{2} + \frac{Vt}{\sqrt{2}} + C_4$

When $t = 0$, $x = 0$ When $t = 0$, $y = 0$

$0 = C_3$ $0 = C_4$

so $x = \frac{Vt}{\sqrt{2}} = \frac{Vt\sqrt{2}}{2}$ (1) so $y = \frac{-gt^2}{2} + \frac{Vt\sqrt{2}}{2}$ (2)

11b From (1):

$$t = \frac{\sqrt{2}x}{V}$$


Substituting for t in (2):

$$y = \frac{-g}{2} \times \left(\frac{\sqrt{2}x}{V}\right)^2 + \frac{V\sqrt{2}}{2} \times \frac{\sqrt{2}x}{V}$$

$$y = \frac{-gx^2}{V^2} + x$$

$$y = x - \frac{gx^2}{V^2} \qquad (3)$$

11c



© Cambridge University Press 2019

The coordinates of P and Q satisfy equation (3),

so $30\sin\theta = 30\cos\theta - \frac{g}{V^2} \times 900\cos^2\theta$

and $-60\sin\theta = 60\cos\theta - \frac{g}{V^2} \times 3600\cos^2\theta$

so $\sin\theta = \cos\theta - \frac{30g\cos^2\theta}{V^2}$

and $-\sin\theta = \cos\theta - \frac{60g\cos^2\theta}{V^2}$

Dividing through by $\cos\theta$,

$\tan\theta = 1 - \frac{30g\cos\theta}{V^2}$ and $-\tan\theta = 1 - \frac{60g\cos\theta}{V^2}$

So $\tan\theta = 1 - \frac{30g\cos\theta}{V^2}$ (4)

and $\tan\theta = \frac{60g\cos\theta}{V^2} - 1$

Hence, $1 - \frac{30g\cos\theta}{V^2} = \frac{60g\cos\theta}{V^2} - 1$

$V^2 - 30g\cos\theta = 60g\cos\theta - V^2$

$2V^2 = 90g\cos\theta$

$V^2 = 45g\cos\theta$

Substituting for V^2 into (4):

$\tan\theta = 1 - \frac{30g\cos\theta}{45g\cos\theta}$

$= 1 - \frac{2}{3}$

$= \frac{1}{3}$

$\theta = \tan^{-1}\frac{1}{3}$

© Cambridge University Press 2019

12a $x = Vt\cos\alpha$ (1)

$y = Vt\sin\alpha - \frac{1}{2}gt^2$ (2)

Rearranging (1):

$t = \dfrac{x}{V\cos\alpha}$

Substituting for t in (2):

$$y = V\left(\frac{x}{V\cos\alpha}\right)\sin\alpha - \frac{1}{2}g\left(\frac{x}{V\cos\alpha}\right)^2$$

$$= \left(\frac{x}{\cos\alpha}\right)\sin\alpha - \frac{gx^2}{2V^2\cos^2\alpha}$$

$$= x\tan\alpha - \frac{gx^2}{2V^2\cos^2\alpha} \quad (3)$$

$y = 0$ when particle lands, so

$$0 = x\tan\alpha - \frac{gx^2}{2V^2\cos^2\alpha}$$

$$0 = \frac{x\sin\alpha}{\cos\theta} - \frac{gx^2}{2V^2\cos^2\alpha}$$

$$0 = \frac{x\sin\alpha\,(2V^2\cos\alpha)}{2V^2\cos^2\alpha} - \frac{gx^2}{2V^2\cos^2\alpha}$$

$$0 = \frac{2V^2x\sin\alpha\cos\alpha}{2V^2\cos^2\alpha} - \frac{gx^2}{2V^2\cos^2\alpha}$$

$gx^2 - 2V^2x\sin\alpha\cos\alpha = 0$ (for non-zero range, $\cos\theta \neq 0$ and $V \neq 0$)

$x\,(gx - 2V^2\sin\alpha\cos\alpha) = 0$

$x = 0$ or $gx = 2V^2\sin\alpha\cos\alpha$

$x = 0$ is not a feasible solution so

$$x = \frac{2V^2\sin\alpha\cos\alpha}{g}$$

$$x = \frac{V^2\sin 2\alpha}{g} \quad (\text{as } \sin 2\alpha = 2\sin\alpha\cos\alpha)$$

So the horizontal range is $\dfrac{V^2\sin 2\alpha}{g}$ as required.

© Cambridge University Press 2019

12b From (3) above,

$$y = x\tan\alpha - \frac{gx^2}{2V^2\cos^2\alpha}$$

$$y = x\tan\alpha - \frac{gx^2}{2V^2\cos^2\alpha} \times \frac{x\tan\alpha}{x\tan\alpha}$$

$$y = x\tan\alpha\left(1 - \frac{gx^2}{2V^2\cos^2\alpha} \times \frac{1}{x\tan\alpha}\right)$$

$$y = x\tan\alpha\left(1 - \frac{gx^2}{2V^2\cos^2\alpha} \times \frac{\cos\alpha}{x\sin\alpha}\right)$$

$$y = x\tan\alpha\left(1 - \frac{gx}{2V^2\cos\alpha\sin\alpha}\right)$$

$$y = x\tan\alpha\left(1 - \frac{gx}{V^2\sin 2\alpha}\right)$$

$$y = x\tan\alpha\left(1 - \frac{x}{R}\right), \text{since } R = \frac{V^2\sin 2\alpha}{g}$$

$$y = x\left(1 - \frac{x}{R}\right)\tan\alpha$$

12c $\alpha = 45°$, so the path of the equation is:

$$y = x\left(1 - \frac{x}{R}\right)$$

$$\text{or } y = x - \frac{x^2}{R}$$

12c i Let $y = 4$, hence:

$$4 = x - \frac{x^2}{R}$$

$$x^2 - Rx + 4R = 0$$

Since $y = 4$ corresponds to $x = x_1$ and $x = x_2$,

x_1 and x_2 must be the roots of the equation $x^2 - Rx + 4R = 0$.

© Cambridge University Press 2019

12c ii $(x_2 - x_1)^2 = (x_2 + x_1)^2 - 4x_2x_1$

Hence,

$6^2 = (x_2 + x_1)^2 - 4(x_2x_1)$

$36 = R^2 - 4(4R)$

$R^2 - 16R - 36 = 0$

$(R - 18)(R + 2) = 0$

$R = 18$, since $R > 0$

So the horizontal range, R, is 18 metres.

© Cambridge University Press 2019

Solutions to Chapter review

1a Initially, $x = y = 0$, and, $\dot{x} = 60\cos 40°$ $\dot{y} = 60\sin 40°$

To begin, $\ddot{x} = 0$ (1) To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{x} = C_1$ Integrating, $\dot{y} = -10t + C_2$

When $t = 0$, $\dot{x} = 60\cos 40°$ When $t = 0$, $\dot{y} = 60\sin 40°$

$60\cos 40° = C_1$ $60\sin 40° = C_2$

so $\dot{x} = 60\cos 40°$ (2) so $\dot{y} = -10t + 60\sin 40°$ (4)

1b To begin, $\dot{x} = 60\cos 40°$ To begin, $\dot{y} = -10t + 60\sin 40°$

Integrating, $x = 60t\cos 40° + C_3$ Integrating, $y = -5t^2 + 60t\sin 40° + C_4$

When $t = 0$, $x = 0$ When $t = 0$, $y = 0$

$0 = C_3$ $0 = C_4$

so $x = 60t\cos 40°$ (5) so $y = -5t^2 + 60t\sin 40°$ (6)

1c i The particle returns to the ground when $y = 0$,

Substituting into (6):

$0 = -5t^2 + 60t\sin 40°$

$0 = -5t(t - 12\sin 40°)$

$t = 0$ or $t = 12\sin 40° \div 7.7$ seconds

Particle returns to the ground at $t > 0$ seconds, so at $t \div 7.7$ seconds.

1c ii When $t = 12\sin 40°$, (5) becomes:

$x = 720\sin 40°\cos 40°$

$\div 354.5$ m

The horizontal distance travelled by the particle is about 354.5 m.

© Cambridge University Press 2019

1c iii Particle reaches greatest height when $\dot{y} = 0$.

Substituting into (4):

$0 = -10t + 60\sin 40°$

$10t = 60\sin 40°$

$t = 6\sin 40°$

Substituting for t in (6):

$y = -5(6\sin 40°)^2 + 60(6\sin 40°)(\sin 40°)$

$= -180\sin^2 40 + 360\sin^2 40°$

$= 180\sin^2 40°$

$\doteqdot 74.4$ m

The greatest height reached by the particle above the ground is about 74.4 m.

2a Initially, $x = y = 0$, and, $\dot{x} = 30\cos 60°$ $\dot{y} = 30\sin 60°$

$\dot{x} = 15$ $\dot{y} = 15\sqrt{3}$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = 15$		When $t = 0$,	$\dot{y} = 15\sqrt{3}$	
	$15 = C_1$			$15\sqrt{3} = C_3$	
so	$\dot{x} = 15$	(2)	so	$\dot{y} = -10t + 15\sqrt{3}$	(5)
Integrating,	$x = 15t + C_2$		Integrating,	$y = -5t^2 + 15t\sqrt{3} + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x = 15t$	(3)	so	$y = -5t^2 + 15t\sqrt{3}$	(6)

Particle reaches greatest height when $\dot{y} = 0$.

Substituting into (5):

$0 = -10t + 15\sqrt{3}$

$10t = 15\sqrt{3}$

© Cambridge University Press 2019

$$t = \frac{3\sqrt{3}}{2}$$

Substituting for t in (6):

$$y = -5\left(\frac{3\sqrt{3}}{2}\right)^2 + 15 \times \frac{3\sqrt{3}}{2} \times \sqrt{3}$$

$$y = -5 \times \frac{27}{4} + 15 \times \frac{9}{2}$$

$$= -33.75 + 67.5$$

$$= 33.75 \text{ m}$$

The greatest height reached by the rock above the point of projection is 33.75 m.

2b The rock will hit the ground when $y = -40$.

Substituting into (6):

$$-40 = -5t^2 + 15t\sqrt{3}$$

$$5t^2 - 15\sqrt{3}t - 40 = 0$$

Using the quadratic formula:

$$t = \frac{15\sqrt{3} \pm \sqrt{\left(15\sqrt{3}\right)^2 - 4 \times 5 \times (-40)}}{2 \times 5}$$

$$t = \frac{15\sqrt{3} \pm \sqrt{1475}}{10}$$

$$t = \frac{15\sqrt{3} \pm 5\sqrt{59}}{10}$$

$$t = \frac{3\sqrt{3} - \sqrt{59}}{2} \text{ or } t = \frac{3\sqrt{3} + \sqrt{59}}{2}$$

$t \div -1.24$ or $t \div 6.44$

Since $t > 0$, the stone hits the ground when $t \div 6.44$ seconds.

When $t = \frac{3\sqrt{3} + \sqrt{59}}{2}$,

$x = 15t$ becomes

© Cambridge University Press 2019

$$x = \frac{15(3\sqrt{3} + \sqrt{59})}{2}$$

$$\div 96.58 \text{ m}$$

The rock lands about 96.58 m from the base of the cliff.

2c The particle's speed can be calculated by applying Pythagoras' theorem to the particle's velocity vector.

That is $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

$$v^2 = \dot{x}^2 + \dot{y}^2$$

$$v = \sqrt{\dot{x}^2 + \dot{y}^2}$$

From (2) and (5):

$$v = \sqrt{15^2 + \left(-10t + 15\sqrt{3}\right)^2}$$

When $t = \frac{3\sqrt{3} + \sqrt{59}}{2}$,

$$v = \sqrt{15^2 + \left(-10\left(\frac{3\sqrt{3} + \sqrt{59}}{2}\right) + 15\sqrt{3}\right)^2}$$

$$= \sqrt{15^2 + \left(-15\sqrt{3} - 5\sqrt{59} + 15\sqrt{3}\right)^2}$$

$$= \sqrt{225 + 1475}$$

$$= \sqrt{1700}$$

$$\div 41.23 \text{ m/s}$$

The rock hits the ground at a speed of about 41.23 m/s.

3a Initially, $x = y = 0$

And $\dot{x} = 25\cos\left(\tan^{-1}\frac{4}{3}\right)$ $\dot{y} = 25\sin\left(\tan^{-1}\frac{4}{3}\right)$

$\dot{x} = 15$ $\dot{y} = 20$

To begin, $\ddot{x} = 0$ (1) To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{x} = C_1$ Integrating, $\dot{y} = -10t + C_2$

© Cambridge University Press 2019

When $t = 0$, $\dot{x} = 15$

$15 = C_1$

so $\dot{x} = 15$ (2)

When $t = 0$, $\dot{y} = 20$

$20 = C_2$

so $\dot{y} = -10t + 20$ (4)

3b To begin, $\dot{x} = 15$

Integrating, $x = 15t + C_3$

When $t = 0$, $x = 0$

$0 = C_3$

so $x = 15t$ (5)

To begin, $\dot{y} = -10t + 20$

Integrating, $y = -5t^2 + 20t + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + 20t$ (6)

3c The particle's distance can be calculated by applying Pythagoras' theorem.

$r^2 = x^2 + y^2$

$r = \sqrt{x^2 + y^2}$

From (5) and (6):

$r = \sqrt{(15t)^2 + (-5t^2 + 20t)^2}$

When $t = 1$,

$r = \sqrt{15^2 + (-5 + 20)^2}$

$= \sqrt{225 + 225}$

$= \sqrt{450}$

$= 15\sqrt{2}$

The distance of the particle from point of projection after one second is $15\sqrt{2}$ m.

3d $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

From (2) and (4):

$\underline{v} = 15\underline{i} + (-10t + 20)\underline{j}$

When $t = 1$,

$\underline{v} = 15\underline{i} + 10\underline{j}$

© Cambridge University Press 2019

$v^2 = 15^2 + 10^2$

$v = \sqrt{15^2 + 10^2}$

$= \sqrt{325}$

$= 5\sqrt{13}$ m/s

$\tan\theta = \frac{10}{15} = \frac{2}{3}$

$\theta = \tan^{-1}\frac{2}{3}$

3e Particle reaches greatest height when $\dot{y} = 0$.

Substituting into (4):

$0 = -10t + 20$

$10t = 20$

$t = 2$

Substituting $t = 2$ into (6):

$y = -20 + 40$

$= 20$ m

So the greatest height, H, is 20 m.

Particle reaches the ground when $y = 0$.

Substituting into (6):

$0 = -5t^2 + 20t$

$5t^2 - 20t = 0$

$5t(t - 4) = 0$

$t = 0$ or $t = 4$

The particle reaches its horizontal range, R, at $t = 4$.

From (5):

$x = 15 \times 4 = 60$, and hence $R = 60$

$\frac{R}{H} = \frac{60}{20} = 3$

© Cambridge University Press 2019

So $R = 3H$.

4a Initially, $x = y = 0$, and, $\dot{x} = 10\cos 45°$ $\dot{y} = 10\sin 45°$

$\dot{x} = 5\sqrt{2}$ $\dot{y} = 5\sqrt{2}$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = 5\sqrt{2}$		When $t = 0$,	$\dot{y} = 5\sqrt{2}$	
	$5\sqrt{2} = C_1$			$5\sqrt{2} = C_3$	
so	$\dot{x} = 5\sqrt{2}$	(2)	so	$\dot{y} = -10t + 5\sqrt{2}$	(5)
Integrating,	$x = 5t\sqrt{2} + C_2$		Integrating,	$y = -5t^2 + 5t\sqrt{2} + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x = 5t\sqrt{2}$	(3)	so	$y = -5t^2 + 5t\sqrt{2}$	(6)

From (3):

$$t = \frac{x}{5\sqrt{2}}$$

Substituting for t in (6):

$$y = -5\left(\frac{x}{5\sqrt{2}}\right)^2 + \frac{5x\sqrt{2}}{5\sqrt{2}}$$

$$y = \frac{-5x^2}{50} + x$$

$$y = x - \frac{1}{10}x^2 \qquad (7)$$

© Cambridge University Press 2019

4b Particle reaches the ground when $y = 0$.

Substituting into (7):

$$0 = x - \frac{x^2}{10}$$

$$x^2 - 10x = 0$$

$$x(x - 10) = 0$$

$x = 0$ or $x = 10$

For horizontal range, $x > 0$, so $x = 10$ m.

By symmetry, the greatest height is reached at half the horizontal range

so, $x = 5$

From (7):

$$y = 5 - \frac{25}{10}$$

$$= 2.5 \text{ m}$$

Hence the horizontal range is 10 m and the greatest height is 2.5 m.

4c i From (7):

$$y = x - \frac{x^2}{10}$$

When $x = 8$,

$$y = 8 - \frac{64}{10}$$

$$= 1.6 \text{ m}$$

The stone hits the wall at a height of 1.6 m.

4c ii From (7):

$$\frac{dy}{dx} = 1 - \frac{x}{5}$$

When $x = 8$,

$$\frac{dy}{dx} = -\frac{3}{5}$$

© Cambridge University Press 2019

$$\tan\theta = -\frac{3}{5}$$

$$\theta = \tan^{-1}\left(-\frac{3}{5}\right)$$

The direction of the object when it hits the wall is $\tan^{-1}\frac{3}{5}$ below the horizontal.

4d i From (7):

$$y = x - \frac{x^2}{10}$$

When $y = 2.1$,

$$2.1 = x - \frac{x^2}{10}$$

$$\frac{x^2}{10} - x + 2.1 = 0$$

$$x^2 - 10x + 21 = 0$$

$$(x-3)(x-7) = 0$$

So $x = 3$ or $x = 7$

It must be the case that $x = 3$ m since the object collided with the ceiling when travelling up.

4d ii From (7):

$$\frac{dy}{dx} = 1 - \frac{x}{5}$$

When $x = 3$,

$$\frac{dy}{dx} = \frac{2}{5}$$

$$\tan\theta = \frac{2}{5}$$

$$\theta = \tan^{-1}\frac{2}{5}$$

The angle at which the stone hits the ceiling is $\tan^{-1}\frac{2}{5}$ above the horizontal.

© Cambridge University Press 2019

5a Initially, $x = y = 0$, and, $\dot{x} = 24$ $\dot{y} = 18$

To begin, $\ddot{x} = 0$ (1) To begin, $\ddot{y} = -10$ (3)

Integrating, $\dot{x} = C_1$ Integrating, $\dot{y} = -10t + C_2$

When $t = 0$, $\dot{x} = 24$ When $t = 0$, $\dot{y} = 18$

$24 = C_1$ $18 = C_2$

so $\dot{x} = 24$ (2) so $\dot{y} = -10t + 18$ (4)

To begin, $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

From (2) and (5): $\underline{v} = 24\underline{i} + (-10t + 18)\underline{j} = 24\underline{i} + (18 - 10t)\underline{j}$

5b To begin, $\dot{x} = 24$ To begin, $\dot{y} = -10t + 18$

Integrating, $x = 24t + C_3$ Integrating, $y = -5t^2 + 18t + C_4$

When $t = 0$, $x = 0$ When $t = 0$, $y = 0$

$0 = C_3$ $0 = C_4$

so $x = 24t$ (5) so $y = -5t^2 + 18t$ (6)

To begin, $\underline{r} = x\underline{i} + y\underline{j}$

From (5) and (6): $\underline{r} = (24t)\underline{i} + (-5t^2 + 18t)\underline{j}$

$\underline{r} = (24t)\underline{i} + (18t - 5t^2)\underline{j}$

5c i When $t = 0$,

$\underline{v} = 24\underline{i} + 18\underline{j}$

$v^2 = 24^2 + 18^2$

$v = \sqrt{24^2 + 18^2}$

$= \sqrt{900}$

$= 30$ m/s

The initial speed of the particle is 30 m/s.

© Cambridge University Press 2019

5c ii When $t = 4$,

$\underline{r} = 96\underline{i} + (72 - 80)\underline{j}$

$\underline{r} = 96\underline{i} - 8\underline{j}$

5c iii Particle reaches greatest height when $\dot{y} = 0$.

From (4):

$0 = -10t + 18$

$10t = 18$

$t = 1.8$

When $t = 1.8$,

$\underline{r} = 43.2\underline{i} + (32.4 - 16.2)\underline{j}$

$\underline{r} = 43.2\underline{i} + 16.2\underline{j}$

6a Initially, $x = y = 0$, and, $\dot{x} = V\cos\alpha$ $\dot{y} = V\sin\alpha$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = V\cos\alpha$		When $t = 0$,	$\dot{y} = V\sin\alpha$	
	$V\cos\alpha = C_1$			$V\sin\alpha = C_3$	
so	$\dot{x} = V\cos\alpha$	(2a)	so	$\dot{y} = -10t + V\sin\alpha$	(5a)
Integrating,	$x = Vt\cos\alpha + C_2$		Integrating,	$y = -5t^2 + Vt\sin\alpha + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x = Vt\cos\alpha$	(3a)	so	$y = -5t^2 + Vt\sin\alpha$	(6a)
When $t = 2$,	$24\sqrt{5} = 2V\cos\alpha$		When $t = 2$,	$28 = -20 + 2V\sin\alpha$	
so	$V\cos\alpha = 12\sqrt{5}$		so	$V\sin\alpha = 24$	
From (2a):	$\dot{x} = 12\sqrt{5}$	(2b)	From (5a):	$\dot{y} = -10t + 24$	(5b)
From (3a):	$x = 12t\sqrt{5}$	(3b)	From (6a):	$y = -5t^2 + 24t$	(6b)

© Cambridge University Press 2019

To begin, $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

From (2b) and (5b):

$\underline{v} = 12\sqrt{5}\underline{i} + (-10t + 24)\underline{j}$

When $t = 0$,

$\underline{v} = 12\sqrt{5}\underline{i} + 24\underline{j}$

6b To begin, $\underline{r} = x\underline{i} + y\underline{j}$

From (3b) and (6b):

$\underline{r} = 12t\sqrt{5}\underline{i} + (-5t^2 + 24t)\underline{j}$

When $t = 3$,

$\underline{r} = 36\sqrt{5}\underline{i} + (-45 + 72)\underline{j}$

$\underline{r} = 36\sqrt{5}\underline{i} + 27\underline{j}$

6c The ball will be rising if $\dot{y} > 0$ or falling if $\dot{y} < 0$.

From (5b):

$\dot{y} = -10t + 24$

When $t = 3$,

$\dot{y} = -30 + 24$

$\dot{y} = -6$

Since $\dot{y} = -6 < 0$, the ball must be falling.

7a Initially, $x = y = 0$, and, $\dot{x} = V\cos\theta$ $\dot{y} = V\sin\alpha$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = V\cos\theta$		When $t = 0$,	$\dot{y} = V\sin\theta$	
	$V\cos\theta = C_1$			$V\sin\theta = C_3$	
so	$\dot{x} = V\cos\theta$	(2a)	so	$\dot{y} = -10t + V\sin\theta$	(5a)

© Cambridge University Press 2019

Integrating,	$x = Vt\cos\theta + C_2$		Integrating,	$y = -5t^2 + Vt\sin\theta + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x = Vt\cos\theta$	(3a)	so	$y = -5t^2 + Vt\sin\theta$	(6a)
When $t = 3$,	$108 = 3V\cos\theta$		When $t = 3$,	$0 = -45 + 3V\sin\theta$	
so	$V\cos\theta = 36$		so	$V\sin\theta = 15$	
From (2a):	$\dot{x} = 36$	(2b)	From (5a):	$\dot{y} = -10t + 15$	(5b)
From (3a):	$x = 36t$	(3b)	From (6a):	$y = -5t^2 + 15t$	(6b)

To begin, $\underline{v} = \dot{x}\underline{i} + \dot{y}\underline{j}$

From (2b) and (5b):

$\underline{v} = 36\underline{i} + (-10t + 15)\underline{j}$

When $t = 0$,

$\underline{v} = 36\underline{i} + 15\underline{j}$

$V^2 = 36^2 + 15^2$

$V = \sqrt{36^2 + 15^2}$

$= \sqrt{1521}$

$= 39$ m/s

$\tan\theta = \dfrac{15}{36} = \dfrac{5}{12}$

$\theta = \tan^{-1}\dfrac{5}{12} = \text{artan}\ \dfrac{5}{12}$

7b Particle reaches greatest height when $\dot{y} = 0$.

Substituting into (5b):

$0 = -10t + 15$

$10t = 15$

$t = 1.5$

Substituting for t in (6b):

© Cambridge University Press 2019

$y = -11.25 + 22.5$

$= 11.25$ m

The greatest height reached by the particle is 11.25 m.

8a $V = 5$ m/s and $\tan\alpha = 2$ or $\alpha = \tan^{-1} 2$

For a right-angled triangle with angle α, opposite side is 2 units, adjacent side is 1 unit and hypotenuse is $\sqrt{2^2 + 1^2} = \sqrt{5}$. Hence $\cos\alpha = \frac{1}{\sqrt{5}}$ and $\sin\alpha = \frac{2}{\sqrt{5}}$.

$\dot{x} = 5\cos\alpha$


$= 5 \times \frac{1}{\sqrt{5}}$

$= \sqrt{5}$ m/s

$\dot{y} = 5\sin\alpha$

$= 5 \times \frac{2}{\sqrt{5}}$

$= 2\sqrt{5}$ m/s



So $\underline{v} = \sqrt{5}\underline{i} + 2\sqrt{5}\underline{j}$ m/s

Hence, the initial values are $\dot{x} = \sqrt{5}$ and $\dot{y} = 2\sqrt{5}$.

8b Initially, $x = y = 0$, and, $\dot{x} = \sqrt{5}$ $\dot{y} = 2\sqrt{5}$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = \sqrt{5}$		When $t = 0$,	$\dot{y} = 2\sqrt{5}$	
	$\sqrt{5} = C_1$			$2\sqrt{5} = C_3$	
so	$\dot{x} = \sqrt{5}$	(2)	so	$\dot{y} = -10t + 2\sqrt{5}$	(5)
Integrating,	$x = \sqrt{5}t + C_2$		Integrating,	$y = -5t^2 + 2t\sqrt{5} + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
	$0 = C_2$			$0 = C_4$	
so	$x = t\sqrt{5}$	(3)	so	$y = -5t^2 + 2t\sqrt{5}$	(6)

© Cambridge University Press 2019

8c From (6):

$y = -5t^2 + 2t\sqrt{5}$

When $t = 1$,

$y = -5 + 2\sqrt{5}$

$\div -0.53$

At one second, the apple would be below the height at which it was thrown. However, since the apple is caught at the same height that it was thrown, it must have been caught before one second. Therefore, it spent less than one second in the air.

8d The apple reaches greatest height when $\dot{y} = 0$.

Substituting into (5):

$0 = -10t + 2\sqrt{5}$

$10t = 2\sqrt{5}$

$t = \frac{\sqrt{5}}{5}$

Substituting for t in (6):

$y = -5\left(\frac{\sqrt{5}}{5}\right)^2 + 2\sqrt{5} \times \frac{\sqrt{5}}{5}$

$= -1 + 2$

$= 1$

The greatest height of the apple above the point of release is 1 metre.

8e The apple lands when $y = 0$.

Substituting into (6):

$0 = -5t^2 + 2t\sqrt{5}$

$5t^2 - 2\sqrt{5}t = 0$

$t(5t - 2\sqrt{5}) = 0$

$t = 0$ or $t = \frac{2\sqrt{5}}{5}$ s

© Cambridge University Press 2019

For the flight of the apple,

$$t = \frac{2\sqrt{5}}{5} = 2\frac{\sqrt{5}}{5} \text{ s}$$

Substituting for t in (3):

$$x = \sqrt{5} \times \frac{2\sqrt{5}}{5}$$


$$x = 2 \text{ m}$$

Hence the horizontal distance travelled by the apple is 2 m.

8f Adam catches the apple at $t = \frac{2\sqrt{5}}{5}$ seconds.

From (2): $\dot{x} = \sqrt{5}$

Substituting for t in (5):



$$\dot{y} = -10 \times \frac{2\sqrt{5}}{5} + 2\sqrt{5}$$

$$\dot{y} = -4\sqrt{5} + 2\sqrt{5}$$

$$\dot{y} = -2\sqrt{5}$$

$$v^2 = \left(\sqrt{5}\right)^2 + \left(-2\sqrt{5}\right)^2$$

$$v = \sqrt{5 + 20}$$

$$v = 5 \text{ m/s}$$

So, the final speed is equal to the initial speed.

$$\tan\theta = \frac{-2\sqrt{5}}{\sqrt{5}}$$

$$\theta = \tan^{-1}\left(\frac{-2\sqrt{5}}{\sqrt{5}}\right)$$

$$\theta = \tan^{-1}(-2)$$

$$\theta = -\tan^{-1} 2$$

The final and initial angles of inclination are equal in magnitude, though it is an angle of inclination initially, and an angle of depression finally.

© Cambridge University Press 2019

8g From (3):

$$t = \frac{x}{\sqrt{5}}$$

Substituting for t in (6):

$$y = -5\left(\frac{x}{\sqrt{5}}\right)^2 + 2 \times \frac{x}{\sqrt{5}} \times \sqrt{5}$$

$$= -x^2 + 2x$$

$$= 2x - x^2$$

9 Let V be the initial speed of the projectile

Initially, $x = y = 0$, and, $\dot{x} = V\cos 26°$ $\dot{y} = V\sin 26°$

To begin,	$\ddot{x} = 0$	(1)	To begin,	$\ddot{y} = -10$	(4)
Integrating,	$\dot{x} = C_1$		Integrating,	$\dot{y} = -10t + C_3$	
When $t = 0$,	$\dot{x} = V\cos 26°$		When $t = 0$,	$\dot{y} = V\sin 26°$	
$V\cos 26° = C_1$				$V\sin 26° = C_3$	
so	$\dot{x} = V\cos 26°$	(2)	so	$\dot{y} = -10t + V\sin 26°$	(5)
To begin,	$\dot{x} = V\cos 26°$		To begin,	$\dot{y} = -10t + V\sin 26°$	
Integrating,	$x = Vt\cos 26° + C_2$		Integrating,	$y = -5t^2 + Vt\sin 26° + C_4$	
When $t = 0$,	$x = 0$		When $t = 0$,	$y = 0$	
$0 = C_2$				$0 = C_4$	
so	$x = Vt\cos 26°$	(3)	so	$y = -5t^2 + Vt\sin 26°$	(6)

From (3):

$$t = \frac{x}{V\cos 26°}$$

Substituting for t in (6):

$$y = -5\left(\frac{x}{V\cos 26°}\right)^2 + \left(\frac{Vx}{V\cos 26°}\right)\sin 26°$$

$$= \frac{-5x^2}{V^2\cos^2 26°} + x\tan 26°$$

If the ball just clears the tree, it will reach a point where $x = 30$ and $y = 12$.

© Cambridge University Press 2019

$$12 = \frac{-5(30)^2}{V^2 \cos^2 26°} + 30 \tan 26°$$

$$\frac{-4500}{V^2 \cos^2 26°} = 12 - 30 \tan 26°$$

$$-\frac{4500}{V^2} = \cos^2 26° \times (12 - 30 \tan 26°)$$

$$V^2 = -\frac{4500}{\cos^2 26° \times (12 - 30 \tan 26°)}$$

$$V = \sqrt{-\frac{4500}{\cos^2 26° \times (12 - 30 \tan 26°)}}$$

$V \doteqdot 46$ m/s

The initial speed of the ball is about 46 m/s.

10a Initially, $x = y = 0$, and, $\dot{x} = V \cos\theta$ $\dot{y} = V \sin\theta$

Assuming $x = Vt \cos\theta$ (1)

$y = -\frac{1}{2} g t^2 + Vt \sin\theta$ (2)

From (1):

$$t = \frac{x}{V \cos\theta}$$

Substituting for t in (2):

$$y = -\frac{1}{2} g \left(\frac{x}{V \cos\theta}\right)^2 + \frac{Vx \sin\theta}{V \cos\theta}$$

$$= -\frac{gx^2}{2V^2 \cos^2\theta} + x \tan\theta$$

$$= -\frac{gx^2 \sec^2\theta}{2V^2} + x \tan\theta$$

$$= x \tan\theta - \frac{gx^2 \sec^2\theta}{2V^2} \quad (3)$$

© Cambridge University Press 2019

10b If initially $V = 25$, then the particle passes through the point $x = 30$ and $y = 6$.

Substituting $x = 30$, $y = 6$ and $g = 10$ in (3):

$$6 = 30\tan\theta - \frac{10 \times 30^2 \sec^2\theta}{2 \times 25^2}$$

$$6 = 30\tan\theta - \frac{9000\sec^2\theta}{1250}$$

$$6 = 30\tan\theta - \frac{36\sec^2\theta}{5}$$

$$30 = 150\tan\theta - 36\sec^2\theta$$

$$36\sec^2\theta - 150\tan\theta + 30 = 0$$

$$6\sec^2\theta - 25\tan\theta + 5 = 0$$

But $\sec^2\theta = 1 + \tan^2\theta$

$$6(1 + \tan^2\theta) - 25\tan\theta + 5 = 0$$

$$6 + 6\tan^2\theta - 25\tan\theta + 5 = 0$$

$$6\tan^2\theta - 25\tan\theta + 11 = 0$$

$$(2\tan\theta - 1)(3\tan\theta - 11) = 0$$

$$\tan\theta = \frac{1}{2} \text{ or } \tan\theta = \frac{11}{3}$$

$$\text{So } \theta = \tan^{-1}\frac{1}{2} \text{ or } \theta = \tan^{-1}\frac{11}{3}$$

$$\text{or } \theta = \arctan\frac{1}{2} \text{ or } \theta = \arctan\frac{11}{3}$$

© Cambridge University Press 2019

11a Initially, $x = y = 0$, and, $\dot{x} = 15\cos 60°$ $\dot{y} = 15\sin 60°$

$\dot{x} = \frac{15}{2}$ $\dot{y} = \frac{15\sqrt{3}}{2}$

To begin, $\ddot{x} = 0$ (1)

Integrating, $\dot{x} = C_1$

When $t = 0$, $\dot{x} = \frac{15}{2}$

$\frac{15}{2} = C_1$

so $\dot{x} = \frac{15}{2}$ (2)

Integrating, $x = \frac{15}{2}t + C_2$

When $t = 0$, $x = 0$

$0 = C_2$

so $x = \frac{15}{2}t$ (3)

To begin, $\ddot{y} = -10$ (4)

Integrating, $\dot{y} = -10t + C_3$

When $t = 0$, $\dot{y} = \frac{15\sqrt{3}}{2}$

$\frac{15\sqrt{3}}{2} = C_3$

so $\dot{y} = -10t + \frac{15\sqrt{3}}{2}$ (5)

Integrating, $y = -5t^2 + \frac{15\sqrt{3}}{2}t + C_4$

When $t = 0$, $y = 0$

$0 = C_4$

so $y = -5t^2 + \frac{15\sqrt{3}}{2}t$ (6)

11b From (3):

$t = \frac{2x}{15}$

Substituting for t in (6):

$$y = -5\left(\frac{2x}{15}\right)^2 + \frac{15\sqrt{3}}{2} \times \frac{2x}{15}$$

$$= -\frac{20x^2}{225} + \sqrt{3}x$$

$$= -\frac{4}{45}x^2 + \sqrt{3}x \quad (7)$$

11c i The ball will land when the equation of its path intersects with the equation of the slope.

$\theta = 30°$

$\tan\theta = \frac{\sqrt{3}}{3}$

© Cambridge University Press 2019

So the gradient is $m = \frac{\sqrt{3}}{3}$.

Hence $y = \frac{\sqrt{3}}{3}x$ is the equation of the slope.

Equating with (7):

$$\frac{\sqrt{3}}{3}x = -\frac{4}{45}x^2 + \sqrt{3}x$$

$$\frac{4}{45}x^2 = \sqrt{3}x - \frac{\sqrt{3}}{3}x$$

$$\frac{4}{45}x = \sqrt{3} - \frac{\sqrt{3}}{3} \qquad \text{assuming } x \neq 0$$

$$\frac{4}{45}x = \frac{3\sqrt{3} - \sqrt{3}}{3}$$

$$x = \frac{45}{4} \times \frac{2\sqrt{3}}{3}$$

$$x = \frac{15\sqrt{3}}{2}$$

When $x = \frac{15\sqrt{3}}{2}$,

$$y = \frac{15\sqrt{3}}{2} \times \frac{\sqrt{3}}{3}$$

$$y = \frac{15}{2}$$

Using Pythagoras' theorem to calculate the distance up the slope:

$$r^2 = \left(\frac{15\sqrt{3}}{2}\right)^2 + \left(\frac{15}{2}\right)^2$$

$$r = \sqrt{\frac{675}{4} + \frac{225}{4}}$$

$$= \sqrt{225}$$

$$= 15 \text{ m}$$

The ball landed 15 m up the hill.

© Cambridge University Press 2019

11c ii From (3):

$$t = \frac{2x}{15}$$

When $x = \frac{15\sqrt{3}}{2}$,

$$t = \frac{2}{15} \times \frac{15\sqrt{3}}{2}$$

$$= \sqrt{3} \text{ seconds}$$

The time of the flight is $\sqrt{3}$ seconds.

© Cambridge University Press 2019