Chapter 13 worked solutions – Differential equations

Solutions to Exercise 13A

Let the integration constants be A, B, C or D.

1a $y' - y = x$

first-order differential equation

1b $y'y = 3x$

first-order differential equation

1c $y'' + 4y' - y = \sin x$

second-order differential equation

1d $y' + y\cos x = e^x$

first-order differential equation

1e $y'' - \frac{1}{2}(y')^2 = 0$

second-order differential equation

1f $y' + y^2 = 1$

first-order differential equation

1g $y' + xy = 0$

first-order differential equation

1h $xy'' + y' = x^2$

second-order differential equation

1i $y'' - xy' + e^x y = 0$

second-order differential equation

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

2a linear

2b non-linear

2d linear

2f linear

2g linear

3a one arbitrary constant

3b one arbitrary constant

3c two arbitrary constants

3d one arbitrary constant

3e two arbitrary constants

3f one arbitrary constant

3g one arbitrary constant

3h two arbitrary constants

3i two arbitrary constants

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4a $y = 5x^3$

$y' = 15x^2$

Substituting for y and y' in $xy' - 3y = 0$:

LHS $= x(15x^2) - 3(5x^3)$

$= 15x^3 - 15x^3$

$= 0$

$=$ RHS

Therefore, $y = 5x^3$ is a solution of $xy' - 3y = 0$.

4b $y = x^2 - 1$

$y' = 2x$

Substituting for y and y' in $xy' - 2y = 2$:

LHS $= x(2x) - 2(x^2 - 1)$

$= 2x^2 - 2x^2 + 2$

$= 2$

$=$ RHS

Therefore, $y = x^2 - 1$ is a solution of $xy' - 2y = 2$.

4c $y = 3e^{-x}$

$y' = -3e^{-x}$

Substituting for y and y' in $y' + y = 0$:

LHS $= -3e^{-x} + 3e^{-x}$

$= 0$

$=$ RHS

Therefore, $y = 3e^{-x}$ is a solution of $y' + y = 0$.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4d $y = \sqrt{x^2 + 4}$

$= (x^2 + 4)^{\frac{1}{2}}$

$y' = \frac{1}{2}(x^2 + 4)^{-\frac{1}{2}} \times 2x$

$= x(x^2 + 4)^{-\frac{1}{2}}$

Substituting for y and y' in $y'y = x$:

LHS $= x(x^2 + 4)^{-\frac{1}{2}} \times (x^2 + 4)^{\frac{1}{2}}$

$= x(x^2 + 4)^0$

$= x \times 1$

$= x$

$=$ RHS

Therefore $y = \sqrt{x^2 + 4}$ is a solution of $y'y = x$.

5a $y = \int (2x - 3)dx$

$= \frac{2x^2}{2} - 3x + C$

$= x^2 - 3x + C$

5b $y = \int (12e^{-2x} + 4)dx$

$= -\frac{12}{2}e^{-2x} + 4x + C$

$= -6e^{-2x} + 4x + C$

5c $y = \int (\sec^2 x)dx$

$= \tan x + C$

5d $y = \int (6\cos 2x + 9\sin 3x)\,dx$

$= \frac{6}{2}\sin 2x - \frac{9}{3}\cos 3x + C$

$= 3\sin 2x - 3\cos 3x + C$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

5e $y = \int \sqrt{1-5x}\ dx$

$= \int (1-5x)^{\frac{1}{2}}\ dx$

$= \dfrac{(1-5x)^{\frac{3}{2}}}{(-5) \times \left(\frac{3}{2}\right)} + C$

$= \dfrac{(1-5x)^{\frac{3}{2}}}{\left(-\frac{15}{2}\right)} + C$

$= -\dfrac{2(1-5x)^{\frac{3}{2}}}{15} + C$

5f $y = \int 4x \cos x^2\ dx$

Let $u = x^2$, $\frac{du}{dx} = 2x$ so $\frac{du}{2x} = dx$

$y = \int 4x \cos u\ \frac{du}{2x}$

$= \int 2 \cos u\ du$

$= 2\int \cos u\ du$

$= 2 \sin u + C$

$= 2 \sin x^2 + C$

6a $y = Ce^x - x - 1$

$y' = Ce^x - 1$

Substituting for y and y' in $y' = x + y$:

LHS $= Ce^x - 1$

RHS $= x + Ce^x - x - 1$

$= Ce^x - 1$

$=$ LHS

Therefore $y = Ce^x - x - 1$ is a solution of $y' = x + y$.

© Cambridge University Press 2019

6b $y = Cxe^{-x}$

$y' = C(e^{-x} \times 1 + x \times -e^{-x})$

$y' = C(e^{-x} - xe^{-x})$

Substituting for y and y' in $xy' = y(1 - x)$:

LHS $= x(C(e^{-x} - xe^{-x}))$

$= Cx(e^{-x} - xe^{-x})$

$= Cxe^{-x}(1 - x)$

$= y(1 - x)$

$=$ RHS

Therefore $y = Cxe^{-x}$ is a solution of $xy' = y(1 - x)$.

6c $y = \sin\ (x + C)$

$y' = \cos\ (x + C)$

Substituting for y and y' in $(y')^2 = 1 - y^2$:

LHS $= (\cos\ (x + C))^2$

$= \cos^2\ (x + C)$

RHS $= 1 - \sin^2\ (x + C)$

$= \cos^2\ (x + C)$

$=$ LHS

Therefore $y = \sin\ (x + C)$ is a solution of $(y')^2 = 1 - y^2$.

6d

$$y = \frac{C}{x} + 2$$

$$= Cx^{-1} + 2$$

$$\frac{dy}{dx} = -Cx^{-2}$$

$$= -\frac{C}{x^2}$$

Substituting for y and $\frac{dy}{dx}$ in $\frac{dy}{dx} = \frac{2-y}{x}$:

© Cambridge University Press 2019

$$\text{RHS} = \frac{2 - \left(\frac{C}{x} + 2\right)}{x}$$

$$= \frac{-\frac{C}{x}}{x}$$

$$= -\frac{C}{x^2}$$

$$= \frac{dy}{dx}$$

$$= \text{LHS}$$

Therefore $y = \frac{C}{x} + 2$ is a solution of $\frac{dy}{dx} = \frac{2-y}{x}$.

7a $y = x^2 - 2x + 3$

$y' = 2x - 2$

$y'' = 2$

Substituting for y, y' and y'' in $x^2y'' - 2xy' + 2y = 6$:

$\text{LHS} = x^2(2) - 2x(2x - 2) + 2(x^2 - 2x + 3)$

$= 2x^2 - 4x^2 + 4x + 2x^2 - 4x + 6$

$= 6$

$= \text{RHS}$

Therefore $y = x^2 - 2x + 3$ is a solution of $x^2y'' - 2xy' + 2y = 6$.

7b $y = 2e^x + e^{5x}$

$y' = 2e^x + 5e^{5x}$

$y'' = 2e^x + 25e^{5x}$

Substituting for y, y' and y'' in $y'' - 6y' + 5y = 0$:

$\text{LHS} = 2e^x + 25e^{5x} - 6(2e^x + 5e^{5x}) + 5(2e^x + e^{5x})$

$= 2e^x + 25e^{5x} - 12e^x - 30e^{5x} + 10e^x + 5e^{5x}$

$= 0$

$= \text{RHS}$

© Cambridge University Press 2019

Therefore $y = 2e^x + e^{5x}$ is a solution of $y'' - 6y' + 5y = 0$.

7c $y = \cos \pi x - 3 \sin \pi x$

$y' = -\pi \sin \pi x - 3\pi \cos \pi x$

$y'' = -\pi^2 \cos \pi x + 3\pi^2 \sin \pi x$

Substituting for y and y'' in $y'' + \pi^2 y = 0$:

$\text{LHS} = -\pi^2 \cos \pi x + 3\pi^2 \sin \pi x + \pi^2(\cos \pi x - 3 \sin \pi x)$

$= -\pi^2 \cos \pi x + 3\pi^2 \sin \pi x + \pi^2 \cos \pi x - 3\pi^2 \sin \pi x$

$= 0$

$= \text{RHS}$

Therefore $y = \cos \pi x - 3 \sin \pi x$ is a solution of $y'' + \pi^2 y = 0$.

7d $y = e^{-2x} \sin x$

$y' = (\sin x)(-2e^{-2x}) + (e^{-2x})(\cos x)$

$= -2e^{-2x} \sin x + e^{-2x} \cos x$

$y'' = (\sin x)(4e^{-2x}) + (-2e^{-2x})(\cos x) + (\cos x)(-2e^{-2x}) + (e^{-2x})(-\sin x)$

$= 4e^{-2x} \sin x - 2e^{-2x} \cos x - 2e^{-2x} \cos x - e^{-2x} \sin x$

$= 3e^{-2x} \sin x - 4e^{-2x} \cos x$

Substituting for y, y' and y'' in $y'' + 4y' + 5y = 0$:

$\text{LHS} = 3e^{-2x} \sin x - 4e^{-2x} \cos x + 4(-2e^{-2x} \sin x + e^{-2x} \cos x) + 5(e^{-2x} \sin x)$

$= 3e^{-2x} \sin x - 4e^{-2x} \cos x - 8e^{-2x} \sin x + 4e^{-2x} \cos x + 5e^{-2x} \sin x$

$= 0$

$= \text{RHS}$

Therefore $y = e^{-2x} \sin x$ is a solution of $y'' + 4y' + 5y = 0$.

© Cambridge University Press 2019

7e $y = \cos(\log_e x)$

$$y' = -\sin(\log_e x) \times \frac{1}{x}$$

$$= -\frac{\sin(\log_e x)}{x}$$

$$y'' = -\frac{x \times (\cos(\log_e x) \times \frac{1}{x} - \sin(\log_e x)(1)}{x^2}$$

$$= -\frac{\cos(\log_e x) - \sin(\log_e x)}{x^2}$$

Substituting for $y,\ y'$ and y'' in $x^2y'' + xy' + y = 0$:

$$\text{LHS} = x^2\left(-\frac{\cos(\log_e x) - \sin(\log_e x)}{x^2}\right) + x\left(-\frac{\sin(\log_e x)}{x}\right) + \cos(\log_e x)$$

$$= -\cos(\log_e x) + \sin(\log_e x) - \sin(\log_e x) + \cos(\log_e x)$$

$$= 0$$

$$= \text{RHS}$$

Therefore $y = \cos(\log_e x)$ is a solution of $x^2y'' + xy' + y = 0$.

8a $y'' = 2$

$$y' = \int 2\, dx = 2x + A$$

$$y = \int (2x + A)\, dx = \frac{2x^2}{2} + A\,x + B = x^2 + Ax + B$$

8b $y'' = \cos 2x$

$$y' = \int \cos 2x\ dx = \frac{1}{2}\sin 2x + A$$

$$y = \int \left(\frac{1}{2}\sin 2x + A\right) dx = -\frac{1}{4}\cos 2x + A\,x + B$$

8c $y'' = e^{\frac{1}{2}x}$

$$y' = \int e^{\frac{1}{2}x}\, dx = 2e^{\frac{1}{2}x} + A$$

$$y = \int \left(2e^{\frac{1}{2}x} + A\right) dx = 4e^{\frac{1}{2}x} + A\,x + B$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

8d $y'' = \sec^2 x$

$y' = \int \sec^2 x \, dx = \tan x + A$

$y = \int (\tan x + A) \, dx$

$y = \int \left(\frac{\sin x}{\cos x} + A\right) dx$

Using substitution:

Let $u = \cos x$

$\frac{du}{dx} = -\sin x$

$dx = -\frac{du}{\sin x}$

$y = \int \left(\frac{\sin x}{\cos x} + A\right) dx$

$= \int \frac{\sin x}{\cos x} \, dx + \int A \, dx$

$= \int \frac{\sin x}{u}\left(-\frac{du}{\sin x}\right) + Ax + B$

$= -\int \frac{1}{u} \, du + Ax + B$

$= -\log_e|u| + Ax + B$

$= -\log_e|\cos x| + Ax + B$

9a For $y = e^{-x}$

$y' = -e^{-x}$

$y'' = e^{-x}$

Substituting y, y' and y'' into $y'' - 2y' - 3y = 0$:

LHS $= e^{-x} - 2(-e^{-x}) - 3e^{-x}$

$= e^{-x} + 2e^{-x} - 3e^{-x}$

$= 0$

$=$ RHS

Therefore $y = e^{-x}$ is a solution of $y'' - 2y' - 3y = 0$.

© Cambridge University Press 2019

For $y = e^{3x}$

$y' = 3e^{3x}$

$y'' = 9e^{3x}$

Substituting y, y' and y'' into $y'' - 2y' - 3y = 0$:

LHS $= 9e^{3x} - 2(3e^{3x}) - 3e^{3x}$

$= 9e^{3x} - 6e^{3x} - 3e^{3x}$

$= 0$

$=$ RHS

Therefore $y = e^{3x}$ is a solution of $y'' - 2y' - 3y = 0$.

9b $y = Ae^{-x} + Be^{3x}$

$y' = -Ae^{-x} + 3Be^{3x}$

$y'' = Ae^{-x} + 9Be^{3x}$

Substituting y, y' and y'' into $y'' - 2y' - 3y = 0$

LHS $= Ae^{-x} + 9Be^{3x} - 2(-Ae^{-x} + 3Be^{3x}) - 3(Ae^{-x} + Be^{3x})$

$= Ae^{-x} + 9Be^{3x} + 2Ae^{-x} - 6Be^{3x} - 3Ae^{-x} - 3Be^{3x}$

$= 0$

$=$ RHS

Therefore $y = Ae^{-x} + Be^{3x}$ is a solution of $y'' - 2y' - 3y = 0$.

10a $y = x^3 + Ax^2 + Bx + C$

$y' = 3x^2 + 2Ax + B$

$y'' = 6x + 2A$

$y''' = 6$

Substituting y''' into $y''' = 6$:

LHS $= 6$

$=$ RHS

Therefore $y = x^3 + Ax^2 + Bx + C$ is a solution of $y''' = 6$.

© Cambridge University Press 2019

10b $y = Ae^{-x} + Be^{-2x} + 2x - 3$

$y' = -Ae^{-x} - 2Be^{-2x} + 2$

$y'' = Ae^{-x} + 4Be^{-2x}$

Substituting y, y' and y'' into $y'' + 3y' + 2y = 4x$:

$\text{LHS} = Ae^{-x} + 4Be^{-2x} + 3(-Ae^{-x} - 2Be^{-2x} + 2) + 2(Ae^{-x} + Be^{-2x} + 2x - 3)$

$= Ae^{-x} + 4Be^{-2x} - 3Ae^{-x} - 6Be^{-2x} + 6 + 2Ae^{-x} + 2Be^{-2x} + 4x - 6$

$= (Ae^{-x} - 3Ae^{-x} + 2Ae^{-x}) + (4Be^{-2x} - 6Be^{-2x} + 2Be^{-2x}) + 6 + 4x - 6$

$= 4x$

$= \text{RHS}$

Therefore $y = Ae^{-x} + Be^{-2x} + 2x - 3$ is a solution of $y'' + 3y' + 2y = 4x$.

10c $y = A\cos 2x + B\sin 2x$

$y' = -2A\sin 2x + 2B\cos 2x$

$y'' = -4A\cos 2x - 4B\sin 2x$

Substituting y and y'' into $y'' + 4y = 0$:

$\text{LHS} = -4A\cos 2x - 4B\sin 2x + 4(A\cos 2x + B\sin 2x)$

$= -4A\cos 2x - 4B\sin 2x + 4A\cos 2x + 4B\sin 2x$

$= 0$

$= \text{RHS}$

Therefore $y = A\cos 2x + B\sin 2x$ is a solution of $y'' + 4y = 0$.

10d $y = A\,e^{-x}\cos x$

$y' = A(\cos x \times (-e^{-x}) + e^{-x} \times (-\sin x))$

$= A(-e^{-x}\cos x - e^{-x}\sin x)$

$= -Ae^{-x}\cos x - Ae^{-x}\sin x$

$y'' = \text{A}\,((e^{-x}\cos x + e^{-x}\sin x) + e^{-x}\sin x - e^{-x}\cos x)$

$= \text{A}\,e^{-x}\cos x + Ae^{-x}\sin x + Ae^{-x}\sin x - A\,e^{-x}\cos x$

$= 2Ae^{-x}\sin 2x$

Substituting y, y' and y'' into $y'' + 2y' + 2y = 0$:

© Cambridge University Press 2019

$\text{LHS} = 2Ae^{-x}\sin 2x + 2(-Ae^{-x}\cos x - Ae^{-x}\sin x) + 2A\,e^{-x}\cos x$

$= 2Ae^{-x}\sin 2x - 2Ae^{-x}\cos x - 2\,Ae^{-x}\sin x + 2A\,e^{-x}\cos x$

$= 0$

$= \text{RHS}$

Therefore $y = A\,e^{-x}\cos x$ is a solution of $y'' + 2y' + 2y = 0$.

10e $y = Ae^{-\frac{1}{2}x^2}$

$y' = A\left(-\frac{1}{2} \times 2xe^{-\frac{1}{2}x^2}\right)$

$= -Axe^{-\frac{1}{2}x^2}$

$y'' = -A\left(e^{-\frac{1}{2}x^2} \times 1 + x \times -\frac{1}{2} \times 2xe^{-\frac{1}{2}x^2}\right)$

$= -Ae^{-\frac{1}{2}x^2} + Ax^2e^{-\frac{1}{2}x^2}$

Substituting y and y'' into $y'' = y(x^2 - 1)$:

$\text{LHS} = -Ae^{-\frac{1}{2}x^2} + Ax^2e^{-\frac{1}{2}x^2}$

$\text{RHS} = Ae^{-\frac{1}{2}x^2}(x^2 - 1)$

$= Ax^2e^{-\frac{1}{2}x^2} - Ae^{-\frac{1}{2}x^2}$

$= -Ae^{-\frac{1}{2}x^2} + Ax^2e^{-\frac{1}{2}x^2}$

$= \text{LHS}$

Therefore $y = A\,e^{-x}\cos x$ is a solution of $y'' = y(x^2 - 1)$.

11a $y' = 1$

$y = x + A$

Substituting (2, 1):

$1 = 2 + \text{A}$

$A = -1$

Therefore $y = x - 1$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

11b $y' = 2x - 3$

$y = \frac{2x^2}{2} - 3x + A$

$y = x^2 - 3x + A$

Substituting $(0, 2)$:

$2 = (0)^2 - 3(0) + A$

$A = 2$

Therefore $y = x^2 - 3x + 2$

11c $y' = 3x^2 + 6x - 9$

$y = \frac{3x^3}{3} + \frac{6x^2}{2} - 9x + A$

$y = x^3 + 3x^2 - 9x + A$

Substituting $(1, 2)$:

$2 = (1)^3 + 3(1)^3 - 9(1) + A$

$2 = 1 + 3 - 9 + A$

$2 = -5 + A$

$A = 7$

Therefore $y = x^3 + 3x^2 - 9x + 7$

11d $y' = \sin x$

$y = -\cos x + A$

Substituting $(\pi, 3)$:

$3 = -\cos \pi + A$

$3 = 1 + A$

$A = 2$

Therefore $y = -\cos x + 2$

$y = 2 - \cos x$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

11e $y' = 6e^{2x}$

$y = 3e^{2x} + A$

Substituting (0, 0):

$0 = 3e^0 + A$

$0 = 3 + A$

$A = -3$

Therefore $y = 3e^{2x} - 3$

$y = 3(e^{2x} - 1)$

11f $y' = 3\sqrt{x} - 2$

$= 3x^{\frac{1}{2}} - 2$

$y = \frac{3}{\frac{3}{2}} \times x^{\frac{3}{2}} - 2x + A$

$= 2x^{\frac{3}{2}} - 2x + A$

Substituting (4, 7):

$7 = 2(4)^{\frac{3}{2}} - 2(4) + A$

$7 = 2(8) - 8 + A$

$7 = 8 + A$

$A = -1$

Therefore $y = 2x^{\frac{3}{2}} - 2x - 1$

$y = 2x\sqrt{x} - 2x - 1$

12a i

$$\text{RHS} = \frac{1}{x} + \frac{1}{1-x}$$

$$= \frac{1(1-x) + 1(x)}{x(1-x)}$$

$$= \frac{1 - x + x}{x(1-x)}$$

© Cambridge University Press 2019

$$= \frac{1}{x(1-x)}$$

$$= \text{LHS}$$

Hence:

$$\frac{1}{x(1-x)} = \frac{1}{x} + \frac{1}{1-x}$$

Alternatively, using a partial fractions approach:

$$\frac{1}{x(1-x)} = \frac{A}{x} + \frac{B}{1-x}$$

$$= \frac{A(1-x) + Bx}{x(1-x)}$$

$$= \frac{A - Ax + Bx}{x(1-x)}$$

$$= \frac{A - x(A-B)}{x(1-x)}$$

Equating coefficients in the numerators:

$1 + 0x = A - (A - B)x$

$A = 1 \qquad (1)$

$A - B = 0 \qquad (2)$

Substituting (1) into (2):

$1 - B = 0$

$B = 1$

Therefore:

$$\frac{1}{x(1-x)} = \frac{1}{x} + \frac{1}{1-x}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

12a ii

$$y = \int \frac{1}{x(1-x)}\,dx$$

$$= \int \left(\frac{1}{x} + \frac{1}{1-x}\right) dx$$

$$= \log_e|x| + (-\log_e|1-x|) + C$$

$$= \log_e|x| - \log_e|1-x| + C$$

$$= \log_e\left|\frac{x}{1-x}\right| + C \qquad (3)$$

12a iii Substituting $y\left(\frac{1}{2}\right) = 0$ into (3):

$$0 = \log_e\left|\frac{\frac{1}{2}}{1-\frac{1}{2}}\right| + C$$

$$0 = \log_e|1| + C$$

$$0 = 0 + C$$

$$C = 0$$

Therefore:

$$y = \log_e\left|\frac{x}{1-x}\right|$$

12b i

$$\text{RHS} = \frac{1}{2-x} + \frac{1}{2+x}$$

$$= \frac{1(2+x) + 1(2-x)}{(2-x)(2+x)}$$

$$= \frac{2+x+2-x}{(2-x)(2+x)}$$

$$= \frac{4}{(2-x)(2+x)}$$

$$= \text{LHS}$$

© Cambridge University Press 2019

Hence:

$$\frac{4}{(2-x)(2+x)} = \frac{1}{2-x} + \frac{1}{2+x}$$

Alternatively, using a partial fractions approach:

$$\begin{aligned}\frac{4}{(2-x)(2+x)} &= \frac{A}{2-x} + \frac{B}{2+x} \\ &= \frac{A(2+x) + B(2-x)}{(2+x)(2-x)} \\ &= \frac{2A + Ax + 2B - Bx}{(2+x)(2-x)} \\ &= \frac{x(A-B) + 2A + 2B}{(2+x)(2-x)}\end{aligned}$$

Equating coefficients in the numerators:

$0x + 4 = x(A-B) + 2A + 2B$

$A - B = 0 \qquad (1)$

$2A + 2B = 4 \qquad (2)$

From (1), $A = B$.

Substituting $A = B$ into (2):

$2B + 2B = 4$

$4B = 4$

$B = 1$

Hence $A = 1$.

Therefore:

$$\frac{4}{(2-x)(2+x)} = \frac{1}{2-x} + \frac{1}{2+x}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

12b ii

$$y = \int \frac{4}{(2-x)(2+x)}\,dx$$

$$= \int \left(\frac{1}{2-x} + \frac{1}{2+x}\right) dx$$

$$= -\log_e|2-x| + \log_e|2+x| + C$$

$$= \log_e|2+x| - \log_e|2-x| + C$$

$$= \log_e\left|\frac{2+x}{2-x}\right| + C \qquad (3)$$

12b iii Substituting $y(0) = 1$ into (3):

$$1 = \log_e\left|\frac{2+0}{2-0}\right| + C$$

$$1 = \log_e|1| + C$$

$$1 = 0 + C$$

$$1 = C$$

Hence:

$$y = \log_e\left|\frac{2+x}{2-x}\right| + 1$$

13a i

$$\frac{d(x^2)}{dx} + \frac{d(y^2)}{dx} = \frac{d(9)}{dx}$$

$$2x + \frac{d(y^2)}{dy} \times \frac{dy}{dx} = 0$$

$$2x + 2y\frac{dy}{dx} = 0$$

(or $2x + 2yy' = 0$)

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

13a ii

From $2x + 2y\frac{dy}{dx} = 0$

$$2y\frac{dy}{dx} = -2x$$

$$\frac{dy}{dx} = -\frac{2x}{2y}$$

$$\frac{dy}{dx} = -\frac{x}{y}$$

13a iii y' is undefined when $y = 0$. That is, at $(3, 0)$ and $(-3, 0)$ where the tangent to the circle is vertical.

13b i $y^2 = x + 4$

$$\frac{d(y^2)}{dx} = \frac{d(x)}{dx} + \frac{d(4)}{dx}$$

$$2y\frac{dy}{dx} = 1 + 0$$

$$\frac{dy}{dx} = \frac{1}{2y}$$

13b ii $xy = c^2$

$$y\frac{d(x)}{dx} + x\frac{d(y)}{dx} = \frac{d(c^2)}{dx}$$

$$y(1) + x\frac{dy}{dx} = 0$$

$$x\frac{dy}{dx} = -y$$

$$\frac{dy}{dx} = -\frac{y}{x}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

13b iii $9x^2 + 16y^2 = 144$

$$\frac{d(9x^2)}{dx} + \frac{d(16y^2)}{dx} = \frac{d(144)}{dx}$$

$$18x + 32y\frac{dy}{dx} = 0$$

$$32y\frac{dy}{dx} = -18x$$

$$\frac{dy}{dx} = -\frac{18x}{32y}$$

$$\frac{dy}{dx} = -\frac{9x}{16y}$$

13b iv $x^2 - 4y^2 = 4$

$$\frac{d(x^2)}{dx} - \frac{d(4y^2)}{dx} = \frac{d(4)}{dx}$$

$$2x - 8y\frac{dy}{dx} = 0$$

$$8y\frac{dy}{dx} = 2x$$

$$\frac{dy}{dx} = \frac{2x}{8y}$$

$$\frac{dy}{dx} = \frac{x}{4y}$$

13b v $xy - y^2 = 1$

$$\frac{d(xy)}{dx} - \frac{d(y^2)}{dx} = \frac{d(1)}{dx}$$

$$y\frac{d(x)}{dx} + x\frac{d(y)}{dx} - 2y\frac{dy}{dx} = 0$$

$$y + x\frac{dy}{dx} - 2y\frac{dy}{dx} = 0$$

$$\frac{dy}{dx}(x - 2y) = -y$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

$$\frac{dy}{dx} = -\frac{y}{x-2y}$$

$$\frac{dy}{dx} = \frac{y}{2y-x}$$

13b vi $x^3 + y^3 = 3xy$

$$\frac{d(x^3)}{dx} + \frac{d(y^3)}{dx} = \frac{d(3xy)}{dx}$$

$$3x^2 + 3y^2\frac{dy}{dx} = 3y\frac{d(x)}{dx} + 3x\frac{d(y)}{dx}$$

$$3x^2 + 3y^2\frac{dy}{dx} = 3y + 3x\frac{dy}{dx}$$

$$x^2 + y^2\frac{dy}{dx} = y + x\frac{dy}{dx}$$

$$y^2\frac{dy}{dx} - x\frac{dy}{dx} = y - x^2$$

$$\frac{dy}{dx}(y^2 - x) = y - x^2$$

$$\frac{dy}{dx} = \frac{y-x^2}{y^2-x}$$

14a $y = \sin x$

$y' = \cos x$

$y'' = -\sin x$

Substituting y and y'' into $y'' + y = 0$:

LHS $= -\sin x + \sin x$

$= 0$

$=$ RHS

Therefore $y = \sin x$ is a solution of $y'' + y' = 0$.

© Cambridge University Press 2019

14b $y'' = -\sin x$

$y'' = -y$

When $y = 12$,

$y'' = -12$

14c $f''(x)$ is the negative of $f(x)$ so it is a reflection in the x-axis.

15 $y'' = \sec^2 x$

$y' = \tan x + A$

$y' = \dfrac{\sin x}{\cos x} + A$

Since $y'(0) = 1$,

$1 = \dfrac{\sin 0}{\cos 0} + A$

$1 = 0 + A$

$A = 1$

Therefore $y' = \dfrac{\sin x}{\cos x} + 1$

Let $u = \cos x$

$\dfrac{du}{dx} = -\sin x$

$dx = -\dfrac{du}{\sin x}$

So:

$y = \displaystyle\int \left(\frac{\sin x}{\cos x} + 1\right) dx$

$= \displaystyle\int \frac{\sin x}{\cos x}\, dx + \int 1\, dx$

$= \displaystyle\int \frac{\sin x}{u}\left(-\frac{du}{\sin x}\right) + x + B$

© Cambridge University Press 2019

$= -\int \frac{1}{u}(du) + x + B$

$= -\log_e|u| + x + B$

$= -\log_e|\cos x| + x + B$

Since $y(0) = 1$,

$1 = -\log_e|\cos 0| + 0 + B$

$1 = -\log_e 1 + 0 + B$

$1 = B$

Therefore $y = -\log_e|\cos x| + x + 1$

$y = 1 + x - \log_e(\cos x)$

16 $y = \tan x$

$y' = \sec^2 x$

$= (\cos x)^{-2}$

$y'' = -2(\cos x)^{-3}(-\sin x)$

$= \frac{2\sin x}{\cos^3 x}$

$= \frac{2\sin x}{\cos x} \times \frac{1}{\cos^2 x}$

$= 2\tan x \sec^2 x$

First check

Substituting $y\left(\frac{\pi}{4}\right) = 1$ into $y = \tan x$:

RHS $= \tan\left(\frac{\pi}{4}\right)$

$= 1$

$=$ LHS

Second check

Substituting $y'\left(\frac{\pi}{4}\right) = 2$ into $y' = \sec^2 x$:

RHS $= \sec^2\left(\frac{\pi}{4}\right)$

© Cambridge University Press 2019

$$= \frac{1}{\cos^2\left(\frac{\pi}{4}\right)}$$

$$= \frac{1}{\left(\frac{1}{\sqrt{2}}\right)^2}$$

$$= 2$$

$$= \text{LHS}$$

Substituting y, y' and y'' into $y'' = 2yy'$:

$$\text{RHS} = 2 \times \tan x \times \sec^2 x$$

$$= 2 \tan x \sec^2 x$$

$$= \text{LHS}$$

Therefore $y = \tan x$ is a solution.

17 $y = Ae^{\lambda x}$

$y' = A\lambda e^{\lambda x}$

$y'' = A\lambda^2 e^{\lambda x}$

17a Substituting y, y' and y'' into $y'' - 4y' + 3y = 0$:

$A\lambda^2 e^{\lambda x} - 4A\lambda e^{\lambda x} + 3\,Ae^{\lambda x} = 0$

$Ae^{\lambda x}(\lambda^2 - 4\lambda + 3\,) = 0$

$Ae^{\lambda x}(\lambda - 1\,)(\lambda - 3) = 0$

$(\lambda - 1\,)(\lambda - 3) = 0 \quad (\text{as } Ae^{\lambda x} \neq 0)$

$\lambda = 1 \text{ or } \lambda = 3$

17b Substituting y, y' and y'' into $y'' + 2y' + y = 0$:

$A\lambda^2 e^{\lambda x} + 2A\lambda e^{\lambda x} + Ae^{\lambda x} = 0$

$Ae^{\lambda x}(\lambda^2 + 2\lambda + 1\,) = 0$

$Ae^{\lambda x}(\lambda + 1)^2 = 0$

$(\lambda + 1)^2 = 0 \quad (\text{as } Ae^{\lambda x} \neq 0)$

© Cambridge University Press 2019

$\lambda = -1$

17c Substituting y, y' and y'' into $y'' - 3y' + 4y = 0$:

$A\lambda^2 e^{\lambda x} - 3A\lambda e^{\lambda x} + 4Ae^{\lambda x} = 0$

$Ae^{\lambda x}(\lambda^2 - 3\lambda + 4\,) = 0$

$\lambda^2 - 3\lambda + 4 \ = 0 \quad$ (as $Ae^{\lambda x} \neq 0$)

Using the discriminant for a quadratic to check for any real solutions:

$\Delta = b^2 - 4ac$

$= (-3)^2 - 4(1)(4)$

$= \ 9 - 16$

$= \ -7$

Since $\Delta < 0$, no real solutions exist.

18a $y = \sec x$

$= (\cos x)^{-1}$

$y' = -(\cos x)^{-2} \times -\sin x$

$= \dfrac{\sin x}{\cos x} \times \dfrac{1}{\cos x}$

$= \tan x \sec x$

$y' - y \tan x$

$= \tan x \sec x - \sec x \tan x$

$= 0$

Therefore required IVP is $y' - y \tan x = 0, y(0) = 1$

18b $y^2 = \sec^2 x$

$(y')^2 = (\tan x \sec x)^2$

$= \tan^2 x \sec^2 x$

$= \tan^2 x \times y^2$

Now $\tan^2 x + 1 = \sec^2 x$

© Cambridge University Press 2019

$\tan^2 x = \sec^2 x - 1$

$\tan^2 x = y^2 - 1$

So $(y')^2 = \tan^2 x \times y^2$

$= (y^2 - 1) \times y^2$

Therefore required IVP is $(y')^2 = y^2(y^2 - 1), y(0) = 1$

19a $y(0) = 1$

19b $y' = -2xy$

Substituting $y(0) = 1$:

$y' = -2(0)(1)$

$= 0$

19c i $y' = -2xy$

$y'' = -2\left(y \times 1 + x \times 1\frac{dy}{dx}\right)$

$= -2y - 2xy'$

$= -2y - 2x(-2xy)$

$= -2y + 4x^2y$

$= y(4x^2 - 2)$

or $y'' = (4x^2 - 2)y$

19c ii $y'' = (4x^2 - 2)y$

At $(0, 1)$:

$y'' = (4 \times 0^2 - 2) \times 1$

$= -2$

Since $y'' < 0, y = f(x)$ is concave down.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

19d $y'' = (4x^2 - 2)y$

$= 4x^2y - 2y$

$y''' = 4y \times \frac{d(x^2)}{dx} + 4x^2 \times \frac{d(y)}{dx} - 2 \times \frac{d(y)}{dx}$

$= 4y \times 2x + 4x^2y' - 2y'$

$= 8xy + 4x^2y' - 2y'$

$= 8xy + 4x^2(-2xy) - 2(-2xy)$

$= 8xy - 8x^3y + 4xy$

At (0, 1):

$y''' = 8 \times 0 \times 1 - 8 \times 0 \times 1 + 4 \times 0 \times 1$

$= 0$

20a $y = cx - c^2$

$y' = c$

Substituting y and y' into $(y')^2 - xy' + y = 0$:

LHS $= (c)^2 - x(c) + cx - c^2$


$= c^2 - cx + cx - c^2$

$= 0$

$=$ RHS

Therefore $y = cx - c^2$ is a general solution of $(y')^2 - xy' + y = 0$.

20b



© Cambridge University Press 2019

They seem to form the outline of a curve (shown dotted)

20c For $c = p$,

$y = px - p^2$

For $c = p + h$,

$y = (p + h)x - (p + h)^2$

At point of intersection:

$(p + h)x - (p + h)^2 = px - p^2$

$px + hx - p^2 - 2ph - h^2 = px - p^2$

$hx - 2ph - h^2 = 0$

$hx = 2ph + h^2$

$hx = h(2p + h)$

$x = 2p + h \qquad (h \neq 0)$

$y = p(2p + h) - p^2$

$= 2p^2 + ph - p^2$

$y = p^2 + ph$

Therefore point of intersection is $(2p + h\,, p^2 + ph)$.

20d $\lim\limits_{h \to 0}(x)$

$= \lim\limits_{h \to 0}(2p + h)$

$= 2p$

$\lim\limits_{h \to 0}(y)$

$= \lim\limits_{h \to 0}(p^2 + ph)$

$= p^2$

Therefore, as $h \to 0$, point of intersection is $(2p\,, p^2)$.

© Cambridge University Press 2019

20e $x = 2p$

so $p = \frac{x}{2}$

Substituting $p = \frac{x}{2}$ into $y = p^2$:

$$y = \left(\frac{x}{2}\right)^2$$

$$y = \frac{1}{4}x^2$$

$$y' = \frac{1}{2}x$$

Substituting y and y' into $(y')^2 - xy' + y = 0$:

$$\text{LHS} = \left(\frac{1}{2}x\right)^2 - x\left(\frac{1}{2}x\right) + \frac{1}{4}x^2$$

$$\text{LHS} = \frac{1}{4}x^2 - \frac{1}{2}x^2 + \frac{1}{4}x^2$$

$$= 0$$

$$= \text{RHS}$$

Therefore $y = \frac{1}{4}x^2$ is a solution of $(y')^2 - xy' + y = 0$.

This is called a singular solution. This means the solution is not obtained from the general solution but obtained by the usual method of solving the differential equation. In solving this differential equation, a general solution consisting of a family of curves is obtained.

Each line in part b is tangent to the parabola. Thus every point in the parabola is also a point in a general solution. Hence the parabola itself is also a solution.

21 $y^{(n)} = 1$

$$y^{(n-1)} = \int 1\,dx$$

$$= x + A_1$$

$$y^{(n-2)} = \int (x + A_1)\,dx$$

$$= \frac{x^2}{2} + A_1x + A_2$$

© Cambridge University Press 2019

$$y^{(n-3)} = \int \left(\frac{x^2}{2} + \frac{A_1 x}{1} + A_2\right) dx$$

$$= \frac{x^3}{3 \times 2} + \frac{A_1 x^2}{2 \times 1} + \frac{A_2 x}{1} + A_3$$

$$y^{(n-4)} = \int \left(\frac{x^3}{3 \times 2} + \frac{A_1 x^2}{2 \times 1} + \frac{A_2 x}{1} + A_3\right) dx$$

$$= \frac{x^4}{4 \times 3 \times 2} + \frac{A_1 x^3}{3 \times 2} + \frac{A_2 x^2}{2 \times 1} + \frac{A_3 x}{1} + A_4$$

Following this pattern gives:

$$y^{(2)} = \frac{x^{n-2}}{(n-2)!} + \frac{A_1 x^{n-3}}{(n-3)!} + \frac{A_2 x^{n-4}}{(n-4)!} + \cdots + \frac{A_{n-3} x}{1!} + A_{n-2}$$

$$y^{(1)} = \frac{x^{n-1}}{(n-1)!} + \frac{A_1 x^{n-2}}{(n-2)!} + \frac{A_2 x^{n-3}}{(n-3)!} + \cdots + \frac{A_{n-2} x}{1!} + A_{n-1}$$

$$y = y^{(0)} = \frac{x^n}{n!} + \frac{A_1 x^{n-1}}{(n-1)!} + \frac{A_2 x^{n-2}}{(n-2)!} + \cdots + \frac{A_{n-1} x}{1!} + A_n$$

Or, absorbing the factorial terms into the constants:

$$y = \frac{x^n}{n!} + C_1 x^{n-1} + C_2 x^{n-2} + \cdots + C_n$$

There are n arbitrary constants.

© Cambridge University Press 2019

Solutions to Exercise 13B

1a $y' = 2(1) - 3 = -1$

1b $y' = 2\cos(0) - 1 = 1$

1c $y' = 4 - (1)^2 = 3$

1d $y' = \frac{1}{1+(1)} = \frac{1}{2}$

1e $y' = \frac{(1)}{(-2)} + 1 = \frac{1}{2}$

1f $y' = (1)(-2) - (1) = -3$

2a

y \ x	-1	0	1	2	3	4	5
3	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$
2	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$
1	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$
0	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$
-1	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$
-2	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$
-3	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$

2d Every entry in that column is the same.

2e Every vertical line $x = k$ is an isocline.


2f concave up with a minimum turning point at $x = 2$

2g a parabola

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

2h



3a $y' = x - y$


3b

y \ x	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2
1	-2	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1
$\frac{1}{2}$	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$
0	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2
$-\frac{1}{2}$	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2	$\frac{5}{2}$
-1	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2	$\frac{5}{2}$	3
$-\frac{3}{2}$	$\frac{1}{2}$	1	$\frac{3}{2}$	2	$\frac{5}{2}$	3	$\frac{7}{2}$
-2	1	$\frac{3}{2}$	2	$\frac{5}{2}$	3	$\frac{7}{2}$	4

3e The lines $y = x + k$ are isoclines.

3f concave up

3g, h



© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

3i $y = x - 1$

3j $y' = x - y$

$\text{LHS} = \frac{d}{dx}(x - 1)$

$= 1$


$\text{RHS} = x - y$

$= x - (x - 1)$

$= 1$

Therefore it is a solution.

4a



4a i x axis

4a ii any horizontal line


4a iii The gradients decrease to zero then increase.

4a iv The gradients are the same. It is an isocline.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4b




4b i There are no points or isoclines where $y' = 0$.

4b ii any vertical line

4b iii The gradients are the same. It is an isocline.

4b iv The gradients decrease to vertical then increase.

4c



4c i $x = -2, x = 2$

4c ii any vertical line


4c iii The gradients are the same. It is an isocline.

4c iv The gradients increase from -1 to 1 then back again.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4d




4d i $y = x - 1$

4d ii $y = x + C$

4d iii The gradients increase.

4d iv The gradients decrease.

4e



4e i $x = 2$

4e ii $y = 0$ (excepting the point (0, 0) where the gradient is undefined)


4e iii The gradients increase.

4e iv The gradients decrease, but the gradient is undefined at the y-axis.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4f




4f i $y = 1$ or $x = 0$

4f ii Undefined at $y = -1$.


4f iii The gradients decrease, but the gradient is undefined at $y = -1$.

4f iv The gradients decrease.

5a




5b




© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations


5c




5d



5e



5f



6a All vertical lines are isoclines so $y' = f(x)$.

6b All vertical lines are isoclines so $y' = f(x)$.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations


6c All horizontal lines are isoclines so $y' = g(y)$.

6d All horizontal lines are isoclines so $y' = g(y)$.

6e Not all horizontals and not all verticals are isoclines, so y' is a combination.

6f Not all horizontals and not all verticals are isoclines, so y' is a combination.

7a Solution curves through the points $(0, -2)$ and $(0, 2)$ are shown.



7b i The gradients decrease.

7b ii The solution curves converge as they cross $x = 1$ from left to right.

7c $y' = -\frac{1}{2}$ everywhere on that line

7d

$$y = \frac{1}{2} - \frac{1}{2}x$$

$$y' = -\frac{1}{2}x - y$$

© Cambridge University Press 2019

$$\text{LHS} = \frac{d}{dx}\left(\frac{1}{2} - \frac{1}{2}x\right)$$

$$= -\frac{1}{2}$$

$$\text{RHS} = -\frac{1}{2}x - y$$


$$= -\frac{1}{2}x - \left(\frac{1}{2} - \frac{1}{2}x\right)$$

$$= -\frac{1}{2}$$

Therefore it is a solution.

7e The isocline is an asymptote for each one.

8a



8b $y = -3, 3$

8c Yes, these constant solutions are isoclines.

8d i converge

8d ii diverge

8d iii yes

8d iv They are the asymptotes for the solution curves.

8e See part a.

9 $y' = -2 - y$

Since gradient depends only on y, all horizontal lines should be isoclines (eliminating options A and B).

Both the remaining solutions show $y' = 0$ at $y = -2$ which fits with the DE given.

Checking D we can confirm that the slope field seems consistent with the equation given.

10 All vertical lines are isoclines but horizontal lines are not, so y' must be a function of x, eliminating options B and D. At $x = 0$, the slope is positive, which eliminates A. Checking C we can confirm that the slope field seems consistent with the equation given.

11 $y' = 1 - \frac{x}{y}$

For $x = 0, y \neq 0$, slope should equal 1. This eliminates options C and D.

For $x = y \neq 0$, slope should equal 0. This eliminates option A.

Checking B we can confirm that the slope field seems consistent with the equation given.


12a Along vertical and horizontal lines, the slope increases with increasing x, and decreases with increasing y. Of the four options given, only B is consistent with this pattern.

12b Along all horizontal lines, the slope is 0 at $x = 0$, and asymptotes to zero again as x becomes large in either direction. However, under options A and C, the slope would tend to $\pm\infty$ as x becomes large, so we can rule these out. At (1, 1), option B predicts a slope of 1 (inconsistent with the graph shown) and D predicts slope of -1 (consistent with the graph shown) so the only valid answer is D.


© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

13a i,ii



13b i,ii



13c i $xy = 4$

Need to show $y' = -\frac{y}{x}$

$$\frac{d}{dx}(xy) = \frac{d}{dx}(4)$$

$$(1)(y) + (x)\frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = -\frac{y}{x}$$

Hyperbola passes through $(2, 2)$ and $(-2, -2)$.

Therefore it is a solution of the IVP in part a.

13c ii $x^2 - y^2 = 4$

Need to show $y' = -\frac{x}{y}$

$$\frac{d}{dx}(x^2 - y^2) = \frac{d}{dx}(4)$$

$$2x - (2y)\frac{dy}{dx} = 0$$

© Cambridge University Press 2019


$$\frac{dy}{dx} = \frac{-2x}{-2y}$$

$$\frac{dy}{dx} = \frac{x}{y}$$

Hyperbola passes through $(2, 0)$ and $(-2, 0)$.


Therefore it is a solution of the IVP in part b.

13c iii



13d At any point where these curves intersect, gradient of the first curve is $-\frac{y}{x}$ and gradient of the second curve is $\frac{x}{y}$. The product of these two is always -1, hence the hyperbolas are perpendicular where they intersect.


14a i, ii



© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

14b i, ii



14c i $y = \frac{1}{2}x^2$

Need to show $y' = \frac{2y}{x}$

LHS $= y'$

$= \frac{d}{dx}(y)$

$= \frac{d}{dx}\left(\frac{1}{2}x^2\right)$

$= x$

RHS $= \frac{2y}{x}$

$= \frac{2\left(\frac{1}{2}x^2\right)}{x}$

$= x$

Also the parabola passes through (2, 2).

Therefore it is a solution of the IVP in part a.

14c ii $x^2 + 2y^2 = 4$

Need to show $y' = -\frac{x}{2y}$

$\frac{d}{dx}(x^2 + 2y^2) = \frac{d}{dx}(4)$

$(2x) + (4y)\frac{dy}{dx} = 0$

© Cambridge University Press 2019


$$\frac{dy}{dx} = -\frac{2x}{4y}$$

$$\frac{dy}{dx} = -\frac{x}{2y}$$

Also the ellipse passes through $(2, 0)$ and $(-2, 0)$.

Therefore it is a solution of the IVP in part b.

14c iii



14d The gradient of the parabola is $\frac{2y}{x}$ and the gradient of the ellipse is $-\frac{x}{27}$.
At any given point the product of these gradients is -1 so the parabola and the ellipse are perpendicular where they intersect.

15a $x^2 + y^2 = 16$

$$\frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) = \frac{d}{dx}(16)$$

$$2x + 2y\frac{dy}{dx} = 0$$

$$2y\frac{dy}{dx} = -2x$$

$$\frac{dy}{dx} = -\frac{2x}{2y}$$

$\frac{dy}{dx} = -\frac{x}{y}$ as required

15b $(x-3)^2 + (y-1)^2 = 4$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

15c $(x-3)^2+(y-1)^2=4$

$$\frac{d}{dx}(x-3)^2+\frac{d}{dx}(y-1)^2=\frac{d}{dx}(16)$$

$$\frac{d}{dx}(x^2-6x+9)+\frac{d}{dx}(y^2-2y+1)=\frac{d}{dx}(16)$$

$$(2x-6)+(2y-2)\frac{dy}{dx}=0$$

$$(2y-2)\frac{dy}{dx}=-(2x-6)$$

$$\frac{dy}{dx}=-\frac{(2x-6)}{(2y-2)}$$


$$\frac{dy}{dx}=-\frac{2(x-3)}{2(y-1)}$$

$$\frac{dy}{dx}=-\frac{x-3}{y-1}$$

15d i

$$\frac{dy}{dx}=\frac{x-1}{y+2}$$


15d ii



© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

16a, b




16c $\Phi(1) \doteq 0.8$; a better approximation is 0.8413.

17a $x = 0$ and $y = 0$

17b The rectangular hyperbola $xy = C$

17c-i



17j i The advantage of this technique is the exact gradient of the integral curve is known as an isocline is crossed.

17j ii The disadvantage of this technique is it takes a lot of time to sketch the isoclines.


18a i The line elements of the slope field for points on the y-axis other than the origin are horizontal.

18a ii The line elements of the slope field for points on the x-axis are vertical.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations


18a iii



18a iv circle

18b i $y = -\frac{1}{C}x$

18b ii



The straight lines at 30° and 60° to the x-axis.

18b iii Each line element is perpendicular to its isocline because the product of the gradients

$$C \times \frac{-1}{C} = -1$$

18b iv See graph in part b ii.

18b v Yes, the shape of part c is clearer now. Notice that the innermost line elements almost join up to give the outline of a circle.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

19a Any two points on the same vertical line have the same x-value.

Therefore, if $y' = f(x)$, then any two points on the same vertical line give the same slope, making this line an isocline.

Let D be the domain of $f(x)$. For each a in D, the point (a, y) gives $y' = f(a)$ for all values of y. Hence $x = a$ is an isocline for each a in D.

19b Any two points on the same horizontal line have the same y-value. Therefore, if $y' = g(y)$, then any two points on the same horizontal line give the same slope, making this line an isocline.

19c If $y = cx + b$ is a solution to this DE, then $y' = \frac{d(cx+b)}{dx} = c$ everywhere along the line. Hence all points on the line give the same gradient, so it is an isocline.

20a $x^3 + y^3 = 3xy$

Differentiating:

$$\frac{d(x^3)}{dx} + \frac{d(y^3)}{dx} = \frac{d(3xy)}{dx}$$

$$3x^2 + 3y^2\frac{dy}{dx} = 3y + 3x\frac{dy}{dx}$$

Grouping terms:

$$\frac{dy}{dx}(3y^2 - 3x) = 3y - 3x^2$$


$$y' = \frac{3y - 3x^2}{3y^2 - 3x}$$

$$= \frac{y - x^2}{y^2 - x}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

20b



20c $y' = 0$ when $y - x^2 = 0$, that is, when $y = x^2$, excluding the points $(0, 0)$ and $(1, 1)$ where the gradient is undefined.

20d $\frac{1}{y'}$ approaches zero as $y^2 - x$ approaches zero, that is, the integral curve is vertical when $x = y^2$, again excluding the points $(0, 0)$ and $(1, 1)$.

20e See part b.

20f Although gradients here are undefined, we can see that the curve crosses itself here.

20g $y = -x - 1$ appears to be an isocline.

Substituting this into the DE from part a, we find:

$$y' = \frac{y - x^2}{y^2 - x}$$

$$= \frac{-x - 1 - x^2}{(-x - 1)^2 - x}$$

$$= \frac{-x^2 - x - 1}{x^2 + 2x + 1 - x}$$

© Cambridge University Press 2019

$$= \frac{-x^2 - x - 1}{x^2 + x + 1}$$

$$= -1$$

Hence the slope field equals -1 everywhere along this line, so it is an isocline. The gradient of the line is also equal to -1 so it is a solution of the DE.

20h The curve is symmetrical in the line $y = x$.

20h i The equation is symmetric in x and y, hence we expect the solution to be symmetric when we swap x and y by reflecting in $y = x$.

20h ii Swapping x and y in the RHS of the differential equation has the same effect as taking the reciprocal of the gradient, hence the gradient field is symmetric around $y = x$ and so we expect the curve to have the same symmetry.

© Cambridge University Press 2019

Solutions to Exercise 13C

Let the integration constants be A, B, C or D.

1a

$$\frac{dy}{dx} = \frac{x-1}{y+1}$$

$$(y+1)\frac{dy}{dx} = x - 1$$

$$(y+1)\,dy = (x-1)\,dx$$

$$\int (y+1)dy = \int (x-1)\,dx$$

1b

$$\int (y+1)dy = \int (x-1)dx$$

$$\frac{(y+1)^2}{2} = \frac{(x-1)^2}{2} + C$$

$$(y+1)^2 = (x-1)^2 + 2C$$

$$(y+1)^2 = (x-1)^2 + D, \text{ where } D = 2C$$

2a

$$\frac{dy}{dx} = xe^{-y}$$

$$\left(\frac{1}{e^{-y}}\right) dy = x\,dx$$

$$\int \left(\frac{1}{e^{-y}}\right) dy = \int x\,dx$$

$$\int (e^{y})\,dy = \int x\,dx$$

$$e^{y} = \frac{x^2}{2} + C$$

$$\ln(e^{y}) = \ln\left|\frac{x^2}{2} + C\right|$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

$$y = \ln\left|\frac{x^2}{2} + C\right|$$

2b

$$\frac{dy}{dx} = 4x^3(1 + y^2)$$

$$\left(\frac{1}{1 + y^2}\right) dy = 4x^3\, dx$$

$$\int \left(\frac{1}{1 + y^2}\right) dy = \int 4x^3\, dx$$

$$\tan^{-1} y = \frac{4x^4}{4} + C$$

$$y = \tan(x^4 + C)$$

3a Substitute $y = 0$, where $x \neq 0$. Then LHS and RHS are both zero.

3b

$$\frac{dy}{dx} = -\frac{y^2}{x}$$

$$\frac{1}{y^2}\, dy = -\frac{1}{x}\, dx$$

$$\int \left(\frac{1}{y^2}\right) dy = \int \left(-\frac{1}{x}\right) dx$$

$$\int (y^{-2})\, dy = -\int \frac{1}{x}\, dx$$

$$\frac{y^{-1}}{-1} = -\ln|x| + C$$

$$-\frac{1}{y} = -\ln|x| + C$$

$$-\frac{y}{1} = -\frac{1}{\ln|x| + C}$$

$$y = \frac{1}{\ln|x| + C}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4a

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$y\,dy = -x\,dx$$

$$\int y\,dy = -\int x\,dx$$

4b

$$\int y\,dy = -\int x\,dx$$

$$\frac{y^2}{2} = -\frac{x^2}{2} + C$$

$$y^2 = -x^2 + 2C$$

$y^2 = -x^2 + D$, where $D = 2C$

4c Substituting $\left(1, \sqrt{3}\right)$:

$$\left(\sqrt{3}\right)^2 = -(1)^2 + D$$

$$3 = -1 + D$$

$$4 = D$$

Therefore $y^2 = -x^2 + 4$ or $y^2 + x^2 = 4$

5a

$$\frac{dy}{dx} = \frac{x}{y}$$

$$y\,dy = x\,dx$$

$$\int y\,dy = \int x\,dx$$

$$\frac{y^2}{2} = \frac{x^2}{2} + C$$

$$y^2 = x^2 + 2C$$

$y^2 = x^2 + D$, where $D = 2C$

© Cambridge University Press 2019

Substituting $(0, 1)$:

$(1)^2 = (0)^2 + D$

$1 = D$

Therefore $y^2 = x^2 + 1$

5b

$\frac{dy}{dx} = (1 + x)\,(1 + y^2)$

$\left(\frac{1}{1 + y^2}\right)\,dy = (1 + x)\,dx$

$\int \left(\frac{1}{1 + y^2}\right) dy = \int (1 + x) dx$

$\tan^{-1} y = \frac{(1 + x)^2}{2} + C$

$y = \tan\left(\frac{1}{2}(1 + x)^2 + C\right)$

Substituting $(-1, 0)$:

$0 = \tan\left(\frac{1}{2}(1 + (-1))^2 + C\right)$

$0 = \tan(C)$

$0 = C$

Therefore, $y = \tan\left(\frac{1}{2}(1 + x)^2\right)$

5c

$\frac{dy}{dx} = -2y^2x$

$\frac{1}{y^2}\,dy = -2x\,dx$

$\int \frac{1}{y^2} dy = \int (-2x) dx$

$\int (y^{-2})\,dy = \int (-2x) dx$

© Cambridge University Press 2019

$$\frac{y^{-1}}{-1} = \frac{-2x^2}{2} + C$$

$$-y^{-1} = -x^2 + C$$

$$y^{-1} = x^2 - C$$

$$\frac{1}{y} = \frac{x^2 - C}{1}$$

$$\frac{y}{1} = \frac{1}{x^2 - C}$$

$$y = \frac{1}{x^2 - C}$$

Substituting $\left(1, \frac{1}{2}\right)$:

$$\frac{1}{2} = \frac{1}{(1)^2 - C}$$

$$2 = 1 - C$$

$$C = -1$$

Therefore $y = \frac{1}{x^2 + 1}$

5d

$$\frac{dy}{dx} = e^{-y}\sec^2 x$$

$$\left(\frac{1}{e^{-y}}\right) dy = \sec^2 x \, dx$$

$$\int \left(\frac{1}{e^{-y}}\right) dy = \int \sec^2 x \, dx$$

$$\int (e^y) \, dy = \int \sec^2 x \, dx$$

$$e^y = \tan x + C$$

$$y = \ln(C + \tan x)$$

Substituting $\left(\frac{\pi}{4}, \ln 2\right)$:

$$\ln 2 = \ln(C + \tan\frac{\pi}{4})$$

© Cambridge University Press 2019

$2 = C + \tan\frac{\pi}{4}$

$2 = C + 1$

$1 = C$

Therefore $y = \ln(1 + \tan x)$

(More strictly, the solution is only valid in the interval $-\frac{\pi}{2} < x < \frac{\pi}{2}$.)

6a

$\frac{dy}{dx} = \frac{2y + 4}{x}$

$y = -2$ is a constant solution

$\text{LHS} = \frac{d(-2)}{dx} = 0$

$\text{RHS} = \frac{2(-2) + 4}{x} = 0$

LHS = RHS

Therefore $y = -2$ is a solution of the DE.

6b

$\frac{dy}{dx} = \frac{2y + 4}{x}$

$\frac{1}{2y + 4}\, dy = \frac{1}{x}\, dx$

$\int \frac{1}{2y + 4} dy = \int \frac{1}{x}\, dx$

$\frac{1}{2}\int \frac{1}{y + 2}\, dy = \int \frac{1}{x}\, dx$

$\frac{1}{2}\ln|y + 2| = \ln|x| + B$

$\ln|y + 2| = 2\ln|x| + 2B$

$\ln|y + 2| = \ln(x^2) + \ln(e^{2B})$

$\ln|y + 2| = \ln(e^{2B}x^2)$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

$y + 2 = e^{2B}x^2$ or $-e^{2B}x^2$

$y + 2 = Cx^2$ where $C = e^{2B}$ or $-e^{2B}$ and $C \neq 0$

$y = Cx^2 - 2$

6c Allow $C = 0$.

7a $y' = -xy$

$y = 0$ is a constant solution

$\text{LHS} = \dfrac{d(0)}{dx} = 0$

$\text{RHS} = -x \times 0 = 0$

LHS = RHS

Therefore $y = 0$ is a solution of the DE.

7b

$\dfrac{dy}{dx} = -xy$

$\dfrac{1}{y}\, dy = -x\, dx$

$\displaystyle\int \frac{1}{y}\, dy = \int (-x)\, dx$

$\displaystyle\int \frac{1}{y}\, dy = -\int x\, dx$

$\ln|y| = -\dfrac{x^2}{2} + B$

$|y| = e^{\left(-\frac{1}{2}x^2 + B\right)}$

$y = e^{-\frac{1}{2}x^2} \times e^B$ or $-e^{-\frac{1}{2}x^2} \times e^B$

$y = Ce^{-\frac{1}{2}x^2}$ where $C = e^B$ or $-e^B$ and $C \neq 0$

7c Allow $C = 0$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

8a

$$\frac{dy}{dx} = \frac{2-y}{x} = \frac{y-2}{-x}$$

$$\frac{1}{y-2}\,dy = -\frac{1}{x}\,dx$$

$$\int \frac{1}{y-2}\,dy = -\int \frac{1}{x}\,dx$$

$$\ln|y-2| = -\ln|x| + B$$

$$\ln|y-2| = \ln|x^{-1}| + \ln(e^B)$$

$$\ln|y-2| = \ln(e^B|x^{-1}|)$$

$$|y-2| = e^B|x^{-1}|$$

$$y-2 = e^Bx^{-1} \text{ or } -e^Bx^{-1}$$

$$y-2 = Cx^{-1} \text{ where } C = e^B \text{ or } -e^B$$

$$y-2 = \frac{C}{x}$$

$$y = \frac{C}{x} + 2$$

8b

$$\frac{dy}{dx} = \frac{xy}{1+x^2}$$

$$\frac{1}{y}\,dy = \frac{x}{1+x^2}\,dx$$

$$\int \frac{1}{y}\,dy = \int \frac{x}{1+x^2}\,dx$$

$$\int \frac{1}{y}\,dy = \frac{1}{2}\int \frac{2x}{1+x^2}\,dx$$

$$2\ln|y| = \ln|1+x^2| + B$$

$$\ln(y^2) = \ln(1+x^2) + B$$

$$\ln(y^2) = \ln(1+x^2) + \ln(C) \quad \text{where } B = \ln(C)$$

$$\ln(y^2) = \ln(C(1+x^2))$$

$$y^2 = C(1+x^2)$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

8c

$$\frac{dy}{dx} = \frac{-2y}{x}$$

$$\frac{1}{y}\,dy = \left(-\frac{2}{x}\right)dx$$

$$\int \frac{1}{y}\,dy = \int \left(-\frac{2}{x}\right)dx$$

$$\ln|y| = -2\ln|x| + B$$

$$\ln|y| = \ln\left(x^{-2}\right) + B$$

$$\ln|y| = \ln\left(x^{-2}\right) + \ln(e^B)$$

$$\ln|y| = \ln(e^B x^{-2})$$

$$|y| = e^B x^{-2}$$

$$y = \frac{e^B}{x^2} \text{ or } -\frac{e^B}{x^2}$$

$$y = \frac{C}{x^2} \text{ where } C = e^B \text{ or } -e^B$$

8d

$$\frac{dy}{dx} = y\sin x$$

$$\frac{1}{y}\,dy = \sin x\;dx$$

$$\int \frac{1}{y}\,dy = \int \sin x\;dx$$

$$\ln|y| = -\cos x + B$$

$$\ln|y| = \ln(e^{-\cos x}) + \ln(e^B)$$

$$\ln|y| = \ln(e^B e^{-\cos x})$$

$$|y| = e^B e^{-\cos x}$$

$$y = e^B e^{-\cos x} \text{ or } -e^B e^{-\cos x}$$

$$y = Ce^{-\cos x} \text{ where } C = e^B \text{ or } -e^B$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

8e

$$\frac{dy}{dx} = \frac{3y}{x^2}$$

$$\frac{1}{3y}\,dy = \frac{1}{x^2}\,dx$$

$$\int \frac{1}{3y}\,dy = \int \frac{1}{x^2}\,dx$$

$$\frac{1}{3}\int \frac{1}{y}\,dy = \int \frac{1}{x^2}\,dx$$

$$\frac{1}{3}\ln|y| = \frac{x^{-1}}{-1} + B$$

$$\frac{1}{3}\ln|y| = -\frac{1}{x} + B$$

$$\ln|y| = -\frac{3}{x} + 3B$$

$$\ln|y| = \ln\left(e^{-\frac{3}{x}}\right) + \ln(e^{3B})$$

$$\ln|y| = \ln\left(e^{3B}e^{-\frac{3}{x}}\right)$$

$$|y| = e^{3B}e^{-\frac{3}{x}}$$

$$y = e^{3B}e^{-\frac{3}{x}} \text{ or } -e^{3B}e^{-\frac{3}{x}}$$

$$y = Ce^{-\frac{3}{x}} \text{ where } C = e^{3B} \text{ or } -e^{3B}$$

8f

$$\frac{dy}{dx} = \frac{y(1-x)}{x}$$

$$\frac{1}{y}\,dy = \frac{1-x}{x}\,dx$$

$$\int \frac{1}{y}\,dy = \int \frac{1-x}{x}\,dx$$

$$\int \frac{1}{y}\,dy = \int \left(\frac{1}{x} - \frac{x}{x}\right)\,dx$$

© Cambridge University Press 2019

$$\int \frac{1}{y}\,dy = \int \left(\frac{1}{x} - 1\right)\,dx$$

$$\ln|y| = \ln|x| - x + B$$

$$\frac{1}{2}\ln|y^2| = \frac{1}{2}\ln|x^2| - x + B$$

$$\ln(y^2) = \ln(x^2) - 2x + 2B$$

$$\ln(y^2) = \ln(x^2) + \ln(e^{-2x}) + \ln(e^{2B})$$

$$\ln(y^2) = \ln(e^{2B}x^2e^{-2x})$$

$$y^2 = e^{2B}x^2e^{-2x}$$

$$y = e^B xe^{-x} \text{ or } -e^B xe^{-x}$$

$$y = Cxe^{-x} \text{ where } C = e^B \text{ or } -e^B$$

9a $\cos y = 0$

$$y = -\frac{3\pi}{2}, -\frac{\pi}{2}, -\frac{3\pi}{2}, \frac{\pi}{2} \text{ and } \frac{3\pi}{2}$$

9b

$$\frac{dy}{dx} = 3x^2\cos^2 y$$

$$\frac{1}{\cos^2 y}\,dy = 3x^2\,dx$$

$$\int \frac{1}{\cos^2 y}\,dy = \int 3x^2\,dx$$

$$\int \sec^2 y\,dy = \int 3x^2\,dx$$

$$\tan y = \frac{3x^3}{3} + C$$

$$\tan y = x^3 + C$$

$$y = \tan^{-1}(x^3 + C)$$

10a Constant solution is $y = 0$ which does not match the initial value.

© Cambridge University Press 2019

10b

$$\frac{dy}{dx} = \frac{2y}{x-1}$$

$$\frac{1}{y}\,dy = \frac{2}{x-1}\,dx$$

$$\int \frac{1}{y}\,dy = \int \frac{2}{x-1}\,dx$$

$$\int \frac{1}{y}\,dy = 2\int \frac{1}{x-1}\,dx$$

$$\ln|y| = 2\ln|x-1| + B$$

$$\ln|y| = \ln((x-1)^2) + \ln(e^B)$$

$$\ln|y| = \ln(e^B(x-1)^2)$$

$$|y| = e^B(x-1)^2 \text{ or } -e^B(x-1)^2$$

$$y = C(x-1)^2 \text{ where } C = e^B \text{ or } -e^B$$

10c Substituting $y(2) = 1$:

$$1 = C(2-1)^2$$

$$1 = C$$

$$y = (x-1)^2$$

11a Constant solution is $y = 1$ which does not satisfy the IVP.

11b

$$\frac{dy}{dx} = (y-1)\tan x$$

$$\frac{1}{y-1}\,dy = \tan x\,dx$$

$$\int \frac{1}{y-1}\,dy = \int \tan x\,dx$$

$$\int \frac{1}{y-1}\,dy = -\int \left(\frac{-\sin x}{\cos x}\right) dx$$

$$\ln|y-1| = -\ln|\cos x| + B$$

© Cambridge University Press 2019

$\ln|y-1| = \ln|\sec x| + \ln(e^B)$

$\ln|y-1| = \ln(e^B|\sec x|)$

$|y-1| = e^B|\sec x|$

$y-1 = e^B \sec x$ or $-e^B \sec x$

$y-1 = C\sec x$ where $C = e^B$ or $-e^B$

$y = 1 + C\sec x$

11c Substituting $y\left(\frac{\pi}{4}\right) = 3$:

$3 = 1 + C\sec\left(\frac{\pi}{4}\right)$

$3 = 1 + C \times \frac{\sqrt{2}}{1}$

$3 - 1 = \sqrt{2}C$

$C = \frac{2}{\sqrt{2}} = \sqrt{2}$

Therefore $y = 1 + \sqrt{2}\sec x$

12a

$\frac{dy}{dx} = \frac{y}{x}$

$\frac{1}{y}\,dy = \frac{1}{x}\,dx$

$\int \frac{1}{y}\,dy = \int \frac{1}{x}\,dx$

$\ln|y| = \ln|x| + B$

$\ln|y| = \ln|x| + \ln(e^B)$

$\ln|y| = \ln(e^B|x|)$

$|y| = e^B|x|$

$y = e^B x$ or $-e^B x$

$y = Cx$ where $C = e^B$ or $-e^B$

© Cambridge University Press 2019

Substituting $y(2) = 1$:

$1 = C \times 2$

$\frac{1}{2} = C$

Therefore $y = \frac{1}{2}x$

12b

$\frac{dy}{dx} = \frac{y}{2x}$

$\frac{1}{y}\,dy = \frac{1}{2x}\,dx$

$\int \frac{1}{y}\,dy = \int \frac{1}{2x}\,dx$

$\int \frac{1}{y}\,dy = \frac{1}{2}\int \frac{1}{x}\,dx$

$\ln|y| = \frac{1}{2}\ln|x| + B$

$\ln|y| = \ln|x|^{\frac{1}{2}} + \ln(e^B)$

$\ln|y| = \ln\left(e^B|x|^{\frac{1}{2}}\right)$

$|y| = e^B|x|^{\frac{1}{2}}$

$y = e^B x^{\frac{1}{2}}$ or $-e^B x^{\frac{1}{2}}$

$y = C\sqrt{x}$ where $C = e^B$ or $-e^B$

Substituting $y(1) = 2$:

$2 = C \times 1$

$2 = C$

Therefore $y = 2\sqrt{x}$

© Cambridge University Press 2019

12c

$$\frac{dy}{dx} = \frac{-2xy}{1+x^2}$$

$$\frac{1}{y}\,dy = \frac{-2x}{1+x^2}\,dx$$

$$\int \frac{1}{y}\,dy = \int \frac{-2x}{1+x^2}\,dx$$

$$\int \frac{1}{y}\,dy = -\int \frac{2x}{1+x^2}\,dx$$

$$\ln|y| = -\ln|1+x^2| + B$$

$$\ln|y| = \ln((1+x^2)^{-1}) + \ln(e^B)$$

$$\ln|y| = \ln(e^B(1+x^2)^{-1})$$

$$|y| = e^B(1+x^2)^{-1}$$

$$y = \frac{e^B}{1+x^2} \text{ or } -\frac{e^B}{1+x^2}$$

$$y = \frac{C}{1+x^2} \text{ where } C = e^B \text{ or } -e^B$$

Substituting $y(1) = 2$:

$$2 = \frac{C}{1+1}$$

$$4 = C$$

Therefore $y = \frac{4}{1+x^2}$

12d

$$\frac{dy}{dx} = -\frac{y}{x}$$

$$\frac{1}{y}\,dy = -\frac{1}{x}\,dx$$

$$\int \frac{1}{y}\,dy = -\int \frac{1}{x}\,dx$$

$$\ln|y| = -\ln|x| + B$$

$$\ln|y| = \ln|x|^{-1} + \ln(e^B)$$

© Cambridge University Press 2019

$\ln|y| = \ln(e^B|x|^{-1})$

$|y| = e^B|x|^{-1}$

$y = \frac{e^B}{x}$ or $-\frac{e^B}{x}$

$y = \frac{C}{x}$ where $C = e^B$ or $-e^B$

Substituting $y(2) = 1$:

$1 = \frac{C}{2}$

$2 = C$

Therefore $y = \frac{2}{x}$

12e

$\frac{dy}{dx} = y\cos x$

$\frac{1}{y}\,dy = \cos x\;dx$

$\int \frac{1}{y}\,dy = \int \cos x\;dx$

$\ln|y| = \sin x + B$

$|y| = e^{(\sin x + B)}$

$|y| = e^{\sin x}e^B$

$y = e^B e^{\sin x}$ or $-e^B e^{\sin x}$

$y = Ce^{\sin x}$ where $C = e^B$ or $-e^B$

Substituting $y\left(\frac{\pi}{2}\right) = 1$:

$1 = Ce^{\sin\frac{\pi}{2}}$

$1 = Ce^1$

$C = \frac{1}{e}$

Therefore $y = \frac{1}{e}e^{\sin x}$

© Cambridge University Press 2019

$$= e^{\sin x} \times e^{-1}$$

$$= e^{\sin x - 1}$$

12f

$$\frac{dy}{dx} = \frac{y(2-x)}{x^2}$$

$$\frac{1}{y}\, dy = \frac{2-x}{x^2}\, dx$$

$$\int \frac{1}{y}\, dy = \int \frac{2-x}{x^2}\, dx$$

$$\int \frac{1}{y}\, dy = \int \left(\frac{2}{x^2} - \frac{x}{x^2}\right) dx$$

$$\int \frac{1}{y}\, dy = \int \left(\frac{2}{x^2} - \frac{1}{x}\right) dx$$

$$\ln|y| = \frac{2x^{-1}}{-1} - \ln|x| + C$$

$$\ln|y| = -\frac{2}{x} - \ln|x| + C$$

$$\ln|y| + \ln|x| = -\frac{2}{x} + C$$

$$\ln|xy| = -\frac{2}{x} + C$$

$$xy = e^{-\frac{2}{x}+C}$$

$$y = \frac{e^{-\frac{2}{x}+C}}{x}$$

Substituting $y(2) = \frac{1}{2}$:

$$\frac{1}{2} = \frac{e^{-\frac{2}{2}+C}}{2}$$

$$\frac{1}{2} = \frac{e^{-1+C}}{2}$$

$$1 = e^{-1-C}$$

$$-1 - C = 0$$

© Cambridge University Press 2019

$C = -1$

$y = \dfrac{e^{-\frac{2}{x}+1}}{x}$

$y = \dfrac{e^{1-\frac{2}{x}}}{x}$

13a

$$\frac{d(\log_e(\log_e x))}{dx}$$

$$= \frac{1}{x} \times \frac{1}{\log_e x}$$

$$= \frac{1}{x \log_e x}$$

13b $(x \log_e x)y' = y$

$$y' = \frac{y}{x \log_e x}$$

$$\frac{dy}{dx} = \frac{y}{x \log_e x}$$

$$\frac{1}{y}\, dy = \frac{1}{x \log_e x}\, dx$$

$$\int \frac{1}{y}\, dy = \int \frac{1}{x \log_e x}\, dx$$

$$\log_e|y| = \log_e(\log_e x) + B$$

$$\log_e|y| = \log_e(\log_e x) + \log_e e^B$$

$$\log_e|y| = \log_e(e^B \log_e x)$$

$$|y| = e^B \log_e x$$

$$y = e^B \log_e x \text{ or } -e^B \log_e x$$

$$y = C \log_e x \quad \text{where } C = e^B \text{ or } -e^B$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

14a

$$\frac{x}{x+2} = \frac{x+2-2}{x+2}$$

$$= \frac{x+2}{x+2} - \frac{2}{x+2}$$

$$= 1 - \frac{2}{x+2}$$

14b $(x+2)y' - xy = 0$

$$\frac{dy}{dx} = \frac{xy}{x+2}$$

$$\frac{1}{y}\,dy = \frac{x}{x+2}\,dx$$

$$\int \frac{1}{y}\,dy = \int \frac{x}{x+2}\,dx$$

$$\int \frac{1}{y}\,dy = \int \left(1 - \frac{2}{x+2}\right)dx$$

$$\ln|y| = x - 2\ln|x+2| + B$$

$$\ln|y| = \ln(e^x) - \ln(|x+2|^2) + \ln(e^B)$$

$$\ln|y| = \ln(e^x) - \ln((x+2)^2) + \ln(e^B)$$

$$\ln|y| = \ln\left(\frac{e^B e^x}{(x+2)^2}\right)$$

$$|y| = \frac{e^B e^x}{(x+2)^2}$$

$$y = \frac{e^B e^x}{(x+2)^2} \text{ or } -\frac{e^B e^x}{(x+2)^2}$$

$$y = \frac{Ce^x}{(x+2)^2} \text{ where } C = e^B \text{ or } -e^B$$

Substituting $y(0) = 1$:

$$1 = \frac{Ce^0}{(0+2)^2}$$

$$1 = \frac{C}{4}$$

© Cambridge University Press 2019

$C = 4$

Therefore $y = \dfrac{4e^x}{(x+2)^2}$

15a Since $\cos 2x = 2\cos^2 x - 1$

$2\cos^2 x = 1 + \cos 2x$

15b

$\dfrac{dy}{dx} = \dfrac{2\cos^2 x}{y}$

$y\,dy = 2\cos^2 x\;dx$

$\int y\,dy = \int 2\cos^2 x\,dx$

$\int y\,dy = \int (1 + \cos 2x)\,dx$

$\dfrac{y^2}{2} = x + \dfrac{1}{2}\sin 2x + C$

$y^2 = 2x + \sin 2x + 2C$

Substituting $y(0) = \sqrt{2}$:

$(\sqrt{2})^2 = 2(0) + \sin(0) + 2C$

$2C = 2$

Therefore $y^2 = 2x + \sin 2x + 2$

16a $y = x \times u$

$y' = u \times 1 + x \times u'$

$y' = u + xu'$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

16b i $xy' = 2x + 2y$

$$y' = \frac{2x + 2y}{x}$$

Substituting for y and y':

$$u + xu' = \frac{2x + 2xu}{x}$$

$$u + xu' = \frac{2x(1 + u)}{x}$$

$$u + xu' = 2(1 + u)$$

$$xu' = 2(1 + u) - u$$

$$xu' = 2 + 2u - u$$

$$xu' = 2 + u$$

$$u' = \frac{2 + u}{x}$$

16b ii

$$\frac{du}{dx} = \frac{2 + u}{x}$$

$$\frac{1}{2 + u}\, du = \frac{1}{x}\, dx$$

$$\int \frac{1}{2 + u}\, du = \int \frac{1}{x}\, dx$$

$$\ln|2 + u| = \ln|x| + B$$

$$\ln|2 + u| = \ln|x| + \ln(e^B)$$

$$\ln|2 + u| = \ln(e^B|x|)$$

$$|2 + u| = e^B|x|$$

$$2 + u = e^B x \text{ or } -e^B x$$

$$2 + u = Cx \text{ where } C = e^B \text{ or } -e^B$$

$$u = Cx - 2$$

© Cambridge University Press 2019

16b iii $u = Cx - 2$

From part a, $u = \frac{y}{x}$.

$\frac{y}{x} = Cx - 2$

$y = Cx^2 - 2x$

17a $(x^2 + 1)y' + (y^2 + 1) = 0$

$(x^2 + 1)\frac{dy}{dx} = -(y^2 + 1)$

$\frac{1}{y^2 + 1}\, dy = -\frac{1}{x^2 + 1}\, dx$

$\int \frac{1}{y^2 + 1}\, dy = -\int \frac{1}{x^2 + 1} dx$

$\tan^{-1} y = -\tan^{-1} x + C$

$\tan^{-1} y + \tan^{-1} x = C$

17b $\tan^{-1} y + \tan^{-1} x = C$

$\tan(\tan^{-1} y + \tan^{-1} x) = \tan C$

Using the formula for tangent of the sum of two angles,

$\frac{y + x}{1 - xy} = D$ where $D = \tan C$

17c Substituting $y(0) = 1$:

$\frac{1 + 0}{1 - 0} = D$

$D = 1$

Therefore $\frac{y + x}{1 - xy} = 1$

$y + x = 1 - xy$

$y + xy = 1 - x$

$y(1 + x) = 1 - x$

© Cambridge University Press 2019

$$y = \frac{1-x}{1+x}$$

18a

$$y_1 = \frac{1}{16}x^4$$

Substituting $x = 2$:

$$y_1(2) = \frac{1}{16} \times 2^4$$
$$= \frac{16}{16}$$
$$= 1$$

$$y_2 = \frac{1}{16}(x^2 - 8)^2$$

Substituting $x = 2$:

$$y_2(2) = \frac{1}{16}(2^2 - 8)^2$$
$$= \frac{1}{16}(-4)^2$$
$$= \frac{16}{16}$$
$$= 1$$

Both y_1 and y_2 satisfy the initial condition $y(2) = 1$.

18b

$$y_1' = \frac{d}{dx}\left(\frac{x^4}{16}\right)$$
$$= \frac{x^3}{4}$$

$$(y_1')^2 = \left(\frac{x^3}{4}\right)^2$$
$$= \frac{x^6}{16}$$

© Cambridge University Press 2019

$$= x^2 \times \frac{x^4}{16}$$

$$= x^2 y_1$$

$$y_2' = \frac{d}{dx}\left(\frac{1}{16}(x^2 - 8)^2\right)$$

$$= \frac{1}{16} \times 2(x^2 - 8) \times 2x$$

$$= \frac{x}{4}(x^2 - 8)$$

$$(y_2')^2 = \left(\frac{x}{4}(x^2 - 8)\right)^2$$


$$= \frac{x^2}{16}(x^2 - 8)^2$$

$$= x^2 \times \frac{1}{16}(x^2 - 8)^2$$

$$= x^2 y_2$$

Both y_1 and y_2 satisfy the DE $(y')^2 = x^2 y$.

18c



© Cambridge University Press 2019

18d $y_2' = xy_2^{\frac{1}{2}}$

So $y_2' = \frac{x}{4}(x^2 - 8)$ becomes

$xy_2^{\frac{1}{2}} = \frac{x}{4}\sqrt{(x^2-8)^2}$

$= \frac{x}{4}|x^2 - 8|$

When $x^2 - 8 < 0$ these two expressions are not equal.

To derive the solution correctly:

$\frac{dy}{dx} = xy^{\frac{1}{2}}$

$y^{-\frac{1}{2}}\,dy = x\,dx$

$\int y^{-\frac{1}{2}}\,dy = \int x\,dx$

$2y^{\frac{1}{2}} = \frac{x^2}{2} + B$

$y^{\frac{1}{2}} = \frac{x^2}{4} + C$ where $C = \frac{B}{2}$

Substituting $y(2) = 1$:

$1^{\frac{1}{2}} = \frac{2^2}{4} + C$

$1 = \frac{4}{4} + C$

$C = 0$

Hence the solution is

$y^{\frac{1}{2}} = \frac{x^2}{4}$ or $y = \frac{x^4}{16}$

19a The result follows from the fundamental theorem of calculus.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

19b i

$$\frac{dy}{dx} = -xy$$

$$\frac{1}{y}\,dy = -x\,dx$$

$$\int \frac{1}{y}\,dy = -\int x\,dx$$

$$\log|y| = -\frac{x^2}{2} + B$$

$$|y| = e^{-\frac{1}{2}x^2+B}$$

$$|y| = e^B e^{-\frac{1}{2}x^2}$$

$y = e^B e^{-\frac{1}{2}x^2}$ or $-e^B e^{-\frac{1}{2}x^2}$

$y = Ce^{-\frac{1}{2}x^2}$ where $C = e^B$ or $-e^B$

Substituting $y(0) = 1$:

$$1 = Ce^{-\frac{1}{2}\times 0^2}$$

$$1 = Ce^0$$

$$C = 1$$

Therefore $y = e^{-\frac{1}{2}x^2}$

19b ii

$$y_1(x) = y_0 + \int_0^x f(t, y_0)\,dt$$

$$= y_0 + \int_0^x (-y_0 t)\,dt$$

$$= 1 - \int_0^x t\,dt \qquad (\text{as } y_0 = 1)$$

$$= 1 - \left[\frac{t^2}{2}\right]_0^x$$

$$= 1 - \frac{1}{2}x^2$$

© Cambridge University Press 2019

$$y_2(x) = y_0 + \int_0^x f\big(t, y_1(t)\big)\, dt$$

$$= y_0 + \int_0^x (-y_1(t)\, t)\, dt$$

$$= 1 - \int_0^x \left(1 - \frac{1}{2}t^2\right) t\, dt$$

$$= 1 - \int_0^x \left(t - \frac{1}{2}t^3\right) dt$$

$$= 1 - \left[\frac{1}{2}t^2 - \frac{1}{8}t^4\right]_0^x$$

$$= 1 - \frac{1}{2}x^2 + \frac{1}{8}x^4$$

$$y_3(x) = y_0 + \int_0^x f\big(t, y_2(t)\big)\, dt$$

$$= y_0 + \int_0^x (-y_2(t)t\,)\, dt$$

$$= 1 - \int_0^x \left(1 - \frac{1}{2}t^2 + \frac{1}{8}t^4\right) t\, dt$$

$$= 1 - \int_0^x \left(t - \frac{1}{2}t^3 + \frac{1}{8}t^5\right) dt$$

$$= 1 - \left[\frac{1}{2}t^2 - \frac{1}{8}t^4 + \frac{1}{48}t^6\right]_0^x$$

$$= 1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6$$

$$y_4(x) = y_0 + \int_0^x f\big(t, y_3(t)\big)\, dt$$

$$= y_0 + \int_0^x (-y_3(t)t\,)dt$$

$$= 1 - \int_0^x \left(1 - \frac{1}{2}t^2 + \frac{1}{8}t^4 - \frac{1}{48}t^6\right) t\, dt$$

© Cambridge University Press 2019

$$= 1 - \int_0^x \left(t - \frac{1}{2}t^3 + \frac{1}{8}t^5 - \frac{1}{48}t^7\right) dt$$

$$= 1 - \left[\frac{1}{2}t^2 - \frac{1}{8}t^4 + \frac{1}{48}t^6 - \frac{1}{384}t^8\right]_0^x$$

$$= 1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \frac{1}{384}x^8$$

19b iii

$y\left(\frac{1}{2}\right) = e^{-\frac{1}{2}\left(\frac{1}{2}\right)^2} = e^{-\frac{1}{8}}$

Therefore $e = \left(y\left(\frac{1}{2}\right)\right)^{-8}$

Applying the formula above,

$y_4\left(\frac{1}{2}\right) \doteqdot 0.882\ 497\ 2$

and $e \doteqdot \left(y_4\left(\frac{1}{2}\right)\right)^{-8} \doteqdot 2.7183$

© Cambridge University Press 2019

Solutions to Exercise 13D

Let the integration constants be A, B, C or D.

1a $y = 0$

1b

$$\frac{dy}{dx} = -y$$

$$\left(\frac{1}{y}\right) dy = (-1)\, dx$$

$$\int \left(\frac{1}{y}\right) dy = \int (-1)\, dx$$

$$\ln|y| = -x + C$$

1c $\ln|y| = \ln e^{-x+C}$

$|y| = e^{-x+C}$

$|y| = e^{C} e^{-x}$

$y = e^{C} e^{-x}$ or $- e^{C} e^{-x}$

$y = Ae^{-x}$ where $A = e^{C}$ or $-e^{C}$

1d $y = 0$ when $A = 0$

1e Substituting $y(0) = 2$:

$2 = Ae^{-0}$

$A = 2$

Therefore $y = 2e^{-x}$

2a $y = 0$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

2b

$$\frac{dx}{dy} = \frac{1}{3y}$$

2c

$$\int \frac{dx}{dy}\, dy = \int \frac{1}{3y}\, dy$$

$$\int \frac{dx}{dy}\, dy = \frac{1}{3}\int \frac{1}{y}\, dy$$

$$x = \frac{1}{3}\ln|y| + C$$

2d

$$x - C = \frac{1}{3}\ln|y|$$

$$\ln|y| = 3(x - C)$$

$$|y| = e^{3(x-C)}$$

$$|y| = e^{-3C}e^{3x}$$

$y = e^{-3C}e^{3x}$ or $-e^{-3C}e^{3x}$

$y = Ae^{3x}$ where $A = e^{-3C}$ or $-e^{-3C}$

2e $y = 0$ when $A = 0$

2f Substituting $y(0) = -1$:

$$-1 = Ae^{3(0)}$$

$$A = -1$$

Therefore $y = -e^{3x}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

3a $y' - y = 0$

$y' = y$

$\frac{dy}{dx} = y$

$\frac{dx}{dy} = \frac{1}{y}$

$\int \frac{dx}{dy}\, dy = \int \frac{1}{y}\, dy$

$x = \ln|y| + C$

$x - C = \ln|y|$

$|y| = e^{x-C}$

$|y| = e^{-C}e^{x}$

$y = e^{-C}e^{x}$ or $-e^{-C}e^{x}$

$y = Ae^{x}$ where $A = e^{-C}$ or $-e^{-C}$

Substituting $y(0) = -3$:

$-3 = Ae^{0}$

$-3 = A$

Therefore $y = -3e^{x}$

3b $y' + 2y = 0$

$y' = -2y$

$\frac{dy}{dx} = -2y$

$\frac{dx}{dy} = -\frac{1}{2y}$

$\int \frac{dx}{dy}\, dy = -\frac{1}{2}\int \frac{1}{y}\, dy$

$x = -\frac{1}{2}\ln|y| + C$

$\ln|y| = 2C - 2x$

$|y| = e^{2C-2x}$

© Cambridge University Press 2019

$|y| = e^{2C}e^{-2x}$

$y = e^{2C}e^{-2x}$ or $-e^{2C}e^{-2x}$

$y = Ae^{-2x}$ where $A = e^{2C}$ or $-e^{2C}$

Substituting $y(0) = 1$:

$1 = Ae^0$

$1 = A$

Therefore $y = e^{-2x}$

3c $y' = -3y$

$\frac{dy}{dx} = -3y$

$\int \frac{dx}{dy}\, dy = -\frac{1}{3}\int \frac{1}{y}\, dy$

$x = -\frac{1}{3}\ln|y| + C$

$\ln|y| = 3C - 3x$

$|y| = e^{3C-3x}$

$|y| = e^{3C}e^{-3x}$

$y = e^{3C}e^{-3x}$ or $-e^{3C}e^{-3x}$

$y = Ae^{-3x}$ where $A = e^{3C}$ or $-e^{3C}$

Substituting $y(0) = 2$:

$2 = Ae^0$

$2 = A$

Therefore $y = 2e^{-3x}$

3d $y' = 2y$

$\frac{dy}{dx} = 2y$

$\frac{dx}{dy} = \frac{1}{2y}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

$$\int \frac{dx}{dy}\, dy = \frac{1}{2}\int \frac{1}{y}\, dy$$

$$x = \frac{1}{2}\ln|y| + C$$

$$2x - 2C = \ln|y|$$

$$|y| = e^{2x-2C}$$

$$|y| = e^{-2C}e^{2x}$$

$y = e^{-2C}e^{2x}$ or $-e^{-2C}e^{2x}$

$y = Ae^{2x}$ where $A = e^{-2C}$ or $-e^{-2C}$

Substituting $y(0) = -1$:

$-1 = Ae^0$

$-1 = A$

Therefore $y = -e^{2x}$

4a $y = 2$

4b

$$\frac{dy}{dx} = 2 - y$$

$$\frac{dy}{dx} = -1(y - 2)$$

$$\frac{1}{y-2}\, dy = (-1)\, dx$$

$$\int \frac{1}{y-2}\, dy = \int (-1)\, dx$$

$$\ln|y - 2| = -x + C$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4c $\ln|y-2| = -x + C$

$|y-2| = e^{-x+C}$

$y - 2 = e^C e^{-x}$ or $-e^C e^{-x}$

$y - 2 = Ae^{-x}$ where $A = e^C$ or $-e^C$

$y = 2 + Ae^{-x}$

4d $y = 2$ when $A = 0$

4e Substituting $y(0) = 3$:

$3 = 2 + Ae^0$

$1 = A$

$y = 2 + e^{-x}$

5a $y' = 1 - y$

$\frac{dy}{dx} = 1 - y$

$\frac{dy}{dx} = -1(y-1)$

$\frac{1}{y-1}\,dy = (-1)\,dx$

$\int \frac{1}{y-1}\,dy = \int (-1)\,dx$

$\ln|y-1| = -x + C$

$|y-1| = e^{-x+C}$

$y - 1 = e^C e^{-x}$ or $-e^C e^{-x}$

$y - 1 = Ae^{-x}$ where $A = e^C$ or $-e^C$

$y = 1 + Ae^{-x}$

Substituting $y(0) = 3$:

$3 = 1 + Ae^0$

$3 = 1 + A$

© Cambridge University Press 2019

$2 = A$

Therefore $y = 1 + 2e^{-x}$

5b $y' = y - 1$

$\frac{dy}{dx} = y - 1$

$\frac{1}{y-1}\,dy = 1\,dx$

$\int \frac{1}{y-1}\,dy = \int 1\,dx$

$\ln|y-1| = x + C$

$|y-1| = e^{x+C}$

$y - 1 = e^C e^x$ or $-e^C e^x$

$y - 1 = Ae^x$ where $A = e^C$ or $-e^C$

$y = 1 + Ae^x$

Substituting $y(0) = 0$:

$0 = 1 + Ae^0$

$0 = 1 + A$

$-1 = A$

Therefore $y = 1 - e^x$

5c $y' = \frac{1}{2}(y+1)$

$\frac{dy}{dx} = \frac{1}{2}(y+1)$

$\frac{1}{y+1}\,dy = \frac{1}{2}\,dx$

$\int \frac{1}{y+1}\,dy = \int \frac{1}{2}\,dx$

$\ln|y+1| = \frac{1}{2}x + C$

$|y+1| = e^{\frac{1}{2}x+C}$

© Cambridge University Press 2019

$y + 1 = e^C e^{\frac{1}{2}x}$ or $- e^C e^{\frac{1}{2}x}$

$y + 1 = Ae^{\frac{1}{2}x}$ where $A = e^C$ or $-e^C$

$y = -1 + Ae^{\frac{1}{2}x}$

Substituting $y(0) = 1$:

$1 = -1 + Ae^0$

$1 = -1 + A$

$2 = A$

Therefore $y = -1 + 2e^{\frac{1}{2}x}$

5d $y' = 2(3 - y)$

$\frac{dy}{dx} = 2(3 - y)$

$\frac{dy}{dx} = -2(y - 3)$

$\frac{1}{y - 3}\, dy = (-2)\, dx$

$\int \frac{1}{y - 3}\, dy = \int (-2)\, dx$

$\ln|y - 3| = -2x + C$

$|y - 3| = e^{-2x+C}$

$y - 3 = e^C e^{-2x}$ or $- e^C e^{-2x}$

$y - 3 = Ae^{-2x}$ where $A = e^C$ or $-e^C$

$y = 3 + Ae^{-2x}$

Substituting $y(0) = 4$:

$4 = 3 + Ae^0$

$4 = 3 + A$

$1 = A$

Therefore $y = 3 + e^{-2x}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

6a $y' = 2y^2$

$\frac{dy}{dx} = 2y^2$

$\int \frac{1}{y^2}\, dy = \int 2\, dx$

$-y^{-1} = 2x + C$

$-\frac{1}{y} = 2x + C$

Substituting $y(0) = 3$:

$-\frac{1}{3} = 0 + C$

$C = -\frac{1}{3}$

Therefore $-\frac{1}{y} = 2x - \frac{1}{3}$

$-\frac{1}{y} = \frac{6x - 1}{3}$

$-\frac{y}{1} = \frac{3}{6x - 1}$

$y = -\frac{3}{6x - 1}$

$y = \frac{3}{1 - 6x}$

6b $y' = -y^2$

$\frac{dy}{dx} = -y^2$

$\int \left(-\frac{1}{y^2}\right) dy = \int 1\, dx$

$y^{-1} = x + C$

$\frac{1}{y} = x + C$

© Cambridge University Press 2019

Substituting $y(0) = 1$:

$\frac{1}{1} = 0 + C$

$C = 1$

Therefore $\frac{1}{y} = x + 1$

$\frac{y}{1} = \frac{1}{x+1}$

$y = \frac{1}{x+1}$

6c $y' = 1 + y^2$

$\frac{dy}{dx} = 1 + y^2$

$\int \frac{1}{1+y^2}\, dy = \int 1\, dx$

$\tan^{-1} y = x + C$

Substituting $y\left(\frac{\pi}{4}\right) = 1$:

$\tan^{-1} 1 = \frac{\pi}{4} + C$

$\frac{\pi}{4} = \frac{\pi}{4} + C$

$0 = C$

Therefore $\tan^{-1} y = x$

$y = \tan x$

6d $y' = -e^y$

$\frac{dy}{dx} = -e^y$

$\int \left(-\frac{1}{e^y}\right) dy = \int 1\, dx$

$\int (-e^{-y})\, dy = \int 1\, dx$

© Cambridge University Press 2019

$e^{-y} = x + C$

$\ln(e^{-y}) = \ln(x + C)$

$-y \ln e = \ln(x + C)$

$-y = \ln(x + C)$

$y = -\ln(x + C)$

Substituting $y(0) = 0$:

$0 = -\ln C$

$C = e^0$

$C = 1$

Therefore $y = -\ln(x + 1)$

6e $y' = e^{-y}$

$\frac{dy}{dx} = e^{-y}$

$\int \frac{1}{e^{-y}}\, dy = \int 1\, dx$

$\int e^y\, dy = \int 1\, dx$

$e^y = x + C$

$\ln e^y = \ln(x + C)$

$y = \ln(x + C)$

Substituting $y(3) = 0$:

$0 = \ln(3 + C)$

$e^0 = 3 + C$

$1 = 3 + C$

$C = -2$

Therefore $y = \ln(x - 2)$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

6f $y' = y^{\frac{2}{3}}$

$\frac{dy}{dx} = y^{\frac{2}{3}}$

$\int \frac{1}{y^{\frac{2}{3}}}\, dy = \int 1\, dx$

$\int y^{-\frac{2}{3}}\, dy = \int 1\, dx$

$\frac{y^{\frac{1}{3}}}{\frac{1}{3}} = x + C$

$3y^{\frac{1}{3}} = x + C$

Substituting $y(0) = 1$:

$3(1^{\frac{1}{3}}) = 0 + C$

$C = 3$

Therefore $3y^{\frac{1}{3}} = x + 3$

$y^{\frac{1}{3}} = \frac{x+3}{3}$

$y = \left(\frac{x+3}{3}\right)^3$

7a $y' = ky$

$\frac{dy}{dx} = ky$

$\int \frac{1}{y}\, dy = \int k\, dx$

$\ln|y| = kx + C$

$|y| = e^{kx+C}$

$y = e^C e^{kx}$ *or* $-e^C e^{kx}$

$y = Ae^{kx}$ where $A = e^C$ or $-e^C$

© Cambridge University Press 2019

7b Substituting $y(0) = 20$:

$20 = Ae^{k(0)}$

$A = 20$

7c $y = 20e^{kx}$

Substituting $y(2) = 5$:

$5 = 20e^{k(2)}$

$\frac{1}{4} = e^{2k}$

$2k = \ln\left(\frac{1}{4}\right)$

$k = \frac{1}{2}\ln\left(\frac{1}{4}\right)$

$k = \frac{1}{2}\ln(2^{-2})$

$k = -2 \times \frac{1}{2}\ln 2$

$k = -\ln 2$

7d $y = 20e^{(-\ln 2)x}$

$y = 20e^{-x\ln 2}$

$y = 20e^{\ln 2^{-x}}$

$y = 20 \times 2^{-x}$

$y(3) = 20 \times 2^{-3}$

$= \frac{20}{8}$

$= 2\frac{1}{2}$

8a $y' = ky$

$\frac{dy}{dx} = ky$

© Cambridge University Press 2019

$$\int \frac{1}{y}\,dy = \int k\,dx$$

$$\ln|y| = kx + C$$

$$|y| = e^{kx+C}$$

$$y = e^{C}e^{kx} \text{ or } -e^{C}e^{kx}$$

$y = Ae^{kx}$ where $A = e^{C}$ or $-e^{C}$

8b Substituting $y(0) = 8$:

$$8 = Ae^{k(0)}$$

$$A = 8$$

8c $y = 8e^{kx}$

Substituting $y(2) = 18$:

$$18 = 8e^{k(2)}$$

$$\frac{18}{8} = e^{2k}$$

$$2k = \ln\left(\frac{9}{4}\right)$$

$$k = \frac{1}{2}\ln\left(\frac{9}{4}\right)$$

$$k = \frac{1}{2}\ln\left(\frac{3}{2}\right)^{2}$$

$$k = 2 \times \frac{1}{2}\ln\frac{3}{2}$$

$$k = \ln\frac{3}{2}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

8d

$$y = 8e^{(\ln\frac{3}{2})x}$$

$$y = 8e^{x\ln\frac{3}{2}}$$

$$y = 8e^{\ln\left(\frac{3}{2}\right)^x}$$

$$y = 8 \times \left(\frac{3}{2}\right)^x$$

$$y(4) = 8 \times \left(\frac{3}{2}\right)^4$$

$$= 8 \times \frac{81}{16}$$

$$= 40\frac{1}{2}$$

9a $y = Ae^{\lambda x} + Be^{-\lambda x} + C\cos\lambda x + D\sin\lambda x$

$$y' = A\lambda e^{\lambda x} - B\lambda e^{-\lambda x} - C\lambda\sin\lambda x + D\lambda\cos\lambda x$$

$$y'' = A\lambda^2 e^{\lambda x} + B\lambda^2 e^{-\lambda x} - C\lambda^2\cos\lambda x - D\lambda^2\sin\lambda x$$

$$y''' = A\lambda^3 e^{\lambda x} - B\lambda^3 e^{-\lambda x} + C\lambda^3\sin\lambda x - D\lambda^3\cos\lambda x$$

$$y'''' = A\lambda^4 e^{\lambda x} + B\lambda^4 e^{-\lambda x} + C\lambda^4\cos\lambda x + D\lambda^4\sin\lambda x$$

$$\text{LHS} = y^{(4)}$$

$$= A\lambda^4 e^{\lambda x} + B\lambda^4 e^{-\lambda x} + C\lambda^4\cos\lambda x + D\lambda^4\sin\lambda x$$

$$\text{RHS} = \lambda^4 y$$

$$= \lambda^4(Ae^{\lambda x} + Be^{-\lambda x} + C\cos\lambda x + D\sin\lambda x)$$

$$= A\lambda^4 e^{\lambda x} + B\lambda^4 e^{-\lambda x} + C\lambda^4\cos\lambda x + D\lambda^4\sin\lambda x$$

LHS = RHS

Therefore y is a solution of the DE.

© Cambridge University Press 2019

9b Substituting $y(0) = 0$:

$0 = Ae^{\lambda(0)} + Be^{-\lambda(0)} + C\cos\lambda(0) + D\sin\lambda(0)$

$0 = A + B + C$ (1)

Substituting $y''(0) = 0$:

$0 = A\lambda^2 e^{\lambda(0)} + B\lambda^2 e^{-\lambda(0)} - C\lambda^2\cos\lambda(0) - D\lambda^2\sin\lambda(0)$

$0 = A\lambda^2 + B\lambda^2 - C\lambda^2$

$0 = \lambda^2(A + B - C)$

$0 = A + B - C$ where $\lambda \neq 0$ (2)

$(1) - (2)$:

$2C = 0$

$C = 0$

9c i Substituting $y(10) = 0$ and $C = 0$:

$0 = Ae^{\frac{n\pi}{10}(10)} + Be^{-\frac{n\pi}{10}(10)} + D\sin\left(\frac{n\pi}{10}(10)\right)$

$0 = Ae^{n\pi} + Be^{-n\pi} + D\sin n\pi$ (1)

Substituting $y''(10) = 0$ and $C = 0$:

$0 = A\lambda^2 e^{10\lambda} + B\lambda^2 e^{-10\lambda} - C\lambda^2\cos 10\lambda - D\lambda^2\sin 10\lambda$

$0 = \lambda^2(Ae^{n\pi} + Be^{-n\pi} - D\sin n\pi)$

$0 = Ae^{n\pi} + Be^{-n\pi} - D\sin n\pi$ where $\lambda \neq 0$ (2)

$(1) + (2)$:

$2Ae^{n\pi} + 2Be^{-n\pi} + D\sin n\pi - D\sin n\pi = 0$

$2Ae^{n\pi} + 2Be^{-n\pi} = 0$

$Ae^{n\pi} + Be^{-n\pi} = 0$

$Ae^{n\pi} = -Be^{-n\pi}$

$A = -Be^{-2n\pi}$ (3)

From part b above, $A + B - C = 0$ and $C = 0$

Therefore $A + B = 0$

$A = -B$ (4)

© Cambridge University Press 2019

Combining (3) and (4):

$-B = -Be^{-2n\pi}$

$B = Be^{-2n\pi}$

Either $B = 0$ or $e^{-2n\pi} = 1$, i.e. $n = 0$.

If $n = 0$ then $\lambda = 0$ and the beam equation simplifies to $y = A + B = 0$.

Without loss of generality we can then set $A = B = 0$.

If $n \neq 0$ then $B = 0$ and therefore $A = 0$.

9c ii Since $A = B = C = 0$,

$y = D \sin\left(\frac{n\pi}{10}x\right)$

10a $y' = e^{-y}$

$\frac{dy}{dx} = e^{-y}$

$\int \frac{1}{e^{-y}}\, dy = \int 1\, dx$

$\int e^{y}\, dy = \int 1\, dx$

$e^{y} = x + C$

$\ln e^{y} = \ln(x + C)$


$y = \ln(x + C)$

10b Log curves with different x-intercepts

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

10c i,ii



10c iii Shift left or right

10c iv The isoclines are horizontal lines.

10d $y = \ln(x + C)$

Substituting $(0, 1)$:

$1 = \ln(0 + C)$

$e^1 = 0 + C$

$C = e$

Therefore $y = \ln(x + e)$

11a $y'y = 2$

$\frac{dy}{dx}y = 2$

$\int y\,dy = \int 2\,dx$

$\frac{y^2}{2} = 2x + B$

$y^2 = 4x + 2B$


$y^2 = 4x + C$ where $C = 2B$

11b Concave right parabolas with vertex on the x-axis.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

11c i, ii



11c iii Shift left or right

11c iv The isoclines are horizontal lines.

11d $y^2 = 4x + C$

Substituting $(0, 1)$:

$1 = 4(0) + C$

$C = 1$

Therefore $y^2 = 4x + 1$

12a

$$L(x) = \frac{1}{1 + e^{-x}}$$

For y-intercept, $x = 0$.

$$L(0) = \frac{1}{1 + e^{0}}$$

$$L(0) = \frac{1}{2}$$

y-intercept at $\left(0, \frac{1}{2}\right)$

12b $e^{-x} > 0$ for all x so $L(x)$ is always positive

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

12c Since $e^{-x} \to 0$ as $x \to \infty$, $\lim_{x\to\infty} L(x) = 1$

and $e^{-x} \to \infty$ as $x \to -\infty$, so $\lim_{x\to-\infty} L(x) = 0$

12d $L(x) = (1 + e^{-x})^{-1}$

$L'(x) = -(1 + e^{-x})^{-2} \times -e^{-x}$

$$= \frac{e^{-x}}{(1 + e^{-x})^2}$$

For stationary points to exist, $L'(x) = 0$.

$$\frac{e^{-x}}{(1 + e^{-x})^2} = 0$$

$e^{-x} = 0$ which has no solution

Therefore, there are no stationary points.

12e i $L'(x)$

$$= \frac{e^{-x}}{(1 + e^{-x})^2}$$

$$= \frac{1}{e^x(1 + e^{-x})^2}$$

$$= \frac{1}{e^x(1 + 2e^{-x} + e^{-2x})}$$

$$= \frac{1}{(e^x + 2 + e^{-x})}$$

$$= \frac{1}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^2}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

12e ii

$$L'(x) = \left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{-2}$$

$$L''(x) = -2\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{-3} \times \left(\frac{1}{2}e^{\frac{x}{2}} - \frac{1}{2}e^{-\frac{x}{2}}\right)$$

$$= -2\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{-3} \times \frac{1}{2}\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)$$

$$= -\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{-3} \times \left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)$$

$$= -\frac{\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{3}}$$

12e iii Point of inflection when $L''(x) = 0$.

$$-\frac{\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{3}} = 0$$

$$e^{\frac{x}{2}} - e^{-\frac{x}{2}} = 0$$

$$e^{-\frac{x}{2}}(e^{-x} - 1) = 0$$

$e^{-\frac{x}{2}} = 0 \implies$ No solution

$$e^{-x} - 1 = 0$$

$$e^{-x} = 1$$

$$-x = \ln 1$$

$$x = 0$$


$$L(0) = (1 + e^0)^{-1} = (2)^{-1} = \frac{1}{2}$$

POI $= \left(0, \frac{1}{2}\right)$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

12f



12g i $y' = y(1 - y)$

$$\text{LHS} = y' = L'(x) = \frac{e^{-x}}{(1 + e^{-x})^2} \qquad \text{(from part a)}$$

$$\text{RHS} = y(1 - y)$$

$$= \left(\frac{1}{1 + e^{-x}}\right)\left(1 - \frac{1}{1 + e^{-x}}\right)$$

$$= \left(\frac{1}{1 + e^{-x}}\right)\left(\frac{1 + e^{-x} - 1}{1 + e^{-x}}\right)$$

$$= \left(\frac{1}{1 + e^{-x}}\right)\left(\frac{e^{-x}}{1 + e^{-x}}\right)$$

$$= \frac{e^{-x}}{(1 + e^{-x})^2}$$

Therefore, since LHS = RHS, $L(x)$ is a solution.

12g ii $y' = y(1 - y)$

$$= \left(\frac{1}{1 + e^{-x}}\right)\left(1 - \frac{1}{1 + e^{-x}}\right)$$

$$= \left(\frac{1}{1 + e^{-x}}\right)\left(\frac{1 + e^{-x} - 1}{1 + e^{-x}}\right)$$

$$= \left(\frac{1}{1 + e^{-x}}\right)\left(\frac{e^{-x}}{1 + e^{-x}}\right)$$

$$= \frac{e^{-x}}{(1 + e^{-x})^2}$$

$$= \frac{1}{e^{x}(1 + e^{-x})^2}$$

© Cambridge University Press 2019

$$= \frac{1}{e^x(1 + 2e^{-x} + e^{-2x})}$$

$$= \frac{1}{(e^x + 2 + e^{-x})}$$

$$= \frac{1}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^2}$$

12g iii $y' = y(1 - y) = y - y^2$

$$y'' = \frac{d}{dy}(y - y^2) \times \frac{dy}{dx}$$

$$= (1 - 2y) \times y'$$

$$= \left(1 - 2\left(\frac{1}{1 + e^{-x}}\right)\right) \times \left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{-2}$$

$$= \left(\frac{e^{-x} - 1}{1 + e^{-x}}\right) \times \left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^{-2}$$

$$= \left(\frac{e^{-x} - 1}{1 + e^{-x}}\right) \times \frac{1}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^2}$$

$$= \left(\frac{1 - e^{-x}}{1 + e^{-x}}\right)\left(\frac{e^{\frac{x}{2}}}{e^{\frac{x}{2}}}\right) \times \frac{-1}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^2}$$

$$= \frac{\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)} \times \frac{-1}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^2}$$

$$= -\frac{\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)}{\left(e^{\frac{x}{2}} + e^{-\frac{x}{2}}\right)^3}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

13a

Let $\frac{1}{y(1-y)} = \frac{A}{y} + \frac{B}{1-y}$

$$\text{RHS} = \frac{A(1-y)+By}{y(1-y)}$$

$$= \frac{A - Ay + By}{y(1-y)}$$

$$= \frac{A + y(-A+B)}{y(1-y)}$$

Equating coefficients of numerators in LHS and RHS:

$A = 1$

$-A + B = 0$

$-1 + B = 0$

$B = 1$

Therefore $\frac{1}{y(1-y)} = \frac{1}{y} + \frac{1}{1-y}$

13b i $y = 0$ and $y = 1$

13b ii $y' = y(1-y)$

$$\frac{dy}{dx} = y(1-y)$$

$$\int \frac{1}{y(1-y)}\,dy = \int 1\ dx$$

$$\int \left(\frac{1}{y} + \frac{1}{1-y}\right) dy = \int 1\ dx$$

$$\ln|y| - \ln|1-y| = x + C$$

$$\ln\left|\frac{y}{1-y}\right| = x + C$$

$$\left|\frac{y}{1-y}\right| = e^{x+C}$$

© Cambridge University Press 2019

$$\left|\frac{y}{1-y}\right| = e^C \times e^x$$

$$\frac{y}{1-y} = e^C \times e^x \text{ or } -e^C \times e^x$$

$$\frac{y}{1-y} = Ae^x \text{ where } A = e^C \text{ or } -e^C$$

$$y = Ae^x(1-y)$$

$$y = Ae^x - yAe^x$$

$$y + yAe^x = Ae^x$$

$$y(1 + Ae^x) = Ae^x$$

$$y = \frac{Ae^x}{1 + Ae^x}$$

$$y = \frac{1}{A^{-1}e^{-x}(1 + Ae^x)}$$

$$y = \frac{1}{A^{-1}e^{-x} + 1}$$

$$y = \frac{1}{Be^{-x} + 1} \text{ where } B = A^{-1}$$

$$y = \frac{1}{1 + Be^{-x}}$$

13c

$$L(x) = \frac{1}{1 + e^{-x}} \longrightarrow y = \frac{1}{1 + Be^{-x}}$$

$$y = \frac{1}{1 + Be^{-x}}$$

$$= \frac{1}{1 + e^{\ln B}e^{-x}}$$

$$= \frac{1}{1 + e^{-x + \ln B}}$$

$$= \frac{1}{1 + e^{-(x - \ln B)}}$$

Therefore there is a shift of $\ln B$ to the right.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

13d i

$$\lim_{B\to\infty} \frac{1}{1+Be^{-x}} = 0$$

Yes, this is one of the constant solutions.

13d ii

$$\lim_{B\to 0^+} \frac{1}{1+Be^{-x}} = 1$$

Yes, this is one of the constant solutions.

14a $y = 0$ and $y = 1$

14b $y' = ry(1-y)$

$$\frac{dy}{dx} = ry(1-y)$$

$$\int \frac{1}{y(1-y)}\, dy = \int r\ dx$$

$$\int \left(\frac{1}{y} + \frac{1}{1-y}\right) dy = \int r\ dx$$

$$\ln|y| - \ln|1-y| = rx + C$$

$$\ln\left|\frac{y}{1-y}\right| = rx + C$$

$$\left|\frac{y}{1-y}\right| = e^{rx+C}$$

$$\left|\frac{y}{1-y}\right| = e^{C} \times e^{rx}$$

$$\frac{y}{1-y} = e^{C} \times e^{rx} \text{ or } -e^{C} \times e^{rx}$$

$$\frac{y}{1-y} = Ae^{rx} \text{ where } A = e^{C} \text{ or } -e^{C}$$

$$y = Ae^{rx}(1-y)$$

$$y = Ae^{rx} - yAe^{rx}$$

$$y + yAe^{rx} = Ae^{rx}$$

© Cambridge University Press 2019

$$y(1 + Ae^{rx}) = Ae^{rx}$$

$$y = \frac{Ae^{rx}}{1 + Ae^{rx}}$$

$$y = \frac{1}{A^{-1}e^{-rx}(1 + Ae^{rx})}$$

$$y = \frac{1}{A^{-1}e^{-rx} + 1}$$

$$y = \frac{1}{Be^{-rx} + 1} \text{ where } B = A^{-1}$$

$$y = \frac{1}{1 + Be^{-rx}}$$

14c Substituting $y(0) = y_0$:

$$y_0 = \frac{1}{1 + Be^{-r(0)}}$$

$$y_0 = \frac{1}{1 + B}$$

$$1 + B = \frac{1}{y_0}$$

$$B = \frac{1}{y_0} - 1$$

14d

$$y = \frac{1}{1 + Be^{-rx}}$$

$$= \frac{1}{1 + e^{\ln B}e^{-rx}}$$

$$= \frac{1}{1 + e^{-rx + \ln B}}$$

$$= \frac{1}{1 + e^{-r\left(x - \frac{1}{r}\ln B\right)}}$$

Therefore, there is a shift of $\frac{1}{r}\ln B$ to the right.

© Cambridge University Press 2019

14e i

$$B = \frac{1}{y_0} - 1$$


As $y_0 \to 0^+, B \to \infty$ so $y = 0$ in the limit.

14e ii

$$B = \frac{1}{y_0} - 1$$

As $y_0 \to 1^-, B \to 0^+$ so $y = 1$ in the limit.

15a,b,d



15c

$$y = \frac{1}{1 - e^{-x}}$$

$$= (1 - e^{-x})^{-1}$$

$$y' = (1 - e^{-x})^{-2} \times -e^{-x}$$

$$= -e^{-x}(1 - e^{-x})^{-2}$$

$$y' = y(1 - y)$$

$$\text{RHS} = \frac{1}{1 - e^{-x}}\left(1 - \frac{1}{1 - e^{-x}}\right)$$

$$= \frac{1}{1 - e^{-x}}\left(\frac{1 - e^{-x} - 1}{1 - e^{-x}}\right)$$

$$= -e^{-x}(1 - e^{-x})^{-2}$$

© Cambridge University Press 2019

$= \text{LHS}$

So $y = \dfrac{1}{1 - e^{-x}}$ is a solution of the DE $y' = y(1 - y)$.

16a

Let $\dfrac{1}{(y-1)(y-3)} = \dfrac{A}{y-1} + \dfrac{B}{y-3}$

$$\text{RHS} = \frac{A(y-3) + B(y-1)}{(y-1)(y-3)}$$

$$= \frac{Ay - 3A + By - B}{(y-1)(y-3)}$$

$$= \frac{y(A+B) - 3A - B}{(y-1)(y-3)}$$

Equating coefficients of numerators in LHS and RHS:

$A + B = 0 \qquad (1)$

$-3A - B = 1 \qquad (2)$

$(1) + (2)$:
$-2A = 1$

$A = -\dfrac{1}{2}$

Substituting $A = -\frac{1}{2}$ into (1):

$-\dfrac{1}{2} + B = 0$

$B = \dfrac{1}{2}$

Therefore

$$\frac{1}{(y-1)(y-3)} = \frac{-\frac{1}{2}}{y-1} + \frac{\frac{1}{2}}{y-3}$$

$$= \frac{1}{2(y-3)} - \frac{1}{2(y-1)}$$

$$= \frac{1}{2}\left(\frac{1}{(y-3)} - \frac{1}{(y-1)}\right)$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

16b i $y = 1$ and $y = 3$

16b ii $y' = -(1-y)(3-y)$

$$\frac{dy}{dx} = -(1-y)(3-y)$$

$$\int \frac{1}{(1-y)(3-y)}\,dy = \int (-1)\,dx$$

$$\int \frac{1}{(y-1)(y-3)}\,dy = \int (-1)\,dx$$

$$\frac{1}{2}\int \left(\frac{1}{(y-3)} - \frac{1}{(y-1)}\right)dy = \int (-1)\,dx$$

$$\int \left(\frac{1}{(y-3)} - \frac{1}{(y-1)}\right)dy = \int (-2)\,dx$$

$$\ln|y-3| - \ln|y-1| = -2x + C$$

$$\ln\left|\frac{y-3}{y-1}\right| = -2x + C$$

$$\left|\frac{y-3}{y-1}\right| = e^{-2x+C}$$

$$\left|\frac{y-3}{y-1}\right| = e^{C} \times e^{-2x}$$

$$\frac{y-3}{y-1} = e^{C} \times e^{-2x} \text{ or } -e^{C} \times e^{-2x}$$

$$\frac{y-3}{y-1} = Ae^{-2x} \text{ where } A = e^{C} \text{ or } -e^{C}$$

$$y - 3 = Ae^{-2x}(y-1)$$

$$y - 3 = yAe^{-2x} - Ae^{-2x}$$

$$y - yAe^{-2x} = 3 - Ae^{-2x}$$

$$y(1 - Ae^{-2x}) = 3 - Ae^{-2x}$$

$$y = \frac{3 - Ae^{-2x}}{1 - Ae^{-2x}}$$

$$y = \frac{3 + Be^{-2x}}{1 + Be^{-2x}} \text{ where } B = -A$$


© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

16c i As $B \to 0^+$, $y \to 3$, so $y = 3$ is the constant solution.

16c ii As $B \to \infty$, $y \to 1$, so $y = 1$ is the constant solution.

16d



16e i $y' = -(1-y)(3-y)$

$$y'' = -\frac{d}{dy}(1-y)(3-y) \times \frac{dy}{dx}$$

$$= -\frac{d}{dy}(3-4y+y^2) \times y'$$

$$= -(-4+2y) \times -(1-y)(3-y)$$

$$= 2(-2+y)(1-y)(3-y)$$

$$= -2(2-y)(1-y)(3-y)$$

$$= -2(1-y)(2-y)(3-y)$$

16e ii For points of inflection, $y'' = 0$

$-2(1-y)(2-y)(3-y) = 0$

$y = 1, 2$ or 3

However, $y = 1$ and $y = 3$ are constant solutions so only point of inflection at $y = 2$.

For $B = 1$

© Cambridge University Press 2019

$$y = \frac{3 + e^{-2x}}{1 + e^{-2x}}$$

$$2 = \frac{3 + e^{-2x}}{1 + e^{-2x}}$$

$$2 + 2e^{-2x} = 3 + e^{-2x}$$

$$e^{-2x} = 1$$

$$x = 0$$

Point of inflection at $(0, 2)$.

17a

$$v = \frac{1}{y}$$

$$v' = -\frac{1}{y^2}y'$$

$$= -v^2 y' \qquad \left(\text{as } v^2 = \frac{1}{y^2}\right)$$

$$= -v^2 ry(1 - y) \qquad (\text{as } y' = ry(1 - y))$$

$$= -v^2 ry^2\left(\frac{1 - y}{y}\right)$$

$$= -rv^2 y^2\left(\frac{1}{y} - 1\right)$$

$$= -r\left(\frac{1}{y} - 1\right) \qquad (\text{as } v^2 y^2 = 1)$$

$$= -r(v - 1)$$

$$= r(1 - v)$$

17b $v' = r(1 - v)$

$$\frac{dv}{dx} = r(1 - v)$$

$$\frac{dv}{dx} = -r(v - 1)$$

$$\int \frac{1}{v - 1}dy = \int (-r)\,dx$$

© Cambridge University Press 2019

$\ln|v-1| = -rx + C$

$|v-1| = e^{-rx+C}$

$|v-1| = e^{C}e^{-rx}$

$v-1 = e^{C}e^{-rx}$ or $-e^{C}e^{-rx}$

$v-1 = Be^{-rx}$ where $B = e^{C}$ or $-e^{C}$

$v = 1 + Be^{-rx}$

17c

$v = \frac{1}{y}$

$y = \frac{1}{v}$

$y = \frac{1}{1+Be^{-rx}}$

18a $y'' = 2(1-y')$

$v' = 2(1-v)$

18b $y'(0) = 0$ so $v(0) = 0$

18c $v' = 2(1-v)$

$\frac{dv}{dx} = 2(1-v)$

$\frac{dv}{dx} = -2(v-1)$

$\int \frac{1}{v-1}dy = \int (-2)\,dx$

$\ln|v-1| = -2x + C$

$|v-1| = e^{-2x+C}$

$|v-1| = e^{C}e^{-2x}$

$v-1 = e^{C}e^{-2x}$ or $-e^{C}e^{-2x}$

© Cambridge University Press 2019

$v - 1 = Be^{-2x}$ where $B = e^C$ or $-e^C$

$v = 1 + Be^{-2x}$

(Alternatively, use $r = 2$ in your answer to Q17b.)

Substituting $v(0) = 0$ from part b:

$0 = 1 + Be^{-2(0)}$

$0 = 1 + B$

$-1 = B$

Therefore $v = 1 - e^{-2x}$

18d Since $v = y'$,

$y' = 1 - e^{-2x}$

18e

$\int y'\,dx = \int (1 - e^{-2x})dx$

$y = x + \frac{1}{2}e^{-2x} + C$

Substituting $y(0) = 1$:

$1 = 0 + \frac{1}{2}e^{-2(0)} + C$

$1 = \frac{1}{2} + C$

$\frac{1}{2} = C$

Therefore $y = x + \frac{1}{2}e^{-2x} + \frac{1}{2}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

19a We are given:

$$\frac{df(x)}{dx} = g\big(f(x)\big)$$

Substitute $u = x - C$

Note that $\frac{du}{dx} = 1$

If $y = f(x - C)$

$y = f(u)$

$$y' = \frac{df(u)}{du} \times \frac{du}{dx}$$

$$= \frac{df(u)}{du} \times 1$$

$$= \frac{df(u)}{du}$$

$$= g\big(f(u)\big)$$

$$= g\big(f(x - C)\big)$$

$$= g\big(f(y)\big)$$

So $y = f(x - C)$ is also a solution.

19b Shift right by C

19c Since translating a solution horizontally gives another solution, all horizontal lines are isoclines.

19d If a graph is shifted right, its gradient at a given height is unchanged by the shift.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

20a i

$$\frac{d^2y}{dx^2} = \frac{d}{dx}\left(\frac{dy}{dx}\right)$$

$$= \frac{dv}{dx} \qquad \left(\text{as } v = \frac{dy}{dx}\right)$$

$$= \frac{dv}{dy} \times \frac{dy}{dx}$$

$$= \frac{dv}{dy} \times v$$

$$= v\frac{dv}{dy}$$

20a ii

$$\frac{d^2y}{dx^2} + y = 0$$

$$v\frac{dv}{dy} + y = 0$$

$$v\frac{dv}{dy} = -y$$

20b

$$v\frac{dv}{dy} = -y$$

$$v\,dv = -y\,dy$$

$$\int v\,dv = -\int y\,dy$$

$$\frac{v^2}{2} = -\frac{y^2}{2} + B$$

$$v^2 = -y^2 + 2B$$

$$v^2 + y^2 = C \text{ where } C = 2B$$

© Cambridge University Press 2019

20c If C is negative, there is no solution.

If C is zero, the solution is a single point, the origin.

If C is positive, the solution is a circle with centre the origin and radius $\sqrt{C}$.

20d Standard parametrisation for a circle: $v = r\cos x\,, y = r\sin x$

20e $y = r\sin x$

$$y' = \frac{d}{dx}(r\sin x)$$
$$= r\cos x$$
$$= v$$

20f The original DE is autonomous. From Question 19 we know that if $y = f(x)$ is a solution to the DE, then $y = f(x - D)$ is also a solution.

Hence the general solution is $y = r\sin(x - D)$.

20g i $y = r\sin(x - D)$

$= r\cos D\sin x - r\sin D\cos x$

$= A\cos x + B\sin x$ where $A = -r\sin D$ and $B = r\cos D$

20g ii $y = A\cos x + B\sin x$

$$\frac{dy}{dx} = -A\sin x + B\cos x$$
$$\frac{d^2y}{dx^2} = -A\cos x - B\sin x$$
$$= -y$$
$$\frac{d^2y}{dx^2} + y = 0$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

21a

$$\frac{dy}{dx} = -\sqrt{1-y^2}$$

$$\int \frac{-1}{\sqrt{1-y^2}}\, dy = \int 1\, dx$$

$$\cos^{-1} y = x + C$$

21b Substituting $y(0) = 1$:

$$\cos^{-1} 1 = 0 + C$$

$$0 = 0 + C$$

$$C = 0$$

Therefore $\cos^{-1} y = x$

21c

$$\frac{dy}{dx} = -\sqrt{1-y^2}$$

If $y = \cos x$,

$$\text{LHS} = \frac{d}{dx}(\cos x) = -\sin x$$

$$\text{RHS} = -\sqrt{1-\cos^2 x} = -|\sin x|$$

So the two solutions differ when $\sin x$ is negative.

21d The DE holds when $\sin x$ is non-negative. The largest region containing the initial point where this is true is $0 \leq x \leq \pi$ so the solution is:

$$y = \cos x\,, 0 \leq x \leq \pi$$

22a $y = x^3$

$$\frac{dy}{dx} = 3x^2$$

© Cambridge University Press 2019

Substituting into the initial value problem $\frac{dy}{dx} = 3y^{\frac{2}{3}}$:

$\text{LHS} = 3x^2$

$\text{RHS} = 3(x^3)^{\frac{2}{3}}$

$= 3x^2$

$= \text{LHS}$

Hence $y = x^3$ is a solution of this IVP.

22b $y = 0$

$\frac{dy}{dx} = 0$

Substituting into the initial value problem $\frac{dy}{dx} = 3y^{\frac{2}{3}}$:

$\text{LHS} = 0$

$\text{RHS} = 3(0)^{\frac{2}{3}}$

$= 0$

$= \text{LHS}$

Hence $y = 0$ is a solution of this IVP.

22c $y(0) = 0$

From parts a and b above, we know the DE is satisfied for $x \neq 0$

For $x < 0, y = 0$ and $\frac{dy}{dx} = 0$

As $x \to 0^+, y$ tends to zero and $\frac{dy}{dx} = 3x^2$ also tends to zero.

Therefore at $x = 0, y = 0$ and $\frac{dy}{dx} = 0$ so the DE and initial values are satisfied here too.

© Cambridge University Press 2019

22d $y(0) = 0$

From parts a and b above, we know the DE is satisfied for $x \neq 0$

For $x > 0, y = 0$ and $\frac{dy}{dx} = 0$

As $x \to 0^-, y$ tends to zero and $\frac{dy}{dx} = 3x^2$ also tends to zero.

Therefore at $x = 0, y = 0$ and $\frac{dy}{dx} = 0$ so the DE and initial values are satisfied here too.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

Solutions to Exercise 13E

Let the integration constants be A, B, C or D.

1a i $y = ax^2 + bx$

$y' = 2ax + b$

Substituting for y' in $y' = 1 - 4x$:

$2ax + b = 1 - 4x$

Equating coefficients:

$2a = -4$ and $b = 1$

$a = -2$ and $b = 1$

Therefore $y = -2x^2 + x$

1a ii $y = e^{-x}(a\cos x + b\sin x)$

$y' = -e^{-x}(a\cos x + b\sin x) + e^{-x}(-a\sin x + b\cos x)$

$y' = e^{-x}\big(-(a\cos x + b\sin x) + (-a\sin x + b\cos x)\big)$

$y' = e^{-x}(-a\cos x - b\sin x - a\sin x + b\cos x)$

Substituting for y' in $y' = 2e^{-x}\sin x$:

$e^{-x}(-a\cos x - b\sin x - a\sin x + b\cos x) = 2e^{-x}\sin x$

Equating coefficients:

$-a - b = 2$ (1)

$-a + b = 0$ (2)

(2) + (3):

$-2a = 2$

$a = -1$

Substituting $a = -1$ into (2):

$1 + b = 0$

$b = -1$

Therefore $y = e^{-x}(-\cos x - \sin x)$ or $y = -e^{-x}(\cos x + \sin x)$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

1a iii $y = ax + b + 3e^{-x}$

$y' = a - 3e^{-x}$

Substituting for $y'$and y in $y' = x - y$:

$a - 3e^{-x} = x - (ax + b + 3e^{-x})$

$a - 3e^{-x} = x - ax - b - 3e^{-x}$

$-x + a = -ax - b$

Equating coefficients:

$-a = -1$ so $a = 1$

$-b = a$ so $b = -1$

Therefore $y = x - 1 + 3e^{-x}$

1b i $y = ax^2 + bx + c + 4e^{-2x}$

$y' = 2ax + b - 8e^{-2x}$

Substituting for $y'$and y in $y' + 2y = x^2 - 3x - 4$:

$2ax + b - 8e^{-2x} + 2(ax^2 + bx + c + 4e^{-2x}) = x^2 - 3x - 4$

$2ax + b - 8e^{-2x} + 2ax^2 + 2bx + 2c + 8e^{-2x} = x^2 - 3x - 4$

$2ax + b + 2ax^2 + 2bx + 2c = x^2 - 3x - 4$

$2ax^2 + 2ax + 2bx + b + 2c = x^2 - 3x - 4$

$2ax^2 + (2a + 2b)x + (b + 2c) = x^2 - 3x - 4$

Equating coefficients:

$2a = 1$ so $a = \frac{1}{2}$

$2a + 2b = -3$

$2\left(\frac{1}{2}\right) + 2b = -3$

$1 + 2b = -3$

$2b = -4$

$b = -2$

$b + 2c = -4$

$-2 + 2c = -4$

© Cambridge University Press 2019

$2c = -2$

$c = -1$

Therefore $y = \frac{1}{2}x^2 - 2x - 1 + 4e^{-2x}$

1b ii $y = ax^2 + bx + c - \sin 2x$

$y' = 2ax + b - 2\cos 2x$

$y'' = 2a + 4\sin 2x$

Substituting for y'', $y'$and y in $y'' + 4y = x^2 + 5x$:

$2a + 4\sin 2x + 4(ax^2 + bx + c - \sin 2x) = x^2 + 5x$

$2a + 4\sin 2x + 4ax^2 + 4bx + 4c - 4\sin 2x = x^2 + 5x$

$2a + 4ax^2 + 4bx + 4c = x^2 + 5x$

$4ax^2 + 4bx + (2a + 4c) = x^2 + 5x$

Equating coefficients:

$4a = 1$

$a = \frac{1}{4}$

$4b = 5$

$b = \frac{5}{4}$

$2a + 4c = 0$

$2\left(\frac{1}{4}\right) + 4c = 0$

$\frac{1}{2} + 4c = 0$

$4c = -\frac{1}{2}$

$c = -\frac{1}{8}$

Therefore $y = \frac{1}{4}x^2 + \frac{5}{4}x - \frac{1}{8} - \sin 2x$

or $y = \frac{1}{8}(2x^2 + 10x - 1) - \sin 2x$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

1c $y = 5e^{\lambda x}$

$y' = 5\lambda e^{\lambda x}$

$y'' = 5\lambda^2 e^{\lambda x}$

Substituting for y'', $y'$and y in $y'' + 5y' + 6y = 0$:

$5\lambda^2 e^{\lambda x} + 5(5\lambda e^{\lambda x}) + 6\left(5e^{\lambda x}\right) = 0$

$5\lambda^2 e^{\lambda x} + 25\lambda e^{\lambda x} + 30e^{\lambda x} = 0$

$5e^{\lambda x}(\lambda^2 + 5\lambda + 6) = 0$

$5e^{\lambda x} = 0$ (no solution)

or $\lambda^2 + 5\lambda + 6 = 0$

$(\lambda + 3)(\lambda + 2) = 0$

$\lambda = -3, -2$

Therefore $y = 5e^{-3x}$ or $5e^{-2x}$

2a

$\frac{dR}{dt} = kR$

$\frac{1}{R}\, dR = k\, dt$

$\int \frac{1}{R}\, dR = \int k\, dt$

$\ln|R| = kt + C$

$R = e^{kt+C}$ since R must be positive

$R = e^C e^{kt}$

$R = Ae^{kt}$ where $A = e^C$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

2b $R = Ae^{kt}$

When $t = 0, R = 100$, so:

$100 = Ae^{k\times 0}$

$A = 100$

2c $R = 100e^{kt}$

When $t = 4, R = 20$, so:

$20 = 100e^{4k}$

$\frac{1}{5} = e^{4k}$

$\ln\left(\frac{1}{5}\right) = 4k$

$k = \frac{1}{4}\ln(5^{-1})$

$k = -\frac{1}{4}\ln 5$

2d $R = 100e^{kt}$ where $k = -\frac{1}{4}\ln 5$

When $t = 12$,

$R = 100e^{\left(-\frac{1}{4}\ln 5\right)\times 12}$

$= 100e^{(-3\ln 5)}$

$= 100e^{\ln 5^{-3}}$

$= 100 \times 5^{-3}$

$= \frac{100}{125}$

$= \frac{4}{5}$

Amount present after 12 days is $\frac{4}{5}$ gram.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

2e $R = 100e^{\left(-\frac{1}{4}\ln 5\right)t}$

$R = 100e^{\left(-\frac{1}{4}t\ln 5\right)}$

$R = 100e^{\left(\ln 5^{-\frac{1}{4}t}\right)}$

$R = 100 \times 5^{-\frac{1}{4}t}$

$R = 100 \times \left(\frac{1}{5}\right)^{\frac{1}{4}t}$

When $t = 12$,

$R = 100 \times \left(\frac{1}{5}\right)^{3} = 100 \times \frac{1}{125} = \frac{4}{5}$

3a

$\frac{dH}{dt} = k(H - 25)$

$\frac{1}{H - 25}\, dH = k\, dt$

$\int \frac{1}{H - 25}\, dR = \int k\, dt$

$\ln|H - 25| = kt + C$

$|H - 25| = e^{kt+C}$

$|H - 25| = e^{C}e^{kt}$

$H - 25 = e^{C}e^{kt}\ or - e^{C}e^{kt}$

$H - 25 = Ae^{kt}$ where $A = e^{C}$ or $-e^{C}$

$H = 25 + Ae^{kt}$

3b When $t = 0, H = 5$, so:

$5 = 25 + Ae^{k(0)}$

$5 = 25 + A$

$-20 = A$

Therefore $H = 25 - 20e^{kt}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

3c When $t = 10, H = 15$, so:

$15 = 25 - 20e^{10k}$

$20e^{10k} = 10$

$e^{10k} = \frac{10}{20}$

$10k = \ln\frac{1}{2}$

$k = \frac{1}{10}\ln(2^{-1})$

$k = -\frac{1}{10}\ln 2$

3d When $H = 24$,

$24 = 25 - 20e^{\left(-\frac{1}{10}\ln 2\right)t}$

$20e^{\left(-\frac{1}{10}\ln 2\right)t} = 1$

$2^{-\frac{1}{10}t} = \frac{1}{20}$

$t \div 43$ mins

4a

$\frac{dV}{dt} = k\pi r^2$, for some constant k

4b

$\frac{dr}{dt} = \frac{dr}{dV} \times \frac{dV}{dt}$

$\frac{dr}{dt} = \frac{dr}{dV} \times k\pi r^2$

$h = 3r$ (from initial proportions) so

$V = \frac{1}{3}\pi r^2 \times 3r = \pi r^3$

© Cambridge University Press 2019

Therefore $\frac{dV}{dr} = 3\pi r^2$

$\frac{dr}{dt} = \frac{1}{3\pi r^2} \times k\pi r^2$

$\frac{dr}{dt} = \frac{1}{3}k$

4c

$\frac{dr}{dt} = \frac{1}{3}k$

$\int 3\,dr = \int k\,dt$

$3r = kt + C$

$r = \frac{1}{3}kt + D$ where $D = \frac{1}{3}C$

Substituting $r(0) = 4$:

$4 = \frac{1}{3}k(0) + D$

$D = 4$

Therefore $r = \frac{1}{3}kt + 4$

4d Substituting $r(6) = 3.5$:

$3.5 = \frac{1}{3}k(6) + 4$

$-0.5 = 2k$

$2k = -\frac{1}{2}$

$k = -\frac{1}{4}$

Therefore $r = \frac{1}{3}\left(-\frac{1}{4}\right)t + 4$

$r = 4 - \frac{1}{12}t$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4e $V = \pi r^3$

$V = \pi\left(4 - \frac{1}{12}t\right)^3$

Since r and V cannot be negative, $4 - \frac{1}{12}t$ must also be non-negative therefore $0 \leq t \leq 48$.

5a $h(0) = 400$

The height decreases so $\frac{dh}{dt} < 0$.

Since $\sqrt{h} \geq 0$, the constant k must be negative.

5b

$\frac{dh}{dt} = k\sqrt{h}$

$\int \frac{1}{\sqrt{h}}\, dh = \int k\, dt$

$\int h^{-\frac{1}{2}}\, dh = \int k\, dt$

$2h^{\frac{1}{2}} = kt + C$

$h^{\frac{1}{2}} = \frac{1}{2}(kt + C)$

$h = \frac{1}{4}(kt + C)^2$

Substituting $h(0) = 400$:

$400 = \frac{1}{4}(k(0) + C)^2$

$1600 = C^2$

$C = 40$

Therefore $h = \frac{1}{4}(kt + 40)^2$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

5c Substituting $h(20) = 100$:

$100 = \frac{1}{4}(k \times 20 + 40)^2$

$400 = (20k + 40)^2$

$20 = 20k + 40$

$20k = -20$

$k = -1$

5d

$h = \frac{1}{4}(-t + 40)^2$

When $h = 0$,

$0 = \frac{1}{4}(-t + 40)^2$

$0 = (-t + 40)^2$

$0 = -t + 40$

$t = 40$ mins

5e No, $h(t)$ increases for $t > 40$, which is impossible.

6a

$$\text{Gradient} = \frac{y - 0}{x - 0} = \frac{y}{x}$$

6b

$$\frac{dy}{dx} = \frac{y}{x}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

6c

$$\int \frac{1}{y}\,dy = \int \frac{1}{x}\,dx$$

$$\ln|y| = \ln|x| + B$$

$$\ln|y| - \ln|x| = B$$

$$\ln\left|\frac{y}{x}\right| = B$$

$$\left|\frac{y}{x}\right| = e^B$$

$$\frac{y}{x} = e^B \text{ or } -e^B$$

$$\frac{y}{x} = C \text{ where } C = e^B \text{ or } -e^B$$

$$y = Cx$$

6d $x = 0$ because then y' undefined.

7a

$$\text{Gradient} = \frac{y-0}{x-0} = \frac{y}{x}$$

7b Normal has gradient of $-\frac{x}{y}$.

$$\frac{dy}{dx} = -\frac{x}{y}$$

7c

$$\int y\ dy = -\int x\ dx$$

$$\frac{y^2}{2} = -\frac{x^2}{2} + C$$

$$\frac{x^2}{2} + \frac{y^2}{2} = C$$

$$x^2 + y^2 = 2C$$

© Cambridge University Press 2019

$x^2 + y^2 = r^2$ where $r^2 = 2C$ and r is the radius of the circle

8a

$$\text{Gradient} = \frac{y - 0}{x - (x - 1)} = y$$

8b

$$\frac{dy}{dx} = y$$

8c

$$\int \frac{1}{y} dy = \int 1 \ dx$$

$$\ln|y| = x + C$$

$$|y| = e^{x+C}$$

$y = e^C e^x$ or $-e^C e^x$

$y = Ae^x$ where $A = e^C$ or $-e^C$

8d Substituting $(0, 1)$:

$1 = Ae^0$

$A = 1$

Therefore $y = e^x$

9a

$$\frac{dP}{dh} = kP$$

$$\int \frac{1}{P} \ dP = \int k \ dh$$

$$ln\,|P| = kh + C$$

$P = e^{kh+C}$ (from the context of the problem, we can ignore the negative case)

$P = e^C e^{kh}$

© Cambridge University Press 2019

$P = Ae^{kh}$ where $A = e^C$

Substituting $(0, P_0)$:

$P_0 = Ae^0$

$A = P_0$

Therefore $P = P_0e^{kh}$

9b Substituting $(10\,000, 40)$:

$40 = P_0e^{10\,000k}$ (1)

Substituting $(6000, 80)$:

$80 = P_0e^{6000k}$ (2)

$(2) \div (1)$:

$$\frac{80}{40} = \frac{e^{6000k}}{e^{10\,000k}}$$

$2 = e^{6000k - 10\,000k}$

$2 = e^{-4000k}$

$\ln 2 = -4000k$

$$k = -\frac{\ln 2}{4000}$$

9c Substituting $k = -\frac{\ln 2}{4000}$ into (1):

$40 = P_0\, e^{10\,000\left(-\frac{\ln 2}{4000}\right)}$

$40 = P_0\, e^{\left(-\frac{5}{2}\ln 2\right)}$

$40 = P_0\, 2^{\left(-\frac{5}{2}\right)}$

$$\frac{40}{2^{\left(-\frac{5}{2}\right)}} = P_0$$

$160\sqrt{2} = P_0$

Therefore $P = 160\sqrt{2}\, e^{\left(-\frac{\ln 2}{4000}\right)h}$

When $h = 0$,

© Cambridge University Press 2019

$P = 160\sqrt{2} \times e^{\left(-\frac{\ln 2}{4000}\right)\times 0}$

$P = 160\sqrt{2}$

$P \div 226 \text{ kPa}$

10a $A = (2x, 0), B = (0, 2y)$

10b

$$\frac{dy}{dx} = \frac{2y - 0}{0 - 2x} = -\frac{y}{x}$$

10c

$$\int \frac{1}{y}\, dy = -\int \frac{1}{x}\, dx$$

$$\ln|y| = -\ln|x| + B$$

$$\ln|y| + \ln|x| = B$$

$$\ln|xy| = B$$

$$|xy| = e^B$$

$xy = e^B$ or $-e^B$

$xy = C$ where $C = e^B$ or $-e^B$

or $y = \frac{C}{x}$

11a

$$\frac{dC}{dt} = k(S - C)$$

$$\int \frac{1}{S - C}\, dC = \int k\ dt$$

$$\int \frac{1}{C - S}\, dC = -\int k\ dt$$

$$ln\,|C - S| = -kt + B$$

$$|C - S| = e^{-kt+B}$$

© Cambridge University Press 2019

$|C - S| = e^B e^{-kt}$

$C - S = e^B e^{-kt}$ or $e^B e^{-kt}$

$C - S = Ae^{-kt}$ where $A = e^{-B}$ or $-e^{-B}$

$C = S + Ae^{-kt}$

11b Substituting $(0, C_0)$:

$C_0 = S + Ae^{-k(0)}$

$C_0 = S + A$

$C_0 - S = A$

Therefore $C = S + (C_0 - S)e^{-kt}$

12a

$$\text{RHS} = \frac{1}{N} + \frac{1}{1000 - N}$$

$$= \frac{1000 - N + N}{N(P - N)}$$

$$= \frac{1000}{N(1000 - N)}$$

$$= \text{LHS}$$

Therefore $\dfrac{1000}{N(1000 - N)} = \dfrac{1}{N} + \dfrac{1}{1000 - N}$

Alternatively, let

$$\frac{1000}{N(1000 - N)} = \frac{A}{N} + \frac{B}{1000 - N}$$

$$= \frac{A(1000 - N) + B(N)}{N(1000 - N)}$$

$$= \frac{1000A - AN + BN}{(y - 1)(y - 3)}$$

$$= \frac{N(-A + B) + 1000A}{(y - 1)(y - 3)}$$

Equating coefficients:

© Cambridge University Press 2019

$1000A = 1000$

$A = 1$

$-A + B = 0$

Since $A = 1$,

$-1 + B = 0$

$B = 1$

Therefore $\dfrac{1000}{N(1000 - N)} = \dfrac{1}{N} + \dfrac{1}{1000 - N}$

12b

$\dfrac{dN}{dt} = kN(1000 - N)$

$\displaystyle\int \frac{1}{N(1000 - N)}\, dN = \int k\, dt$

$\dfrac{1}{1000}\displaystyle\int \frac{1000}{N(1000 - N)}\, dN = \int k\, dt$

$\displaystyle\int \left(\frac{1}{N} + \frac{1}{1000 - N}\right) dN = \int 1000k\, dt$

$\ln|N| - \ln|1000 - N| = 1000kt + C$

$\ln\left|\dfrac{N}{1000 - N}\right| = 1000kt + C$

$\left|\dfrac{N}{1000 - N}\right| = e^{1000kt + C}$

$\dfrac{N}{1000 - N} = e^C e^{1000kt}$ or $-e^C e^{1000kt}$

$\dfrac{N}{1000 - N} = Ae^{1000kt}$ where $A = e^C$ or $-e^C$

$N = Ae^{1000kt}(1000 - N)$

$N = 1000Ae^{1000kt} - NAe^{1000kt}$

$N + NAe^{1000kt} = 1000Ae^{1000kt}$

$N(1 + Ae^{1000kt}) = 1000Ae^{1000kt}$

$N = \dfrac{1000Ae^{1000kt}}{1 + Ae^{1000kt}}$

© Cambridge University Press 2019

$$N = \frac{1000Ae^{1000kt}}{1 + Ae^{1000kt}} \times \frac{A^{-1}e^{-1000kt}}{A^{-1}e^{-1000kt}}$$

$$N = \frac{1000}{A^{-1}e^{-1000kt} + 1}$$

$$N = \frac{1000}{Be^{-1000kt} + 1} \text{ where } B = A^{-1}$$

$$N = \frac{1000}{1 + Be^{-1000kt}}$$

12c Substituting $N(0) = 40$:

$$40 = \frac{1000}{1 + Be^{-1000k(0)}}$$

$$40 = \frac{1000}{1 + B}$$

$$40(1 + B) = 1000$$

$$40 + 40B = 1000$$

$$B = \frac{1000 - 40}{40}$$

$$B = 24$$

Therefore $N = \frac{1000}{1 + 24e^{-1000kt}}$

12d Substituting $N(1) = 80$:

$$80 = \frac{1000}{1 + 24e^{-1000k(1)}}$$

$$80 + 1920e^{-1000k} = 1000$$

$$1920e^{-1000k} = 920$$

$$e^{-1000k} = \frac{920}{1920}$$

$$-1000k = \ln\left(\frac{23}{48}\right)$$

$$k = -\frac{1}{1000}\ln\left(\frac{23}{48}\right)$$

$$k \div 7.357 \times 10^{-4}$$

© Cambridge University Press 2019

12e When $t = 5$,

$$N = \frac{1000}{1 + 24e^{-1000k(5)}}$$

$$\div 622.57$$

Population will be 623 after 5 years.

13a

$$\text{RHS} = \frac{1}{N} + \frac{1}{P - N}$$

$$= \frac{P - N + N}{N(P - N)}$$

$$= \frac{P}{N(P - N)}$$

$$= \text{LHS}$$

Therefore $\frac{P}{N(P - N)} = \frac{1}{N} + \frac{1}{P - N}$

Alternatively, let

$$\frac{P}{N(P - N)} = \frac{A}{N} + \frac{B}{P - N}$$

$$= \frac{A(P - N) + B(N)}{N(P - N)}$$

$$= \frac{AP - AN + BN}{N(P - N)}$$

$$= \frac{N(-A + B) + AP}{N(P - N)}$$

© Cambridge University Press 2019

Equating coefficients:

$AP = P$

$A = 1$

$-A + B = 0$

Since $A = 1$,

$-1 + B = 0$

$B = 1$

Therefore $\frac{P}{N(P-N)} = \frac{1}{N} + \frac{1}{P-N}$

13b (Same as Q12b with $P = 1000$.)

$\frac{dN}{dt} = kN(P-N)$

$\int \frac{1}{N(P-N)}\, dN = \int k\, dt$

$\frac{1}{P}\int \frac{P}{N(P-N)}\, dN = \int k\, dt$

$\int \left(\frac{1}{N} + \frac{1}{P-N}\right) dN = \int kP\, dt$

$\ln|N| - \ln|P-N| = kPt + C$

$\ln\left|\frac{N}{P-N}\right| = kPt + C$

$\left|\frac{N}{P-N}\right| = e^{kPt+C}$

$\frac{N}{P-N} = e^C e^{kPt}$ or $-e^C e^{kPt}$

$\frac{N}{P-N} = Ae^{kPt}$ where $A = e^C$ or $-e^C$

$N = Ae^{kPt}(P-N)$

$N = PAe^{kPt} - NAe^{kPt}$

$N + NAe^{kPt} = PAe^{kPt}$

$N(1 + Ae^{kPt}) = PAe^{kPt}$

© Cambridge University Press 2019

$$N = \frac{PAe^{kPt}}{1 + Ae^{kPt}}$$

$$N = \frac{PAe^{kPt}}{1 + Ae^{kPt}} \times \frac{A^{-1}e^{-kPt}}{A^{-1}e^{-kPt}}$$

$$N = \frac{P}{A^{-1}e^{-kPt} + 1}$$

$$N = \frac{P}{Be^{-kPt} + 1} \text{ where } B = A^{-1}$$

$$N = \frac{P}{1 + Be^{-kPt}}$$

13c Substituting $N(0) = 23.2$ and $P = 187.5$:

$$23.2 = \frac{187.5}{1 + Be^{-187.5k(0)}}$$

$$23.2 = \frac{187.5}{1 + B}$$

$$23.2(1 + B) = 187.5$$

$$23.2 + 23.2B = 187.5$$

$$B = \frac{187.5 - 23.2}{23.2}$$

$$B = \frac{164.3}{23.2}$$

Therefore $N = \dfrac{187.5}{1 + \frac{164.3}{23.2}e^{-187.5kt}}$

$$N = \frac{187.5}{1 + \frac{164.3}{23.2}e^{-187.5kt}} \times \frac{23.2}{23.2}$$

$$N = \frac{187.5 \times 23.2}{23.21 + 164.3e^{-187.5kt}}$$

© Cambridge University Press 2019

13d Substituting $N(40, 63.0)$:

$$63.0 = \frac{187.5 \times 23.2}{23.21 + 164.3e^{-187.5(40)k}}$$

$$63.0(23.21 + 164.3e^{-7500k}) = 187.5 \times 23.2$$

$$23.21 + 164.3e^{-7500k} = \frac{4350}{63.0}$$

$$164.3e^{-7500k} = \frac{4350}{63.0} - 23.21$$

$$e^{-7500k} = \frac{4350 - 23.21 \times 63.0}{63.0 \times 164.3}$$

$$-7500k = \ln\left(\frac{4350 - 1462.23}{10\,350.9}\right)$$

$$k = -\frac{1}{7500}\ln\left(\frac{2887.77}{10\,350.9}\right)$$

$$k \doteqdot 1.702 \times 10^{-4}$$

13e When $t = 80$ and $k = 1.702 \times 10^{-4}$,

$$N = \frac{187.5 \times 23.2}{23.21 + 164.3e^{-187.5(1.702 \times 10^{-4})(80)}}$$

$$N = \frac{4350}{23.21 + 164.3e^{-2.553}}$$

$$N = 120.832 \ldots$$

Predicted population is about 120.9 million.

13f The mathematical model needs to be revised. Clearly the carrying capacity is much larger than the figure of 187.5 estimated in 1850.

Using the model, when $t = 168$ and $k = 1.702 \times 10^{-4}$,

$$N = \frac{187.5 \times 23.2}{23.21 + 164.3e^{-187.5(1.702 \times 10^{-4})(168)}}$$

$$N = \frac{4350}{23.21 + 164.3e^{-5.3613}}$$

$$N = 181.401 \ldots$$

Predicted population from the model is about 181.4 million.

© Cambridge University Press 2019

14a Using the results from Question 13a and 13b,

$$N = \frac{P}{1 + Be^{-kPt}}$$

Alternatively, the full working is shown below.

From Q13a, $\frac{P}{N(P-N)} = \frac{1}{N} + \frac{1}{P-N}$

$$\frac{dN}{dt} = kN(P-N)$$

$$\int \frac{1}{N(P-N)}\, dN = \int k\, dt$$

$$\frac{1}{P}\int \frac{P}{N(P-N)}\, dN = \int k\, dt$$

$$\int \left(\frac{1}{N} + \frac{1}{P-N}\right) dN = \int kP\, dt$$

$$\ln|N| - \ln|P-N| = kPt + C$$

$$\ln\left|\frac{N}{P-N}\right| = kPt + C$$

$$\left|\frac{N}{P-N}\right| = e^{kPt+C}$$

$$\frac{N}{P-N} = e^C e^{kPt} \text{ or } -e^C e^{kPt}$$

$$\frac{N}{P-N} = Ae^{kPt} \text{ where } A = e^C \text{ or } -e^C$$

$$N = Ae^{kPt}(P-N)$$

$$N = PAe^{kPt} - NAe^{kPt}$$

$$N + NAe^{kPt} = PAe^{kPt}$$

$$N(1 + Ae^{kPt}) = PAe^{kPt}$$

$$N = \frac{PAe^{kPt}}{1 + Ae^{kPt}}$$

$$N = \frac{PAe^{kPt}}{1 + Ae^{kPt}} \times \frac{A^{-1}e^{-kPt}}{A^{-1}e^{-kPt}}$$

$$N = \frac{P}{A^{-1}e^{-kPt} + 1}$$

© Cambridge University Press 2019

$$N = \frac{P}{Be^{-kPt} + 1} \text{ where } B = A^{-1}$$

$$N = \frac{P}{1 + Be^{-kPt}}$$

14b Substituting $N(0) = N_0$:

$$N_0 = \frac{P}{1 + Be^{-kP(0)}}$$

$$N_0 = \frac{P}{1 + B}$$

$$N_0(1 + B) = P$$

$$N_0 + BN_0 = P$$

$$BN_0 = P - N_0$$

$$B = \frac{P - N_0}{N_0}$$

$$\text{Therefore } N = \frac{P}{1 + \left(\frac{P - N_0}{N_0}\right)e^{-kPt}}$$

$$N = \frac{P}{1 + \left(\frac{P - N_0}{N_0}\right)e^{-kPt}} \times \frac{N_0}{N_0}$$

$$N = \frac{N_0 P}{N_0 + (P - N_0)e^{-kPt}}$$

14c Substituting $N(t_1) = N_1$:

$$N_1 = \frac{N_0 P}{N_0 + (P - N_0)e^{-kPt_1}}$$

$$\frac{N_1}{N_0 P} = \frac{1}{N_0 + (P - N_0)e^{-kPt_1}}$$

$$\frac{N_0 P}{N_1} = \frac{N_0 + (P - N_0)e^{-kPt_1}}{1}$$

$$\frac{N_0 P}{N_1} - N_0 = (P - N_0)e^{-kPt_1}$$

© Cambridge University Press 2019

$$\frac{N_0P - N_0N_1}{N_1} = (P - N_0)e^{-kPt_1}$$

$$\frac{N_0P - N_0N_1}{N_1(P - N_0)} = e^{-kPt_1}$$

$$\frac{N_0(P - N_1)}{N_1(P - N_0)} = e^{-kPt_1}$$

$$\frac{N_1(P - N_0)}{N_0(P - N_1)} = e^{kPt_1}$$

$$kPt_1 = \ln\left(\frac{N_1(P - N_0)}{N_0(P - N_1)}\right)$$

$$k = \frac{1}{t_1P}\ln\left(\frac{N_1(P - N_0)}{N_0(P - N_1)}\right)$$

14d

$$N = \frac{N_0P}{N_0 + (P - N_0)e^{-Pk(t)}} \text{ and } k = \frac{1}{t_1P}\ln\left(\frac{N_1(P - N_0)}{N_0(P - N_1)}\right)$$

By the same logic (as in part c),

$$k = \frac{1}{t_2P}\ln\left(\frac{N_2(P - N_0)}{N_0(P - N_{2)}}\right)$$

Substituting $t_1 = 1$ and $t_2 = 2$:

$$\frac{1}{P}\ln\left(\frac{N_1(P - N_0)}{N_0(P - N_1)}\right) = k = \frac{1}{2P}\ln\left(\frac{N_2(P - N_0)}{N_0(P - N_2)}\right)$$

$$2\ln\left(\frac{N_1(P - N_0)}{N_0(P - N_1)}\right) = \ln\left(\frac{N_2(P - N_0)}{N_0(P - N_2)}\right)$$

$$\ln\left(\frac{N_1(P - N_0)}{N_0(P - N_1)}\right)^2 = \ln\left(\frac{N_2(P - N_0)}{N_0(P - N_2)}\right)$$

$$\left(\frac{N_1(P - N_0)}{N_0(P - N_1)}\right)^2 = \frac{N_2(P - N_0)}{N_0(P - N_2)}$$

$$N_0(P - N_2)\big(N_1(P - N_0)\big)^2 = N_2(P - N_0)\big(N_0(P - N_1)\big)^2$$

$$(P - N_2)(P - N_0)(N_1)^2 = N_0N_2(P - N_1)^2$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

15a It represents the 200 fish harvested each year.

15b

$$\frac{dy}{dt} = -2 + \frac{1}{24}y(16 - y)$$

$$\frac{dy}{dt} = \frac{1}{24}(-48 + 16y - y^2)$$

$$\frac{dy}{dt} = -\frac{1}{24}(y^2 - 16y + 48)$$

$$\frac{dy}{dt} = -\frac{1}{24}(y - 4)(y - 12)$$

Initial condition: $y(0) = 5 - 2 = 3$ (where y is measured in hundreds)

15c RHS

$$= \frac{3}{y - 12} - \frac{3}{y - 4}$$

$$= \frac{3(y - 4) - 3(y - 12)}{(y - 12)(y - 4)}$$

$$= \frac{3y - 12 - 3y + 36}{(y - 12)(y - 4)}$$

$$= \frac{24}{(y - 12)(y - 4)}$$

$$= \text{LHS}$$

15d

$$\frac{dy}{dt} = -\frac{1}{24}(y - 4)(y - 12)$$

$$\int \frac{24}{(y - 12)(y - 4)}\,dy = -\int 1\,dt$$

$$\int \left(\frac{3}{y - 12} - \frac{3}{y - 4}\right)dy = -\int 1\,dt$$

$$\int \left(\frac{1}{y - 12} - \frac{1}{y - 4}\right)dy = -\frac{1}{3}\int 1\,dt$$

© Cambridge University Press 2019

$$\ln|y-12| - \ln|y-4| = -\frac{1}{3}t + C$$

$$\ln\left|\frac{y-12}{y-4}\right| = -\frac{1}{3}t + C$$

$$\left|\frac{y-12}{y-4}\right| = e^{-\frac{1}{3}t+C}$$

$$\frac{y-12}{y-4} = e^{C}e^{-\frac{1}{3}t} \text{ or } e^{C}e^{-\frac{1}{3}t}$$

$$\frac{y-12}{y-4} = Ae^{-\frac{1}{3}t} \text{ where } A = e^{C} \text{ or } -e^{C}$$

$$y-12 = (y-4)Ae^{-\frac{1}{3}t}$$

$$y-12 = yAe^{-\frac{1}{3}t} - 4Ae^{-\frac{1}{3}t}$$

$$y\left(1 - Ae^{-\frac{1}{3}t}\right) = 12 - 4Ae^{-\frac{1}{3}t}$$

$$y = \frac{12 - 4Ae^{-\frac{1}{3}t}}{1 - Ae^{-\frac{1}{3}t}}$$

Substituting $y(0) = 3$:

$$3 = \frac{12 - 4Ae^{0}}{1 - Ae^{0}}$$

$$3 = \frac{12 - 4A}{1 - A}$$

$$3 - 3A = 12 - 4A$$

$$A = 9$$

Therefore $y = \dfrac{12 - 36e^{-\frac{1}{3}t}}{1 - 9e^{-\frac{1}{3}t}}$

$$y = \frac{12\left(1 - 3e^{-\frac{1}{3}t}\right)}{1 - 9e^{-\frac{1}{3}t}}$$

© Cambridge University Press 2019

15e Substituting $y = 0$:

$$0 = \frac{12\left(1 - 3e^{-\frac{1}{3}t}\right)}{1 - 9e^{-\frac{1}{3}t}}$$

$$1 - 3e^{-\frac{1}{3}t} = 0$$

$$e^{-\frac{1}{3}t} = \frac{1}{3}$$

$$t = -3\ln\frac{1}{3}$$

$$= 3\ln 3$$

$$= 3.295\,8\ldots$$

$$\div 3.3$$

Fish will die out in approximately 3.3 years.

15f i $y(0) = 5$

15f ii

$$y = \frac{4\left(3 + 7e^{-\frac{1}{3}t}\right)}{1 + 7e^{-\frac{1}{3}t}}$$

As $t \to \infty$, $e^{-\frac{1}{3}t} \to 0$

Therefore y approaches:

$$\frac{4(3 + 7(0))}{1 + 7(0)}$$

$$= \frac{12}{1}$$

$$= 12$$

So skipping the harvest for the first year allows the fish population to stabilise at 1200. This suggests the harvest should be stopped for one year to save the species.

© Cambridge University Press 2019

16a i Let $u = log_e x$

$$\frac{du}{dx} = \frac{1}{x}$$

$$x\,du = dx$$

$$I = \int \frac{dx}{x \log_e x}$$

$$I = \int \frac{x\,du}{xu}$$

$$I = \int \frac{du}{u}$$

16a ii $I = log_e|u| + C$

$= log_e|log_e x| + C$

16b

$$\frac{dN}{dt} = kN \log_e N$$

$$\int \frac{1}{N log_e N}\,dN = \int k\,dt$$

$log_e|log_e N| = kt + C$ (using part a)

$|log_e N| = e^{kt+C}$

$log_e N = e^C e^{kt}$ or $-e^C e^{kt}$

$log_e N = Ae^{kt}$ where $A = e^C$ or $-e^C$

$N = e^{Ae^{kt}}$

17 $M_0 = 200$

$$\frac{dM}{dt} = -\text{outflow} - \text{decay}$$

$$\frac{dM}{dt} = -\frac{5}{100}M - \frac{10}{100}M$$

$$= -\frac{15}{100}M$$

© Cambridge University Press 2019

$= -\frac{3}{20}M$

$\int \frac{1}{M}\, dM = -\frac{3}{20}\int 1\, dt$

$\ln M = -\frac{3}{20}t + C$ (we can omit absolute value signs since M cannot be negative)

$M = e^{-\frac{3}{20}t+C}$

$M = e^{-\frac{3}{20}}e^{C}$

$M = Ae^{-\frac{3}{20}t}$ where $A = e^{C}$

Substituting $M(0) = 200$:

$200 = Ae^{0}$

$200 = A$

Therefore $M = 200e^{-\frac{3}{20}t}$

18a

$\frac{dh}{dt} = k(h - 20)$

$\int \frac{1}{h-20}\, dh = \int k\, dt$

$\ln|h - 20| = kt + C$

$|h - 20| = e^{kt+C}$

$|h - 20| = e^{kt}e^{C}$

$h - 20 = Ae^{kt}$ where $A = e^{C}$ or $-e^{C}$

$h = 20 + Ae^{kt}$

Substituting $h(0) = 100$:

$100 = 20 + Ae^{k(0)}$

$A = 80$

Therefore $h = 20 + 80e^{kt}$

$h = 20(1 + 4e^{kt})$

© Cambridge University Press 2019

Substituting $h(10) = 80$:

$80 = 20(1 + 4e^{10k})$

$1 + 4e^{10k} = 4$

$4e^{10k} = 3$

$e^{10k} = \frac{3}{4}$

$10k = \ln\frac{3}{4}$

$k = \frac{1}{10}\ln\frac{3}{4}$

18b i

$\frac{dH}{dt} = k(H - 20 + t)$

$H(0) = 80$

18b ii $y = H - 20 + t$

$\frac{dy}{dt} = \frac{dH}{dt} + 1$

$\frac{dy}{dt} = k(H - 20 + t) + 1$

$\frac{dy}{dt} = ky + 1$

$y(0) = 80 - 20 = 60$

18b iii

$\frac{dy}{dt} = ky + 1$

$\int \frac{1}{ky + 1}\, dy = \int 1\, dt$

$\frac{1}{k}\int \frac{1}{y + \frac{1}{k}} dy = \int 1\, dt$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

$$\int \frac{1}{y+\frac{1}{k}}dy = k\int 1\,dt$$

$$\ln\left|y+\frac{1}{k}\right| = kt + C$$

$$\left|y+\frac{1}{k}\right| = e^{kt+C}$$

$$\left|y+\frac{1}{k}\right| = e^{C}e^{kt}$$

$y+\frac{1}{k} = Ae^{kt}$ where $A = e^{C}$ or $-e^{C}$

$$y = Ae^{kt} - \frac{1}{k}$$

Substituting $y(0) = 60$:

$$60 = Ae^{0t} - \frac{1}{k}$$

$$A = 60 + \frac{1}{k}$$

Therefore $y = \left(60+\frac{1}{k}\right)e^{kt} - \frac{1}{k}$

Since $y = H - 20 + t$, $\quad H = y + 20 - t$

$$H = \left(60+\frac{1}{k}\right)e^{kt} - \frac{1}{k} + 20 - t$$

or $H(t) = 20 - \frac{1}{k} - t + \left(60+\frac{1}{k}\right)e^{kt}$

19a

$$\frac{dN}{dt} = kN(P-N)$$

$$\frac{d(Py)}{dt} = kPy(P-Py)$$

$$P\frac{dy}{dt} = kP^2y(1-y)$$

$$\frac{dy}{dt} = kPy(1-y)$$

© Cambridge University Press 2019

$$\frac{dy}{dt} = ry(1-y) \qquad (\text{since } r = kP)$$

$$y(0) = y_0 = \frac{N_0}{P}$$

19b $x = rt$

$$\frac{dx}{dt} = r$$

Using the chain rule:

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$$

$$= ry(1-y) \times \frac{1}{r}$$

$$= y(1-y)$$

Initial condition: $y(0) = y_0$

19c

$$v = \frac{1}{y}$$

$$\frac{dv}{dy} = -\frac{1}{y^2}$$

Using the chain rule:

$$\frac{dv}{dx} = \frac{dv}{dy} \times \frac{dy}{dx}$$

$$= -\frac{1}{y^2} \times y(1-y)$$

$$= -\frac{1}{y}(1-y)$$

$$= -v\left(1 - \frac{1}{v}\right)$$

$$= 1 - v$$

or $v' = 1 - v$

© Cambridge University Press 2019

19d $v' = 1 - v$

$$\int \frac{1}{1-v}\, dv = \int 1\, dx$$

$$\int \frac{1}{v-1}\, dv = -\int 1\, dx$$

$$\ln|v-1| = -x + C$$

$$|v-1| = e^{-x+C}$$

$v - 1 = e^C e^{-x}$ or $-e^C e^{-x}$

$v - 1 = Ae^{-x}$ where $A = e^C$ or $-e^C$

$$v = 1 + Ae^{-x}$$

Since $v = \frac{1}{y}$ or $y = \frac{1}{v}$

$$y = \frac{1}{1 + Ae^{-x}}$$

19e Substituting $y(0) = y_0 = \frac{N_0}{P}$:

$$\frac{N_0}{P} = \frac{1}{1 + Ae^{-0}}$$

$$1 + A = \frac{P}{N_0}$$

$$A = \frac{P}{N_0} - 1$$

Therefore $y = \dfrac{1}{1 + \left(\frac{P}{N_0} - 1\right)e^{-x}}$

$$y = \frac{1}{1 + \left(\frac{P}{N_0} - 1\right)e^{-x}} \times \frac{N_0}{N_0}$$

$$y = \frac{N_0}{N_0 + (P - N_0)e^{-x}}$$

© Cambridge University Press 2019

19f Substituting $x = rt = kPt$:


$$y = \frac{N_0}{N_0 + (P - N_0)e^{-kPt}}$$

$$N = Py$$

$$= P \times \frac{N_0}{N_0 + (P - N_0)e^{-kPt}}$$

$$= \frac{N_0 P}{N_0 + (P - N_0)e^{-kPt}}$$

20a The curves all have an asymptote $y = 0$ on the left and an asymptote $y = 1$ on the right, an the curves become steeper at $\left(0, \frac{1}{2}\right)$ as the value of r increases.




Chapter 13 worked solutions – Differential equations

20b i As $r \to \infty, -rx \to -\infty$ so $e^{-rx} \to 0$ and $y \to 1$

20b ii As $r \to \infty, -rx = 0$ so $e^{-rx} = 1$ and $y = \frac{1}{2}$

20b iii As $r \to \infty, -rx \to \infty$ so $e^{-rx} \to \infty$ and $y \to 0$

20c



© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

Solutions to Chapter review

Let the integration constants be A, B, C or D.

1a 1st-order, linear, one arbitrary constant

1b 2nd-order, two arbitrary constants

1c 3rd-order, three arbitrary constants

2a $xy' = y(1 - x^2)$

$y' = \frac{y}{x}(1 - x^2)$

2b

y \ x	-2	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2
2	3	$\frac{4}{3}$	0	-3	*	3	0	$-\frac{4}{3}$	-3
$\frac{3}{2}$	$\frac{9}{4}$	$\frac{5}{4}$	0	$-\frac{5}{4}$	*	$\frac{5}{4}$	0	$-\frac{5}{4}$	$-\frac{9}{4}$
1	$\frac{3}{2}$	$\frac{5}{6}$	0	$-\frac{3}{2}$	*	$\frac{3}{2}$	0	$-\frac{5}{6}$	$-\frac{3}{2}$
$\frac{1}{2}$	$\frac{3}{4}$	$\frac{5}{12}$	0	$-\frac{5}{12}$	*	$\frac{5}{12}$	0	$-\frac{5}{12}$	$-\frac{3}{4}$
0	0	0	0	0	*	0	0	0	0
$-\frac{1}{2}$	$-\frac{3}{4}$	$-\frac{5}{12}$	0	$\frac{5}{12}$	*	$-\frac{5}{12}$	0	$\frac{5}{12}$	$\frac{3}{4}$
-1	$-\frac{3}{2}$	$-\frac{5}{6}$	0	$\frac{3}{2}$	*	$-\frac{3}{2}$	0	$\frac{5}{6}$	$\frac{3}{2}$
$-\frac{3}{2}$	$-\frac{9}{4}$	$-\frac{5}{4}$	0	$\frac{5}{4}$	*	$-\frac{5}{4}$	0	$\frac{5}{4}$	$\frac{9}{4}$
-2	-3	$-\frac{4}{3}$	0	3	*	-3	0	$\frac{4}{3}$	3

2c See part h.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations


2d See part h.

2e The lines $x = 1$ and $x = -1$ are isoclines. $y = 0$ is also an isocline, ignoring the point $(0, 0)$ where the gradient is undefined.

2f $y = 0$. Yes, $y = 0$ is a constant solution.

2g Odd: y' is unchanged when x is replaced with $-x$ and y is replaced with $-y$.

2c,d,h



3a $y = Cx^2 e^x$

$y' = C(e^x \times 2x + x^2 \times e^x)$

Substituting for y and y' in $xy' = y(2 + x)$:

LHS $= xy'$

$= Cx(2xe^x + x^2e^x)$

$= Cx^2e^x(2 + x)$

$= y(2 + x)$

$=$ RHS

Therefore, $y = Cx^2 e^x$ is a solution of $xy' = y(2 + x)$.

© Cambridge University Press 2019

3b $y = \sqrt{x^2 + C} = (x^2 + C)^{\frac{1}{2}}$

$y' = \frac{1}{2}(x^2 + C)^{-\frac{1}{2}} \times 2x$

$y' = x(x^2 + C)^{-\frac{1}{2}}$

Substituting for y and y' in $yy' = x$:

LHS $= yy'$

$= (x^2 + C)^{\frac{1}{2}} \times x(x^2 + C)^{-\frac{1}{2}}$

$= x(x^2 + C)^0$

$= x$

$=$ RHS

Therefore, $y = \sqrt{x^2 + C}$ is a solution of $yy' = x$.

3c

$y = \dfrac{1}{x^2 + C} = (x^2 + C)^{-1}$

$y' = -(x^2 + C)^{-2} \times 2x$

$= -2x(x^2 + C)^{-2}$

$= -\dfrac{2x}{(x^2 + C)^2}$

Substituting for y and y' in $y' = -2xy^2$

RHS $= -2x \times \left(\dfrac{1}{x^2 + C}\right)^2$

$= -\dfrac{2x}{(x^2 + C)^2}$

$=$ LHS

Therefore, $y = \dfrac{1}{x^2 + C}$ is a solution of $y' = -2xy^2$.

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

4a $y = 0$

4b Taking the reciprocal of the DE gives

$$\frac{dx}{dy} = -\frac{2}{y}$$

4c

$$\int \frac{dx}{dy}\, dy = -\int \frac{2}{y}\, dy$$

$$x = -2\ln|y| + C$$

4d $x = -2\ln|y| + C$

$x - C = -2\ln|y|$

$\frac{C - x}{2} = \ln|y|$

$|y| = e^{\left(\frac{C-x}{2}\right)}$

$y = e^{\left(\frac{C}{2}\right)} \times e^{\left(-\frac{x}{2}\right)}$ or $-e^{\left(\frac{C}{2}\right)} \times e^{\left(-\frac{x}{2}\right)}$

$y = Ae^{-\frac{x}{2}}$ where $A = e^{\left(\frac{C}{2}\right)}$ or $-e^{\left(\frac{C}{2}\right)}$

4e $y = 0$ when $A = 0$

4f Substituting $y(0) = 3$:

$3 = Ae^{-\frac{0}{2}}$

$A = 3$

Therefore $y = 3e^{-\frac{x}{2}}$


5a i $y' = 0$ when $x = 2$ or $x = -2$

© Cambridge University Press 2019

5a ii It is an isocline.

5a iii Gradients decrease to -1 then increase.

5a iv




5b i $y' = 0$ when $y = 2$ or $y = -2$

5b ii Gradients decrease to $-\frac{1}{2}$ then increase.

5b iii It is an isocline.

5b iv




5c i $y' = 0$ when $y = -x$

© Cambridge University Press 2019

5c ii Gradients decrease

5c iii Gradients increase

5c iv



6a $y = Ae^{-x} + Be^{-2x}$

$y' = -Ae^{-x} - 2Be^{-2x}$

$y'' = Ae^{-x} + 4Be^{-2x}$

Substituting for y'', y' and y in $y'' + 3y' + 2y = 0$:

$\text{LHS} = Ae^{-x} + 4Be^{-2x} + 3(-Ae^{-x} - 2Be^{-2x}) + 2(Ae^{-x} + Be^{-2x})$

$= Ae^{-x} + 4Be^{-2x} - 3Ae^{-x} - 6Be^{-2x} + 2Ae^{-x} + 2Be^{-2x}$

$= 3Ae^{-x} - 3Ae^{-x} + 6Be^{-2x} - 6Be^{-2x}$

$= 0$

$= \text{RHS}$

Therefore, $y = Ae^{-x} + Be^{-2x}$ is a solution of $y'' + 3y' + 2y = 0$.

© Cambridge University Press 2019

6b $y = Ae^{-x}\cos 2x + Be^{-x}\sin 2x$

$y' = A(\cos 2x \times -e^{-x} + e^{-x} \times -2\sin 2x) + B(\sin 2x \times -e^{-x} + e^{-x} \times 2\cos 2x)$

$= A(-e^{-x}\cos 2x - 2e^{-x}\sin 2x) + B(-e^{-x}\sin 2x + 2e^{-x}\cos 2x)$

$= -Ae^{-x}(\cos 2x + 2\sin 2x) - Be^{-x}(\sin 2x - 2\cos 2x)$

$y'' = -A\big(-e^{-x}(\cos 2x + 2\sin 2x) + e^{-x}(-2\sin 2x + 4\cos 2x)\big) - B\big(-e^{-x}(\sin 2x - 2\cos 2x) + e^{-x}(2\cos 2x + 4\sin 2x)\big)$

$= -Ae^{-x}(-\cos 2x - 2\sin 2x - 2\sin 2x + 4\cos 2x) - Be^{-x}(-\sin 2x + 2\cos 2x + 2\cos 2x + 4\sin 2x)$

$= -Ae^{-x}(3\cos 2x - 4\sin 2x) - Be^{-x}(3\sin 2x + 4\cos 2x)$

Substituting y'', y' and y in $y'' + 2y' + 5y = 0$:

LHS

$= -Ae^{-x}(3\cos 2x - 4\sin 2x) - Be^{-x}(3\sin 2x + 4\cos 2x) + 2\big(-Ae^{-x}(\cos 2x + 2\sin 2x) - Be^{-x}(\sin 2x - 2\cos 2x)\big) + 5Ae^{-x}\cos 2x + 5Be^{-x}\sin 2x$

$= e^{-x}\cos 2x\,(-3A - 4B - 2A + 4B + 5A) + e^{-x}\sin 2x\,(4A - 3B - 4A - 2B + 5B)$

$= e^{-x}\cos 2x\,(0) + e^{-x}\sin 2x\,(0)$

$= 0$

$=$ RHS

Therefore, $y = Ae^{-x}\cos 2x + Be^{-x}\sin 2x$ is a solution of $y'' + 2y' + 5y = 0$.

6c $y = A\cos x + B\sin x + Ce^{2x}$

$y' = -A\sin x + B\cos x + 2Ce^{2x}$

$y'' = -A\cos x - B\sin x + 4Ce^{2x}$

$y''' = A\sin x - B\cos x + 8Ce^{2x}$

Substituting y''', y'', y' and y in $y''' - 2y'' + y' - 2y = 0$:

LHS

$= A\sin x - B\cos x + 8Ce^{2x} - 2(-A\cos x - B\sin x + 4Ce^{2x}) + (-A\sin x + B\cos x + 2Ce^{2x}) - 2(A\cos x + B\sin x + Ce^{2x})$

$= A\sin x - B\cos x + 8Ce^{2x} + 2A\cos x + 2B\sin x - 8Ce^{2x} - A\sin x + B\cos x + 2Ce^{2x} - 2A\cos x - 2B\sin x - 2Ce^{2x}$

© Cambridge University Press 2019

$= 0$

$=$ RHS

Therefore, $y = A\cos x + B\sin x + Ce^{2x}$ is a solution of $y''' - 2y'' + y' - 2y = 0$.

6d $y = Ae^{2x} + Be^{-2x} + C\cos 2x + D\cos 2x$

$y' = 2Ae^{2x} - 2Be^{-2x} - 2C\sin 2x - 2D\sin 2x$

$y'' = 4Ae^{2x} + 4Be^{-2x} - 4C\cos 2x - 4D\cos 2x$

$y''' = 8Ae^{2x} - 8Be^{-2x} + 8C\sin 2x + 8D\sin 2x$

$y'''' = 16Ae^{2x} + 16Be^{-2x} + 16C\cos 2x + 16D\cos 2x$

Substituting y'''' and y in $y'''' - 16y = 0$:

$\text{LHS} = 16Ae^{2x} + 16Be^{-2x} + 16C\cos 2x + 16D\cos 2x$

$-16(Ae^{2x} + Be^{-2x} + C\cos 2x + D\cos 2x)$

$= 0$

$=$ RHS

Therefore, $y = Ae^{2x} + Be^{-2x} + C\cos 2x + D\cos 2x$ is a solution of $y'''' - 16y = 0$.

7a

$\frac{dy}{dx} = \frac{-2xy}{1+x^2}$

$\int \frac{1}{y}\,dy = -\int \frac{2x}{1+x^2}\,dx$

$\ln|y| = -\ln|1+x^2| + B$

$\ln|y| + \ln|1+x^2| = B$

$\ln|y(1+x^2)| = B$

$|y(1+x^2)| = e^B$

$y(1+x^2) = e^B$ or $-e^B$

$y(1+x^2) = C$ where $C = e^B$ or $-e^B$

$y = \frac{C}{1+x^2}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

7b

$$\frac{dy}{dx} = \frac{1-x}{2+y}$$

$$\int (2+y)\,dy = \int (1-x)\,dx$$

$$2y + \frac{y^2}{2} = x - \frac{x^2}{2} + B$$

$$y^2 + 4y = -x^2 + 2x + 2B$$

$$y^2 + 4y + 4 - 4 = -\left((x^2 - 2x + 1) - 1\right) + 2B$$

$$(y+2)^2 - 4 = -((x-1)^2 - 1) + 2B$$

$$(x-1)^2 + (y+2)^2 = 1 + 4 + 2B$$

$$(x-1)^2 + (y+2)^2 = 5 + 2B$$

$(x-1)^2 + (y+2)^2 = C$ where $C = 5 + 2B$

7c

$$\frac{dy}{dx} = \frac{y(x-1)}{x}$$

$$\int \frac{1}{y}\,dy = \int \frac{x-1}{x}\,dx$$

$$\int \frac{1}{y}\,dy = \int \left(1 - \frac{1}{x}\right) dx$$

$$\ln|y| = x - \ln|x| + B$$

$$\ln|x| + \ln|y| = x + B$$

$$\ln|xy| = x + B$$

$$|xy| = e^{x+B}$$

$xy = e^x \times e^B$ or $-e^x \times e^B$

$xy = e^x \times C$ where $C = e^B$ or $-e^B$

$$y = \frac{Ce^x}{x}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

8a

$$\frac{dy}{dx} = \frac{1}{2}(1-y) = \frac{1-y}{2}$$

$$\frac{dx}{dy} = \frac{2}{1-y}$$

$$\frac{dx}{dy} = \frac{-2}{y-1}$$

$$\int \frac{dx}{dy}\,dy = -\int \frac{2}{y-1}\,dy$$

$$x = -2\log_e|y-1| + C$$

$$\log_e|y-1| = -\frac{1}{2}(x-C)$$

$$|y-1| = e^{-\frac{1}{2}(x-C)}$$

$$y-1 = e^{-\frac{1}{2}(x-C)} \text{ or } -e^{-\frac{1}{2}(x-C)}$$

$$y-1 = e^{-\frac{1}{2}x} \times e^{\frac{1}{2}C} \text{ or } -e^{-\frac{1}{2}x} \times e^{\frac{1}{2}C}$$

$$y-1 = Ae^{-\frac{1}{2}x} \text{ where } A = e^{\frac{1}{2}C} \text{ or } -e^{\frac{1}{2}C}$$

$$y = 1 + Ae^{-\frac{1}{2}x}$$

Substituting $y(0) = 2$:

$$2 = 1 + Ae^{-\frac{1}{2}\times 0}$$

$$A = 2 - 1$$

$$A = 1$$

Therefore $y = 1 + e^{-\frac{1}{2}x}$

© Cambridge University Press 2019

8b

$$\frac{dy}{dx} = \frac{1}{5}(5 - y) = \frac{5 - y}{5}$$

$$\frac{dx}{dy} = \frac{5}{5 - y}$$

$$\frac{dx}{dy} = \frac{-5}{y - 5}$$

$$\int \frac{dx}{dy}\, dy = -\int \frac{5}{y - 5}\, dy$$

$$x = -5 \log_e |y - 5| + C$$

$$\log_e |y - 5| = -\frac{1}{5}(x - C)$$

$$|y - 5| = e^{-\frac{1}{5}(x - C)}$$

$$y - 5 = e^{-\frac{1}{5}(x - C)} \text{ or } -e^{-\frac{1}{5}(x - C)}$$

$$y - 5 = e^{-\frac{1}{5}x} \times e^{\frac{1}{5}C} \text{ or } -e^{-\frac{x}{5}} \times e^{\frac{1}{5}C}$$

$$y - 5 = Ae^{-\frac{1}{5}x} \text{ where } A = e^{\frac{1}{5}C} \text{ or } -e^{\frac{1}{5}C}$$

$$y = 5 + Ae^{-\frac{1}{5}x}$$

Substituting $y(0) = 2$:

$$2 = 5 + Ae^{-\frac{1}{5} \times 0}$$

$$A = 2 - 5$$

$$A = -3$$

Therefore $y = 5 - 3e^{-\frac{1}{5}x}$

9a Let

$$\frac{1}{1 - x^2} = \frac{A}{1 - x} + \frac{B}{1 + x}$$

$$= \frac{A(1 + x) + B(1 - x)}{(1 + x)(1 - x)}$$

$$= \frac{A + Ax + B - Bx}{(1 + x)(1 - x)}$$

© Cambridge University Press 2019

$$= \frac{(A-B)x + A + B}{(1+x)(1-x)}$$

Equating coefficients in the numerators:

$A + B = 1 \qquad (1)$

$A - B = 0 \qquad (2)$

$(1) + (2)$:

$2A = 1$

$A = \frac{1}{2}$

Substituting $A = \frac{1}{2}$ into (1):

$\frac{1}{2} + B = 1$

$B = \frac{1}{2}$

So $\frac{1}{1-x^2} = \frac{1}{2(1-x)} + \frac{1}{2(1+x)}$

or $\frac{1}{1-x^2} = \frac{1}{2}\left(\frac{1}{1-x} + \frac{1}{1+x}\right)$

9b

$\frac{dy}{dx} = \frac{y}{1-x^2}$

$\frac{1}{y}\,dy = \frac{1}{1-x^2}\,dx$

$\int \frac{1}{y}\,dy = \int \frac{1}{1-x^2}\,dx$

$\int \frac{1}{y}dy = \int \left(\frac{\frac{1}{2}}{1-x} + \frac{\frac{1}{2}}{1+x}\right)dx$

$\log_e |y| = -\frac{1}{2}\log_e |1-x| + \frac{1}{2}\log_e |1+x| + B$

$\log_e |y| = \frac{1}{2}\log_e \left|\frac{1+x}{1-x}\right| + B$

© Cambridge University Press 2019

$$\log_e |y| = \log_e \left|\frac{1+x}{1-x}\right|^{\frac{1}{2}} + \log_e e^B$$

$$\log_e |y| = \log_e \left(e^B \left|\frac{1+x}{1-x}\right|^{\frac{1}{2}}\right)$$

$$|y| = e^B \left|\frac{1+x}{1-x}\right|^{\frac{1}{2}}$$

$$y = e^B \left|\frac{1+x}{1-x}\right|^{\frac{1}{2}} \text{ or } -e^B \left|\frac{1+x}{1-x}\right|^{\frac{1}{2}}$$

$$y = C\left|\frac{1+x}{1-x}\right|^{\frac{1}{2}} \text{ where } C = e^B \text{ or } -e^B$$

$$y = C\sqrt{\left|\frac{1+x}{1-x}\right|}$$

10a For domain: $1 - e^{-x} \neq 0$

$e^{-x} \neq 1$

$-x \neq \log_e 1$

$x \neq 0$

Also $L(x) \neq 0$, so there are no intercepts.

10b

As $x \to \infty, e^{-x} \to 0$ so $L(x) \to \frac{1}{1-0}$. Hence $\lim_{x\to\infty} L(x) = 1$

As $x \to -\infty, e^{-x} \to \infty$ so $L(x) \to \frac{1}{-\infty}$. Hence $\lim_{x\to-\infty} L(x) = 0$

10c $x \neq 0$ (from part a) so there is a vertical asymptote at $x = 0$.

As $x \to 0^+, e^{-x} \to 1^-$ so $L(x) \to \frac{1}{0^+}$. That is $L(x) \longrightarrow \infty$.

As $x \to 0^-, e^{-x} \to 1^+$ so $L(x) \to \frac{1}{0^-}$. That is $L(x) \longrightarrow -\infty$.

© Cambridge University Press 2019

10d For $x = \log_e 2$:

$$y = L(\log_e 2)$$

$$= \frac{1}{1 - e^{-(\log_e 2)}}$$

$$= \frac{1}{1 - e^{\log_e 2^{\frac{1}{2}}}}$$

$$= \frac{1}{1 - \frac{1}{2}}$$

$$= 2$$

For $x = -\log_e 2$:

$$y = L(-\log_e 2)$$

$$= \frac{1}{1 - e^{\log_e 2}}$$

$$= \frac{1}{1 - 2}$$

$$= -1$$

10e i $L(x) = (1 - e^{-x})^{-1}$

$$L'(x) = -(1 - e^{-x})^{-2} \times e^{-x}$$

$$= -\frac{e^{-x}}{(1 - e^{-x})^2}$$

$$= -\frac{e^{-x}}{1 - 2e^{-x} + e^{-2x}}$$

$$= -\frac{1}{e^x(1 - 2e^{-x} + e^{-2x})}$$

$$= \frac{-1}{e^x - 2 + e^{-x}}$$

$$= \frac{-1}{e^x - 2\left(e^{\frac{x}{2}}\right)\left(e^{-\frac{x}{2}}\right) + e^{-x}}$$

$$= \frac{-1}{\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)^2}$$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

10e ii $L'(x) = -\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)^{-2}$

$$L''(x) = 2\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)^{-3} \times \left(\frac{e^{\frac{x}{2}}}{2} + \frac{e^{-\frac{x}{2}}}{2}\right)$$


$$= \frac{2}{\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)^{3}} \times \left(\frac{e^{\frac{x}{2}} + e^{-\frac{x}{2}}}{2}\right)$$

$$= \frac{e^{\frac{x}{2}} + e^{-\frac{x}{2}}}{\left(e^{\frac{x}{2}} - e^{-\frac{x}{2}}\right)^{3}}$$

10f When $x > 0$, $L''(x) > 0$ and when $x < 0, L''(x) < 0$.

Hence graph is concave up for $x > 0$ and concave down for $x < 0$.

10g



10h Substituting for y' and y in $y' = y(1 - y)$:

$$\text{LHS} = -\frac{e^{-x}}{(1 - e^{-x})^2}$$

$$\text{RHS} = \frac{1}{1 - e^{-x}}\left(1 - \frac{1}{1 - e^{-x}}\right)$$

$$= \frac{1}{1 - e^{-x}} - \frac{1}{(1 - e^{-x})^2}$$

© Cambridge University Press 2019

$$= \frac{(1 - e^{-x}) - 1}{(1 - e^{-x})^2}$$

$$= -\frac{e^{-x}}{(1 - e^{-x})^2}$$

$$= \text{LHS}$$

Therefore, since LHS = RHS, $y = L(x)$ is a solution of the DE.

11 The graph in option C corresponds to $y' = \frac{1}{4}(x^2 + y^2)$.

Slope must be zero at the origin (eliminates A) and positive everywhere else (eliminates B and D which have zero slope on the x and y-axes respectively).

12 Option B $y' = 1 + \frac{y}{x}$ corresponds to the graph.

The diagram shows slope of zero on the isocline $y = -x$, which eliminates options C and D. For $x = 1$, slope grows more extreme with increasing y, which eliminates A.

13a

$$\frac{dy}{dx} = -\frac{y}{x}$$

$$\int \frac{1}{y}\, dy = -\int \frac{1}{x}\, dx$$

$$\log_e |y| = -\log_e |x| + C$$

$$\log_e |x| + \log_e |y| = C$$

$$\log_e |xy| = C$$

$$|xy| = e^C$$

$$xy = e^C \text{ or } -e^C$$

$$xy = A \text{ where } A = e^C \text{ or } -e^C$$

$$y = \frac{A}{x}$$

© Cambridge University Press 2019

Substituting $y(2) = 1$:

$1 = \frac{A}{2}$

$A = 2$

Therefore $y = \frac{2}{x}$

13b

$\frac{dy}{dx} = (1 + 2x)e^{-y}$

$\int \frac{1}{e^{-y}}\, dy = \int (1 + 2x)\, dx$

$\int e^{y}\, dy = \int (1 + 2x)\, dx$

$e^{y} = x + x^2 + C$

$y = \ln(x + x^2 + C)$

Substituting $y(1) = 0$:

$0 = \ln(1 + 1 + C)$

$0 = \ln(2 + C)$

$1 = 2 + C$

$C = -1$

Therefore $y = \ln(x + x^2 - 1)$

13c

$\frac{dy}{dx} = \frac{y^2}{\sqrt{x}}$

$\int \frac{1}{y^2}\, dy = \int \frac{1}{\sqrt{x}}\, dx$

$\int y^{-2}\, dy = \int x^{-\frac{1}{2}}\, dx$

$-y^{-1} = 2x^{\frac{1}{2}} + C$

© Cambridge University Press 2019

$$-\frac{1}{y} = 2x^{\frac{1}{2}} + C$$

$$-y = \frac{1}{2x^{\frac{1}{2}} + C}$$

$$y = -\frac{1}{2\sqrt{x} + C}$$

Substituting $y(0) = -1$:

$$-1 = -\frac{1}{2\sqrt{0} + C}$$

$$-1 = -\frac{1}{C}$$

$$C = 1$$

Therefore $y = -\frac{1}{2\sqrt{x} + 1}$

14a Let

$$\frac{1}{y(1-y)} = \frac{A}{y} + \frac{B}{1-y}$$

$$= \frac{A(1-y) + By}{y(1-y)}$$

$$= \frac{A - Ay + By}{y(1-y)}$$

$$= \frac{(B-A)y + A}{y(1-y)}$$

Equating coefficients in the numerators:

$A = 1$

$B - A = 0 \qquad (1)$

Substituting $A = 1$ into (1):

$B - 1 = 0$

$B = 1$

Therefore $\frac{1}{y(1-y)} = \frac{1}{y} + \frac{1}{1-y}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

14b i $y = 0$ and $y = 1$

14b ii

$$\frac{dy}{dx} = y(1-y)$$

$$\int \frac{1}{y(1-y)} dy = \int 1\, dx$$

$$\int \left(\frac{1}{y} + \frac{1}{1-y}\right) dy = \int 1\, dx$$

$$\log_e |y| - \log_e |1-y| = x + C$$

$$\log_e \left|\frac{y}{1-y}\right| = x + C$$

$$\left|\frac{y}{1-y}\right| = e^{x+C}$$

$$\frac{y}{1-y} = e^x e^C \text{ or } -e^x e^C$$

$$\frac{y}{1-y} = Ae^x \text{ where } A = e^C \text{ or } -e^C$$

$$y = Ae^x(1-y)$$

$$y = Ae^x - Ae^x y$$

$$Ae^x y + y = Ae^x$$

$$y(Ae^x + 1) = Ae^x$$

$$y = \frac{Ae^x}{Ae^x + 1}$$

$$y = \frac{Ae^x}{Ae^x + 1} \times \frac{A^{-1}e^{-x}}{A^{-1}e^{-x}}$$

$$y = \frac{1}{A^{-1}e^{-x}(Ae^x + 1)}$$

$$y = \frac{1}{1 + A^{-1}e^{-x}}$$

$$y = \frac{1}{1 + Be^{-x}} \text{ where } B = A^{-1}$$

© Cambridge University Press 2019

14b iii Substituting $\left(0, \frac{1}{4}\right)$:

$$\frac{1}{4} = \frac{1}{1 + Be^0}$$

$$\frac{1}{4} = \frac{1}{1 + B}$$

$$B = 3$$

Therefore $y = \frac{1}{1 + 3e^{-x}}$

15a Let

$$\frac{1}{(2-y)(3-y)} = \frac{A}{2-y} + \frac{B}{3-y}$$

$$= \frac{A(3-y) + B(2-y)}{(2-y)(3-y)}$$

$$= \frac{3A - Ay + 2B - By}{(2-y)(3-y)}$$

$$= \frac{(-A-B)y + 3A + 2B}{(2-y)(1-y)}$$

Equating coefficients in the numerators:

$-A - B = 0$ (1)

$3A + 2B = 1$ (2)

$2 \times (1) + (2)$:

$A = 1$

Substituting $A = 1$ into (1):

$-1 - B = 0$

$B = -1$

Therefore $\frac{1}{(2-y)(3-y)} = \frac{1}{2-y} - \frac{1}{3-y}$

or $\frac{1}{(2-y)(3-y)} = \frac{1}{y-3} - \frac{1}{y-2}$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

15b i $y = 2$ and $y = 3$

15b ii $y' = -\frac{1}{5}(3-y)(2-y)$

$$\frac{dy}{dx} = -\frac{1}{5}(3-y)(2-y)$$

$$\frac{1}{(3-y)(2-y)}\,dy = -\frac{1}{5}\,dx$$

$$\int \left(\frac{1}{y-3} - \frac{1}{y-2}\right) dy = \int -\frac{1}{5}\,dx$$

$$\log_e |y-3| - \log_e |y-2| = -\frac{1}{5}x + C$$

$$\log_e \left|\frac{y-3}{y-2}\right| = -\frac{1}{5}x + C$$

$$\left|\frac{y-3}{y-2}\right| = e^{-\frac{1}{5}x+C}$$

$$\frac{y-3}{y-2} = e^{-\frac{1}{5}x}e^C \text{ or } -e^{-\frac{1}{5}x}e^C$$

$$\frac{y-3}{y-2} = Ae^{-\frac{1}{5}x} \text{ where } A = e^C \text{ or } -e^C$$

$$y-3 = Ae^{-\frac{1}{5}x}(y-2)$$

$$y-3 = Ae^{-\frac{1}{5}x}y - 2Ae^{-\frac{1}{5}x}$$

$$y - Ae^{-\frac{1}{5}x}y = 3 - 2Ae^{-\frac{1}{5}x}$$

$$y\left(1 - Ae^{-\frac{1}{5}x}\right) = 3 - 2Ae^{-\frac{1}{5}x}$$

$$y = \frac{3 - 2Ae^{-\frac{1}{5}x}}{1 - Ae^{-\frac{1}{5}x}}$$

Substituting $y(0) = 1$:

$$1 = \frac{3 - 2Ae^0}{1 - Ae^0}$$

$$1 = \frac{3-2A}{1-A}$$

© Cambridge University Press 2019

$$1 - A = 3 - 2A$$

$$A = 2$$

Therefore $y = \dfrac{3 - 4e^{-\frac{1}{5}x}}{1 - 2e^{-\frac{1}{5}x}}$

15b iii $y = 0$ when:

$$0 = \frac{3 - 4e^{-\frac{1}{5}x}}{1 - 2e^{-\frac{1}{5}x}}$$

$$3 - 4e^{-\frac{1}{5}x} = 0$$

$$e^{-\frac{1}{5}x} = \frac{3}{4}$$

$$-\frac{1}{5}x = \ln\frac{3}{4}$$

$$x = -5\ln\frac{3}{4}$$

$$x = 5\ln\left(\frac{3}{4}\right)^{-1}$$

$$x = 5\ln\frac{4}{3}$$

or $x \div 1.44$

16a Let

$$\frac{5}{N(5-N)} = \frac{A}{N} + \frac{B}{5-N}$$

$$= \frac{A(5-N) + BN}{N(5-N)}$$

$$= \frac{5A - AN + BN}{N(5-N)}$$

$$= \frac{(B-A)N + 5A}{N(5-N)}$$

© Cambridge University Press 2019

Equating coefficients in the numerators:

$5A = 5$ so $A = 1$

$B - A = 0 \qquad (1)$

Substituting $A = 1$ into (1):

$B - 1 = 0$

$B = 1$

Therefore $\dfrac{5}{N(5-N)} = \dfrac{1}{N} + \dfrac{1}{5-y}$

16b

$\dfrac{dN}{dt} = kN(5-N)$

$\dfrac{1}{N(5-N)}\, dN = k\, dt$

$\displaystyle\int \frac{1}{N(5-N)} dN = \int k\, dt$

$\displaystyle\int \frac{5}{N(5-N)} dN = \int 5k\, dt$

$\displaystyle\int \left(\frac{1}{N} + \frac{1}{5-N}\right) dN = \int 5k\, dt$

$\log_e |N| - \log_e |5-N| = 5kt + C$

$\log_e \left|\dfrac{N}{5-N}\right| = 5kt + C$

$\left|\dfrac{N}{5-N}\right| = e^{5kt+C}$

$\dfrac{N}{5-N} = e^{5kt+C}$ or $-e^{5kt+C}$

$\dfrac{N}{5-N} = e^C e^{5kt}$ or $-e^C e^{5kt}$

$\dfrac{N}{5-N} = Ae^{5kt}$ where $A = e^C$ or $-e^C$

$N = Ae^{5kt}(5-N)$

$N = 5Ae^{5kt} - NAe^{5kt}$

© Cambridge University Press 2019

$$NAe^{5kt} + N = 5Ae^{5kt}$$

$$N(Ae^{5kt} + 1) = 5Ae^{5kt}$$

$$N = \frac{5Ae^{5kt}}{Ae^{5kt} + 1}$$

$$N = \frac{5Ae^{5kt}}{Ae^{5kt} + 1} \times \frac{A^{-1}e^{-5kt}}{A^{-1}e^{-5kt}}$$

$$N = \frac{5}{A^{-1}e^{-5kt}(Ae^{5kt} + 1)}$$

$$N = \frac{5}{1 + A^{-1}e^{-5kt}}$$

$$N = \frac{5}{1 + Be^{-5kt}} \text{ where } B = A^{-1}$$

16c Substituting $N(0) = 1$:

$$1 = \frac{5}{1 + Be^{0}}$$

$$1 + B = 5$$

$$B = 4$$

Therefore $N = \frac{5}{1 + 4e^{-5kt}}$

16d Total sales at $t = 2$ is 1 million phones already sold at $t = 0$, plus 0.4 million phones sold between $t = 0$ and $t = 2$. So $N(2) = 1.4$ (as N is the number of people in millions).

Substituting $N(2) = 1.4$:

$$1.4 = \frac{5}{1 + 4e^{-5k(2)}}$$

$$1 + 4e^{-10k} = \frac{5}{1.4}$$

$$1 + 4e^{-10k} = \frac{25}{7}$$

$$4e^{-10k} = \frac{25}{7} - 1$$

© Cambridge University Press 2019

$$4e^{-10k} = \frac{18}{7}$$

$$e^{-10k} = \frac{9}{14}$$

$$-10k = \ln\frac{9}{14}$$

$$k = -\frac{1}{10}\ln\frac{9}{14}$$

$$k \doteqdot 4.418 \times 10^{-2}$$

16e When $t = 3$,

$$N = \frac{5}{1 + 4e^{-5\left(-\frac{1}{10}\ln\frac{9}{14}\right)(3)}}$$

$$\doteqdot 1.633$$

Sales during year 3 equals total sales in three years (1.633 million) minus sales up to year 2 (1.4 million).

That is, the projected sales in the third year will be 0.233 million or 233,000.

17a $y' = x(1 - 2y)$

$= x - 2xy$

$y'' = 1 - 2y \times 1 - 2x \times y'$

$= 1 - 2y - 2xy'$

17b Turning point when $y' = 0$:

$x(1 - 2y) = 0$

$x = 0$

When $x = 0, y = 1$, so

$y'' = 1 - 2 \times 1 - 2 \times 0 \times 0$

$= 1 - 2$

$= -1$

Since $y'' < 0$, the turning point at $x = 0$ is a maximum.

© Cambridge University Press 2019

17c $y' = x(1 - 2y)$

$\frac{dy}{dx} = x(1 - 2y)$

$\int \frac{1}{1-2y} dy = \int x\, dx$

$-\frac{1}{2}\int \frac{2}{2y-1} dy = \int x\, dx$

$\int \frac{2}{2y-1} dy = -2\int x\, dx$

$\ln|2y - 1| = -x^2 + C$

$|2y - 1| = e^{-x^2+C}$

$|2y - 1| = e^C e^{-x^2}$

$2y - 1 = e^C e^{-x^2}$ or $-e^C e^{-x^2}$

$2y - 1 = Ae^{-x^2}$ where $A = e^C$ or $-e^C$

$2y = 1 + Ae^{-x^2}$

$y = \frac{1}{2}(1 + Ae^{-x^2})$

Substituting $y(0) = 1$:

$1 = \frac{1}{2}(1 + Ae^0)$

$2 = 1 + A$

$A = 1$

Therefore $y = \frac{1}{2}(1 + e^{-x^2})$

18a The solutions are $y = F(x) + C$ where $F'(x) = f(x)$.

Each is a vertical shift of the other.

18b The solutions are $y = G^{-1}(x + C)$ where $G'(y) = \frac{1}{g(y)}$.

Each is a horizontal shift of the other.

© Cambridge University Press 2019

19a $y = 1, y = -1$

19b $y_1 = \cos(x + A)$

$y_1' = -\sin(x + A)$

Substituting y' and y in $(y')^2 = 1 - y^2$:

$\text{LHS} = (-\sin(x + A))^2$

$= \sin^2(x + A)$

$= 1 - \cos^2(x + A)$

$= 1 - y^2$

$= \text{RHS}$

Therefore $y_1 = \cos(x + A)$ is a solution of $(y')^2 = 1 - y^2$.

$y_2 = \sin(x + B)$

$y_2' = \cos(x + B)$

Substituting y' and y in $(y')^2 = 1 - y^2$:

$\text{LHS} = (\cos(x + B))^2$

$= \cos^2(x + B)$

$= 1 - \sin^2(x + B)$

$= 1 - y^2$

$= \text{RHS}$

Therefore $y_2 = \sin(x + B)$ is a solution of $(y')^2 = 1 - y^2$.

19c $\cos(x + A) = \cos x \cos A - \sin x \sin A$

$\sin(x + B) = \sin x \cos B + \sin B \cos x$

Let $B = A + \frac{\pi}{2} + 2n\pi$ for integer n

Then $\cos B = \cos\left(A + \frac{\pi}{2} + 2n\pi\right)$

$\cos B = \cos\left(A + \frac{\pi}{2}\right)$

$= \cos A \cos\frac{\pi}{2} - \sin A \sin\frac{\pi}{2}$

© Cambridge University Press 2019

$= \cos A \times 0 - \sin A \times 1$

$= -\sin A$

$\sin B = \sin\left(A + \frac{\pi}{2} + 2n\pi\right)$

$\sin B = \sin\left(A + \frac{\pi}{2}\right)$

$= \sin A \cos\frac{\pi}{2} + \cos A \sin\frac{\pi}{2}$

$= \sin A \times 0 + \cos A \times 1$

$= \cos A$

Substituting $\cos B = -\sin A$ and $\sin B = \cos A$ in the expansion for $\sin(x + B)$:

$\sin(x + B) = \sin x \times -\sin A + \cos A \times \cos x$

$= -\sin x \sin A + \cos A \cos x$

$= \cos(x + A)$

So the two solutions are identical.

19d $B = A + \frac{\pi}{2}$

(More precisely, $B = A + \frac{\pi}{2} + 2n\pi$, for some integer n.)

20a Using Pythagoras' theorem:

$AB^2 + AC^2 = BC^2$

$y^2 + AC^2 = 4^2$

$AC^2 = 16 - y^2$

$AC = \sqrt{16 - y^2}$

$\text{Gradient} = -\frac{AB}{AC}$

$\frac{dy}{dx} = -\frac{y}{\sqrt{16 - y^2}}$


20b $-4 \leq y \leq 4; y(0) = 4$

© Cambridge University Press 2019

Chapter 13 worked solutions – Differential equations

20c $y = 0$, not a solution of the IVP

20d



© Cambridge University Press 2019