1999 HIGHER SCHOOL CERTIFICATE SOLUTIONS

3 UNIT (ADDITIONAL) AND 3/4 UNIT (COMMON) MATHEMATICS

QUESTION 1

(a) $\displaystyle\int_0^{\sqrt{3}} \frac{dx}{\sqrt{4-x^2}} = \left[\sin^{-1}\frac{x}{2}\right]_0^{\sqrt{3}}$ (standard integral)

$= \sin^{-1}\frac{\sqrt{3}}{2} - \sin^{-1} 0$

$= \frac{\pi}{3} - 0$

$= \frac{\pi}{3}.$

(b) $\frac{d}{dx}\sin^3 x = 3\sin^2 x . \frac{d}{dx}\sin x$

$= 3\sin^2 x \cos x.$

(c) $A(-2, 7)$, $B(8, -8)$

Ratio 2 : 3

For P, $x = \frac{2\times 8 + 3\times(-2)}{2+3}$

$= \frac{10}{5}$

$= 2,$

and $y = \frac{2\times(-8)+3\times 7}{2+3}$

$= \frac{5}{5}$

$= 1.$

$\therefore$ P is $(2, 1)$.

(d) Asymptote when denominator is zero, that is, $x - 3 = 0$ or $x = 3$.

(e) $P(x) = x^3 - 4x$

$P(-3) = -27 + 12$

$= -15$ is the remainder.

(f) If $u = \tan x$

$\frac{du}{dx} = \sec^2 x \Rightarrow du = \sec^2 x\, dx.$

When $x = 0$, $u = 0$.

When $x = \frac{\pi}{3}$, $u = \sqrt{3}$.

$\therefore \displaystyle\int_0^{\frac{\pi}{3}} \tan^2 x \sec^2 x\, dx = \int_0^{\sqrt{3}} u^2\, du$

$= \left[\frac{u^3}{3}\right]_0^{\sqrt{3}}$

$= \sqrt{3}.$

QUESTION 2

(a) Number of ways of choosing 3 females from 7 is ${}^7C_3$. The other two must be male. The number of ways of choosing 2 from 4 is ${}^4C_2$.

$\therefore$ Number of committees $= {}^7C_3 \times {}^4C_2$

$= 210.$

(b) **Method 1:**

$\cos\theta + \sqrt{3}\sin\theta = 1$

Now $R\cos(\theta - \alpha) = R\cos\theta\cos\alpha + R\sin\theta\sin\alpha$

$\therefore$ $R\cos\alpha = 1$

$R\sin\alpha = \sqrt{3}.$

$\therefore$ $R^2\left(\sin^2\alpha + \cos^2\alpha\right) = 3 + 1 = 4.$

$\therefore$ $R = 2$

and $\frac{R\sin\alpha}{R\cos\alpha} = \frac{\sqrt{3}}{1},$

$\therefore$ $\tan\alpha = \sqrt{3}$

$\alpha = \frac{\pi}{3}.$

$\therefore$ $2\cos\left(\theta - \frac{\pi}{3}\right) = 1, \quad -\frac{\pi}{3} \le \left(\theta - \frac{\pi}{3}\right) \le \frac{5\pi}{3}$

$\cos\left(\theta - \frac{\pi}{3}\right) = \frac{1}{2}$

$\theta - \frac{\pi}{3} = -\frac{\pi}{3}, \frac{\pi}{3}, \frac{5\pi}{3}.$

$\therefore$ $\theta = 0, \frac{2\pi}{3}, 2\pi.$

Method 2:

If $\theta = \pi$, $\cos\pi + \sqrt{3}\sin\pi = -1 + 0 \neq 1$,

$\therefore$ $\theta = \pi$ is not a solution.

If $\theta \neq \pi$, let $t = \tan\frac{\theta}{2}$.

$\therefore$ $\sin\theta = \frac{2t}{1+t^2}$ and $\cos\theta = \frac{1-t^2}{1+t^2}$.

$\therefore$ $\frac{1-t^2}{1+t^2} + \sqrt{3}\times\frac{2t}{1+t^2} = 1$

$1 - t^2 + 2\sqrt{3}t = 1 + t^2$

$2t^2 - 2\sqrt{3}t = 0$

$2t\left(t - \sqrt{3}\right) = 0.$

$\therefore$ $t = 0, \sqrt{3}.$

That is, $\tan\frac{\theta}{2} = 0,\ \sqrt{3}$.

$\therefore\ \frac{\theta}{2} = 0, \frac{\pi}{3}, \pi \quad (0 \le \theta \le 2\pi)$.

$\therefore\ \theta = 0, \frac{2\pi}{3}, 2\pi$.

c) $f(x) = x + \log_e x$

(i) The natural domain is $x > 0$ since $\log_e x$ is defined only for $x > 0$.

(ii) $y = f(x)$ is increasing if $f'(x) > 0$.

$\therefore\ f'(x) = 1 + \frac{1}{x} > 0$, since $x > 0$.

(iii) $f(0{\cdot}5) = 0{\cdot}5 + \log_e 0{\cdot}5$
$\doteqdot -0{\cdot}193 < 0$.

$f(1) = 1 + \log_e 1$
$= 1 > 0$.

The curve cuts the x axis between $x = 0{\cdot}5$ and $x = 1$, since the sign of $f(x)$ changes and $f(x)$ is continuous.

(iv) Let $f(x) = x + \log_e x$

$f'(x) = 1 + \frac{1}{x}$.

Let x_2 be a second approximation to the root of $x + \log_e x = 0$.

$\therefore\ x_2 = 0{\cdot}5 - \frac{f(0{\cdot}5)}{f'(0{\cdot}5)}$, by Newton's method,

$= 0{\cdot}5 - \frac{0{\cdot}5 + \log_e 0{\cdot}5}{1 + \frac{1}{0{\cdot}5}}$

$= 0{\cdot}564\ldots$

N.B. You need to use Newton's method again to see how many of these digits are significant, but this is not required by the question.

QUESTION 3

(a) $V = \pi\int_0^{\frac{\pi}{2}} (3\sin x)^2\ dx$

$= 9\pi\int_0^{\frac{\pi}{2}} \sin^2 x\ dx$

$= \frac{9\pi}{2}\int_0^{\frac{\pi}{2}} (1 - \cos 2x)\ dx$

$= \frac{9\pi}{2}\left[x - \frac{\sin 2x}{2}\right]_0^{\frac{\pi}{2}}$

$= \frac{9\pi}{2}\left[\left(\frac{\pi}{2} - 0\right) - (0 - 0)\right]$

$= \frac{9\pi^2}{4}$.

$\therefore$ Volume $= \frac{9\pi^2}{4}$ cubic units.

(b) $P(6) = \frac{1}{6},\ P(\bar{6}) = \frac{5}{6}$.

Probability of '6' on exactly 2 of 7 throws


$= {}^7C_2\left(\frac{1}{6}\right)^2\left(\frac{5}{6}\right)^5$

$= \frac{7\times 6}{1\times 2} \times \frac{1}{6} \times \frac{5^5}{6^5}$

$= \frac{21875}{93312}$

$\doteqdot 0{\cdot}2344$.

(c)



Data: AC is a diameter.

Construction: Join AB, PQ.

Proof: $\angle ABC = 90°$ (angle in semicircle, given AC is diameter)

$\angle CPQ = \angle ABC$ (exterior angle of cyclic quadrilateral equals interior opposite angle)

$\therefore\ \angle CPQ$ is a right angle.

(d) (i) $A(2\sin x + \cos x) + B(2\cos x - \sin x)$
$\equiv \sin x + 8\cos x$

$\therefore\ (2A - B)\sin x + (A + 2B)\cos x$
$\equiv \sin x + 8\cos x$.

Equating coefficients of $\sin x$ and $\cos x$,

$2A - B = 1$ —①

$A + 2B = 8$ —②

① × 2 → $4A - 2B = 2$ —③

② + ③ → $5A = 10$

$A = 2$.

Substitute $A = 2$ in ②:

$2B = 6$

$B = 3$.

$\therefore\ A = 2,\ B = 3$.

(ii) $\int \frac{\sin x + 8\cos x}{2\sin x + \cos x}\ dx$

$= \int \frac{2(2\sin x + \cos x) + 3(2\cos x - \sin x)}{2\sin x + \cos x}\ dx$ from (i)

$= \int 2\ dx + 3\int \frac{2\cos x - \sin x}{2\sin x + \cos x}\ dx$


$= 2x + 3\ln(2\sin x + \cos x) + C$.

$\left[\text{Note: } \frac{d}{dx}(2\sin x + \cos x) = 2\cos x - \sin x\right]$

QUESTION 4

(a) $\sum_{k=2}^{5}(-1)^k k = (-1)^2\times 2+(-1)^3\times 3$
$+(-1)^4\times 4+(-1)^5\times 5$
$= -2.$

(b)



(i) $x^2 = 4y$
$y = \frac{x^2}{4}$
$\frac{dy}{dx} = \frac{x}{2}.$

When $x = 2p$, $\frac{dy}{dx} = \frac{2p}{2} = p.$

Equation of tangent AP is
$y - y_1 = m(x - x_1)$
$y - p^2 = p(x - 2p)$
$y = px - p^2$ —①

(ii) Equation of normal BP is
$y - p^2 = -\frac{1}{p}(x - 2p).$

B lies on BP at $x = 0$.
When $x = 0$, $y = p^2 - \frac{1}{p}(-2p)$
$= p^2 + 2.$
$\therefore$ B is $(0,\ p^2+2)$.

(iii) Substitute $y = 0$ in ①: $0 = px - p^2$
$x = p.$
$\therefore$ A is $(p, 0)$.

If $C(x, y)$ is the midpoint of $A(p, 0)$ and $B(0, p^2+2)$, $x = \frac{p+0}{2}$ and $y = \frac{0+(p^2+2)}{2}$.

$x = \frac{p}{2}$ —②

$y = \frac{p^2+2}{2}$ —③

From ②, $p = 2x$.

Substitute in ③: $y = \frac{4x^2+2}{2} = 2x^2 + 1.$

But $p > 0$, $\therefore$ $x > 0$.

$\therefore$ Cartesian equation of locus of C is $y = 2x^2 + 1$, $x > 0$.

(c) (i) $\int_1^2 \frac{dx}{x} = \left[\ln x\right]_1^2$
$= \ln 2 - \ln 1$
$= \ln 2.$

(ii) $\int_1^2 \frac{dx}{x} \doteqdot \frac{b-a}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]$,
where $f(x) = \frac{1}{x}$, $a = 1$, $b = 2$.
$= \frac{2-1}{6}\left[\frac{1}{1}+4\times\frac{1}{1.5}+\frac{1}{2}\right]$
$= \frac{25}{36}$ $(= 0{\cdot}694\dot{}).$

(iii) $\ln 2 \doteqdot \frac{25}{36}$
$2 \doteqdot e^{\frac{25}{36}}$
$2^{\frac{36}{25}} \doteqdot e$ (raising both sides to power $\frac{36}{25}$)
$\therefore$ $e \doteqdot 2{\cdot}7132\ldots$
$= 2{\cdot}713$ (3 dec. places).

QUESTION 5

(a) Prove $(n+1)(n+2)\cdots(2n-1)2n$
$= 2^n[1\times 3\times\cdots\times(2n-1)]$

If $n = 1$, LHS $= 1 + 1 = 2$
RHS $= 2^1\times 1 = 2.$
$\therefore$ The statement is true for $n = 1$.

Assume statement is true for $n = k$, that is,
assume $(k+1)(k+2)\cdots(2k-1)2k$
$= 2^k[1\times 3\times\cdots\times(2k-1)].$ —①

Hence prove statement is true for $n = k+1$, that is, prove
$(k+2)(k+3)\cdots(2k+1)(2k+2)$
$= 2^{k+1}[1\times 3\times\cdots\times(2k+1)].$ —②

Now LHS
$= (k+2)(k+3)\cdots(2k+1)(2k+2)$
$= \frac{(k+1)(k+2)(k+3)\cdots(2k-1)2k(2k+1)(2k+2)}{k+1}$
$= \frac{2^k}{k+1}[1\times 3\times\cdots\times(2k-1)](2k+1)(2k+2)$, from ①
$= \frac{2^k}{\cancel{k+1}}[1\times 3\times\cdots\times(2k-1)](2k+1)2\cancel{(k+1)}$
$= 2^{k+1}[1\times 3\times\cdots\times(2k-1)(2k+1)]$
$=$ RHS.

$\therefore$ If the statement is true for $n = k$, it is also true for $n = k+1$. But it is true for $n = 1$.

$\therefore$ It is true for $n = 1+1 = 2$ and so on, that is, it is true for all integers $n \geq 1$.

(b) $f(x) = e^x - 1 - x$

(i) $f'(x) = e^x - 1$

$= 0$ only when $x = 0$.

$\therefore$ There is only one stationary point (at $x = 0$).

$f''(x) = e^x > 0$ for all x.

$\therefore$ The graph of $f(x)$ is concave up for all x.


Since $f(x)$ is continuous for all x (being made up of the sum and difference of continuous functions), the stationary point at $x = 0$ is both a local and absolute minimum.

(ii) When $x = 0$, $f(x) = e^0 - 1 - 0 = 0$.

$\therefore$ The least value of $f(x) = 0$.

$\therefore$ $f(x) \geq 0$ for all x.

(iii)



N.B. The gradient of $y = e^x - 1$ at $x = 0$ is 1, so $y = x$ is a tangent at $(0, 0)$.
This is also implied by (ii).

(iv) Inverse relation of $y = e^x - 1$ is $x = e^y - 1$.

That is, $e^y = x + 1$

$y = \log_e(x + 1)$

$\therefore$ $g^{-1}(x) = \log_e(x + 1)$.

(v) Domain of $g^{-1}(x)$ is $x + 1 > 0$, that is, $x > -1$.

(vi) $g(x) = e^x - 1$

$g^{-1}(x) = \log_e(1 + x)$.

The graphs of a pair of inverse functions are symmetrical about the line $y = x$.
The graph of $y = g(x)$ is above the graph of $y = x$ except at $x = 0$ where they coincide.

$\therefore$ The graph of $y = g^{-1}(x)$ is below the graph of $y = x$ except at $x = 0$ where they coincide.

$\therefore$ $\log_e(1 + x) \leq x$ for all $x > -1$.

QUESTION 6

(a) $x = \cos^2 3t, \quad t > 0.$ —①

(i) Substitute $x = \frac{3}{4}$ in ①:

$$\frac{3}{4} = \cos^2 3t$$

$$\cos 3t = \pm\frac{\sqrt{3}}{2}$$

$$3t = \frac{\pi}{6}, \ \ldots$$

$$t = \frac{\pi}{18}, \ \ldots$$

Particle is first at $x = \frac{3}{4}$ after $\frac{\pi}{18}$ seconds.

(ii) $v = \frac{dx}{dt} = 2\cos 3t \,.\, -3\sin 3t$

$= -3\sin 6t.$

When $t = \frac{\pi}{18}$, $v = -3\sin\left(6 \times \frac{\pi}{18}\right)$

$= -3\sin\frac{\pi}{3}$

$= \frac{-3\sqrt{3}}{2} < 0.$

Since $v < 0$, the particle is travelling in the negative direction.

(iii) $a = \frac{dv}{dt} = -3 \times 6\cos 6t$

$= -18\cos 6t.$

$\cos 6t = 2\cos^2 3t - 1$

(using $\cos 2x = 2\cos^2 x - 1$)

$= 2x - 1$, from ①.

$\therefore$ $a = -18(2x - 1)$.

(iv) $a = -18(2x - 1)$

$= -36\left(x - \frac{1}{2}\right)$

$= -6^2\left(x - \frac{1}{2}\right),$


which is of the form $\ddot{x} = -n^2(x - b)$, indicating simple harmonic motion with centre of oscillation at $x = \frac{1}{2}$.

(v) Period $= \frac{2\pi}{n}$ seconds

$= \frac{2\pi}{6}$ seconds

$= \frac{\pi}{3}$ seconds.


(b) (i)



$OD = 10$ cm (radius)

$\therefore$ $FD = 8$ cm (Pythagoras' theorem).

(ii) OBC is a sector of a circle, centre O, radius 10 cm.



$l = r\theta$


$\therefore \quad 4 = 10 \times \angle BOC.$

$\therefore \angle BOC = 0{\cdot}4$ radians

$\angle DFE = \angle BOC$

$\therefore \angle DFE = 0{\cdot}4$ radians.

(iii)




$ED^2 = FD^2 + FE^2 - 2 \times FD \times FE \cos 0{\cdot}4$ (by cosine rule)

$= 8^2 + 8^2 - 2 \times 8 \times 8 \cos 0{\cdot}4$


$= 128(1 - \cos 0{\cdot}4).$

{ *Alternatively:*



$ED = 2 \times 8 \sin 0{\cdot}2$

$= 16 \sin 0{\cdot}2.$ }



$\cos \angle EOD = \dfrac{OE^2 + OD^2 - ED^2}{2 \times OE \times OD}$


$= \dfrac{10^2 + 10^2 - 128(1 - \cos 0{\cdot}4)}{2 \times 10 \times 10}$

$\angle DOE = 0{\cdot}3192 \ldots$

$\approx 0{\cdot}319$ radians (3 dec. places).

QUESTION 7

(a) Initial conditions for velocity:



When $t = 0$,

$\dot{x} = 30\cos(5°), \quad \dot{y} = -30\sin(5°).$ —①

(i) $\ddot{x} = 0$

$\therefore \quad \dot{x} = C_1$ (constant).

$\therefore \quad \dot{x} = 30\cos(5°)$ from ①. —②

$x = \int 30\cos(5°)\ dt$

$= 30t\cos(5°) + C_2.$

When $t = 0, \ x = 0, \quad \therefore \ C_2 = 0.$

$\therefore \quad x = 30t\cos(5°).$

$\ddot{y} = -10$

$\therefore \quad \dot{y} = \int -10\ dt$

$= -10t + D_1.$

When $t = 0, \ \dot{y} = -30\cos(5°)$ from ①.

$\therefore \quad D_1 = -30\sin(5°)$

$\therefore \quad \dot{y} = -10t - 30\sin(5°).$ —③

$y = \int -10t - 30\sin(5°)\ dt$

$= -5t^2 - 30t\sin(5°) + D_2.$

When $t = 0, \ y = 0, \quad \therefore \ D_2 = 0.$

$\therefore \quad y = -30t\sin(5°) - 5t^2.$ —④

(ii) Ball strikes the ground when $y = -2$.

Substitute $y = -2$ in ④:

$-2 = -30t\sin 5° - 5t^2$

$5t^2 + 30t\sin 5° - 2 = 0$

$t = \dfrac{-30\sin 5° \pm \sqrt{(-30\sin 5°)^2 - 4 \times 5 \times (-2)}}{2 \times 5}$

$= \dfrac{-30\sin 5° + \sqrt{900\sin^2 5° + 40}}{10}$

(other answer negative and therefore irrelevant)


$= 0{\cdot}4229 \ldots$

$\therefore$ The ball strikes the ground after 0·42 seconds (2 dec. places).

(iii) When $t = 0{\cdot}4229$,

$\dot{x} = 30\cos(5°)$ from ②,

$\dot{y} = -4{\cdot}229 - 30\sin(5°)$, from ③.



$\tan\theta = \dfrac{4{\cdot}229 + 30\sin 5°}{30\cos 5°}$

$= 0{\cdot}228\,99 \ldots$

$\theta \doteqdot 12{\cdot}9°.$

Angle at which the ball strikes the ground is 13° (nearest degree).

(b) $(1-x)^n = \binom{n}{0} - \binom{n}{1}x^1 + \binom{n}{2}x^2 + \cdots + \binom{n}{n}(-1)^n x^n$

$$\left(1+\frac{1}{x}\right)^n = \binom{n}{0}\left(\frac{1}{x}\right)^0 + \binom{n}{1}\left(\frac{1}{x}\right)^1 + \binom{n}{2}\left(\frac{1}{x}\right)^2 + \cdots + \binom{n}{n}(-1)^n\left(\frac{1}{x}\right)^n.$$

The term in x^2 in $(1-x)^n\left(1+\frac{1}{x}\right)^n$ is

$$\binom{n}{2}\binom{n}{0}x^2\left(\frac{1}{x}\right)^0 - \binom{n}{3}\binom{n}{1}x^3\left(\frac{1}{x}\right)^1 + \binom{n}{5}\binom{n}{3}x^5\left(\frac{1}{x}\right)^3 + \cdots + (-1)^n\binom{n}{n}\binom{n}{n-2}x^n\left(\frac{1}{x}\right)^n.$$

$\therefore$ The coefficient of x^2 in $(1-x)^n\left(1+\frac{1}{x}\right)^n$ is

$$\binom{n}{2}\binom{n}{0} - \binom{n}{3}\binom{n}{1} + \cdots + (-1)^n\binom{n}{n}\binom{n}{n-2},$$

and this is the expression given in the question.

Now $(1-x)^n\left(1+\frac{1}{x}\right)^n = \left[(1-x)\left(1+\frac{1}{x}\right)\right]^n$

$$= \left(\frac{1}{x}-x\right)^n.$$

The general term of $\left(\frac{1}{x}-x\right)^n$ is

$$\binom{n}{r}\left(\frac{1}{x}\right)^{n-r}(-x)^r = \binom{n}{r}(-1)^r x^{2r-n}.$$

The term in x^2 has $2r-n=2$

$$r = \frac{n+2}{2}.$$

$\therefore$ The coefficient of $x^2 = \binom{n}{\frac{n+2}{2}}(-1)^{\frac{n+2}{2}}$,

and only exists if n is even,

$\left(\frac{n+2}{2}\right.$ must be an integer$\left.\right)$.

$$\therefore \binom{n}{2}\binom{n}{0} - \binom{n}{3}\binom{n}{1} + \cdots + (-1)^n\binom{n}{n}\binom{n}{n-2}$$

$$= \begin{cases} \binom{n}{\frac{n+2}{2}}(-1)^{\frac{n+2}{2}} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

END OF 3 UNIT (ADDITIONAL) AND 3/4 UNIT (COMMON) MATHEMATICS SOLUTIONS