itute.com

NSW BOS Mathematics Extension 1 Solutions 2007
Free download and print from www.itute.com
Do not photocopy © Copyright 2007 itute.com

Q1a $\left(1+\sqrt{5}\right)^3 = 1^3 + 3(1)^2\left(\sqrt{5}\right) + 3(1)\left(\sqrt{5}\right)^2 + \left(\sqrt{5}\right)^3 = 16 + 8\sqrt{5}$.

Q1b $x = \dfrac{(4)(3)+(19)(2)}{2+3} = 10$; $y = \dfrac{(5)(3)+(-5)(2)}{2+3} = 1$.

Q1c $\dfrac{d}{dx}\left(\tan^{-1}\left(x^4\right)\right) = \left(\dfrac{d}{d\left(x^4\right)}\left(\tan^{-1}\left(x^4\right)\right)\right)\left(\dfrac{d}{dx}\left(x^4\right)\right)$

$= \left(\dfrac{1}{1+\left(x^4\right)^2}\right)\left(4x^3\right) = \dfrac{4x^3}{1+x^8}$.

Q1d Line 1: $x-2y+3=0$, $y = \dfrac{1}{2}x + \dfrac{3}{2}$, $\dfrac{dy}{dx} = \dfrac{1}{2}$, $m_1 = \dfrac{1}{2}$.

Curve 2: $y = x^3+1$, $\dfrac{dy}{dx} = 3x^2$. At $x=1$, $\dfrac{dy}{dx} = 3$, $m_2 = 3$.

$\tan\theta = \dfrac{m_2 - m_1}{1+m_2m_1} = \dfrac{5/2}{1+3/2} = 1$, $\therefore \theta = \dfrac{\pi}{4}$.

Q1e Given $u = 25-x^2$, $\therefore -\dfrac{du}{dx} = 2x$.


When $x=3$, $u=16$; when $x=4$, $u=9$.

$\displaystyle\int_3^4 \frac{2x}{\sqrt{25-x^2}}dx = -\int_3^4 \frac{1}{\sqrt{u}}\frac{du}{dx}dx = -\int_{16}^9 u^{-\frac{1}{2}}du = \int_9^{16} u^{-\frac{1}{2}}du$

$= \left[2u^{\frac{1}{2}}\right]_9^{16} = 2$.

Q2a $LHS = \dfrac{1-\cos\theta}{\sin\theta} = \dfrac{2\sin^2\dfrac{\theta}{2}}{2\sin\dfrac{\theta}{2}\cos\dfrac{\theta}{2}} = \dfrac{\sin\dfrac{\theta}{2}}{\cos\dfrac{\theta}{2}} = \tan\dfrac{\theta}{2} = RHS$.

Q2bi



Q2bii The range is $[0, 2\pi]$.

Q2c $P(x) = x^2 + ax + b$,

$P(2) = 4 + 2a + b = 0$, $\therefore 2a + b = -4$.

$P(-1) = 1 - a + b = 18$, $\therefore a - b = -17$.

Solve simultaneously, $a = -7$ and $b = 10$.

Q2di $v = 50\left(1-e^{-0.2t}\right)$, $a = \dfrac{dv}{dt} = 50\left(0.2e^{-0.2t}\right) = 10e^{-0.2t}$.

When $t=10$, $a = 10e^{-2} \approx 1.4\ \text{ms}^{-2}$.


Q2dii Displacement in the first 10 seconds

$= \displaystyle\int_0^{10} 50\left(1-e^{-0.2t}\right)dt = \left[50\left(t+5e^{-0.2t}\right)\right]_0^{10} = 50\left(10+5e^{-2}-5\right) \approx 284$.

Distance fallen = 284 m.

Q3a $V = \displaystyle\int_0^3 \pi y^2 dx = \int_0^3 \frac{\pi}{9+x^2}dx = \left[\frac{\pi}{3}\tan^{-1}\frac{x}{3}\right]_0^3 = \frac{\pi}{3}\tan^{-1}1 = \frac{\pi^2}{12}$

Q3bi $y = \dfrac{x-2}{x-4} = 1 - \dfrac{2}{x-4}$. Vertical asymptote: $x=4$; horizontal asymptote: $y=1$.



Q3bii $\dfrac{x-2}{x-4} = 3$ when $x-2 = 3x-12$, i.e. $x=5$.

From graph, $\dfrac{x-2}{x-4} \le 3$ when $x<4$ or $x \ge 5$.

Q3ci $\ddot{x} = -e^{-2x}$, $\dfrac{d}{dx}\left(\dfrac{1}{2}v^2\right) = -e^{-2x}$,

$\dfrac{1}{2}v^2 = \displaystyle\int\left(-e^{-2x}\right)dx = \frac{1}{2}e^{-2x} + c_1$.

Given $x=0$ and $v=1$ when $t=0$,

$\therefore c_1 = 0$, $\therefore v^2 = e^{-2x}$, $\therefore \dot{x} = v = e^{-x}$. Note: $\dot{x} = -e^{-x}$ does not meet the initial conditions.

Q3cii $\dot{x} = e^{-x}$, $\therefore \dfrac{dx}{dt} = e^{-x}$, $\dfrac{dt}{dx} = e^x$, $\therefore t = \displaystyle\int e^x dx = e^x + c_2$.

Given $x=0$ and $v=1$ when $t=0$,

$\therefore c_2 = -1$ and $t = e^x - 1$. Hence $x = \log_e(t+1)$.

Q4ai $\Pr(X=2) = 0.1^2 = 0.01$

Q4aii $\Pr(X=2) = {}^{20}C_2(0.1)^2(0.9)^{18} \approx 0.285$

Q4aiii $\Pr(X>2) = 1 - \Pr(X=0) - \Pr(X=1) - \Pr(X=2)$

$= 1 - 0.9^{20} - {}^{20}C_1(0.1)(0.9)^{19} - 0.285 \approx 0.32$.

Q4b For $n=1$, $7^{2n-1}+5=12$ is divisible by 12.


For $n=k$, assume that $7^{2k-1}+5$ is divisible by 12.

For $n=k+1$,

$7^{2(k+1)-1}+5=7^2 7^{2k-1}+5=49\left(7^{2k-1}+5\right)-48\times 5$ is divisible by 12 because both $7^{2k-1}+5$ and 48 are divisible by 12.

Hence $7^{2n-1}+5=12$ is divisible by 12 for all integers $n\geq 1$.

Q4ci



$\angle QBX=\angle XAP$ (angles subtended on the circumference by the same arc).

$\angle XAP+\angle AXP=\angle PXD+\angle AXP=90°$, $\therefore \angle XAP=\angle PXD$.

$\angle PXD=\angle QXB$ (vertically opposite angles).

$\therefore \angle QXB=\angle QBX$.

Q4cii $\angle QXB=\angle QBX$, $\therefore \Delta QXB$ is isosceles and $QB=QX$.

Similarly, $\angle QXC=\angle QCX$, $\therefore \Delta QXC$ is isosceles and $QX=QC$. $\therefore QB=QC$ and hence Q bisects BC.

Q5ai Area of sector $OPQ=\frac{1}{2}r^2\theta$.

Area of $\Delta OPT=\frac{1}{2}r(PT)=\frac{1}{2}r(r\tan\theta)=\frac{1}{2}r^2\tan\theta$.

$\therefore \frac{1}{2}r^2\tan\theta=2\times\frac{1}{2}r^2\theta$, $\therefore \tan\theta=2\theta$.

Q5aii Let $f(\theta)=2\theta-\tan\theta$. $f'(\theta)=2-\sec^2\theta$.

Newton's method:

$$\theta_1\approx\theta_0-\frac{f(\theta_0)}{f'(\theta_0)}=1.15-\frac{2\times 1.15-\tan 1.15}{2-\sec^2 1.15}=1.1664.$$

Q5b There are 4! permutations of the four children together, and there are 3! ways in arranging a particular permutation of the four children, Mr Roberts and Mrs Roberts.

$\therefore$ total number of arrangements of the family of six with the children together is 3!4!.

Total number of arrangements of the family of six without restriction is 6!.

$\therefore \Pr(children\,together)=\frac{3!4!}{6!}=\frac{1}{5}$.

Q5c $\sin^{-1}x+\frac{1}{2}\cos^{-1}y=\frac{\pi}{3}$ (1)

$3\sin^{-1}x-\frac{1}{2}\cos^{-1}y=\frac{2\pi}{3}$ (2)

(1) + (2): $4\sin^{-1}x=\pi$, $\sin^{-1}x=\frac{\pi}{4}$, $\therefore x=\frac{1}{\sqrt{2}}$.

From (1), $\cos^{-1}y=2\left(\frac{\pi}{3}-\sin^{-1}x\right)=2\left(\frac{\pi}{3}-\frac{\pi}{4}\right)=\frac{\pi}{6}$,

$\therefore y=\frac{\sqrt{3}}{2}$.

Q5di Substitute $x=2aq$ and $y=aq^2$ into the equation of PQ:

$2aq+paq^2-2ap-ap^3=0$,

$a\left[2(q-p)+p\left(q^2-p^2\right)\right]=0$,

$a\left[2(q-p)+p(q-p)(q+p)\right]=0$,

$a(q-p)\left[2+p(q+p)\right]=0$.

Since $q-p\neq 0$ $\because q\neq p$, and $a\neq 0$,

$\therefore 2+p(q+p)=0$, hence $p^2+pq+2=0$.

Q5dii Gradient of $OP=\frac{ap^2}{2ap}=\frac{p}{2}$,

gradient of $OQ=\frac{aq^2}{2aq}=\frac{q}{2}$.

If $OP\perp OQ$, then $\frac{p}{2}\times\frac{q}{2}=-1$, $\therefore pq=-4$.

From Q5di, $p^2+pq+2=0$, $\therefore p^2-4+2=0$, hence $p^2=2$.

Q6ai $x=\sqrt{3}\sin 2t-\cos 2t+3$, $\dot{x}=2\sqrt{3}\cos 2t+2\sin 2t$,

$\ddot{x}=-4\sqrt{3}\sin 2t+4\cos 2t=-4\left(\sqrt{3}\sin 2t-\cos 2t\right)=-4(x-3)$.

Q6aii Period $=\frac{2\pi}{2}=\pi$ seconds.

Q6aiii $\dot{x}=2\sqrt{3}\cos 2t+2\sin 2t$,

$\dot{x}=A\cos(2t-\alpha)=A\cos 2t\cos\alpha+A\sin 2t\sin\alpha$.

$\therefore A\cos\alpha=2\sqrt{3}$ and $A\sin\alpha=2$.

Hence $A^2\cos^2\alpha+A^2\sin^2\alpha=16$,

$\therefore A=4$ and $\sin\alpha=\frac{1}{2}$, i.e. $\alpha=\frac{\pi}{6}$.

$\therefore \dot{x}=4\cos\left(2t-\frac{\pi}{6}\right)$.

Q6aiv $0\leq t\leq\pi$, $0\leq 2t\leq 2\pi$.

When $\dot{x}=2$, $4\cos\left(2t-\frac{\pi}{6}\right)=2$, $\cos\left(2t-\frac{\pi}{6}\right)=\frac{1}{2}$,

$2t-\frac{\pi}{6}=\frac{\pi}{3}, \frac{5\pi}{3}$. $\therefore t=\frac{\pi}{4}, \frac{11\pi}{12}$.

When $\dot{x}=-2$, $4\cos\left(2t-\frac{\pi}{6}\right)=-2$, $\cos\left(2t-\frac{\pi}{6}\right)=-\frac{1}{2}$,

$2t-\frac{\pi}{6}=\frac{2\pi}{3}, \frac{4\pi}{3}$. $\therefore t=\frac{5\pi}{12}, \frac{3\pi}{4}$.

© Copyright 2007 itute.com

Q6bi $f(x)=e^x-e^{-x}$, $f'(x)=e^x+e^{-x}>0$ for all values of x,
$\therefore f(x)$ is increasing for all values of x.

Q6bii Equation of $f(x)$: $y=e^x-e^{-x}$

Equation of inverse: $x=e^y-e^{-y}$.

$xe^y=(e^y)^2-1$, $(e^y)^2-xe^y-1=0$.

Apply quadratic formula: $e^y=\dfrac{x+\sqrt{x^2+4}}{2}$ since $e^y>0$.

$\therefore y=\log_e\left(\dfrac{x+\sqrt{x^2+4}}{2}\right)$, $\therefore f^{-1}(x)=\log_e\left(\dfrac{x+\sqrt{x^2+4}}{2}\right)$.

Q6biii $y=e^x-e^{-x}$, $\therefore x=\log_e\left(\dfrac{y+\sqrt{y^2+4}}{2}\right)$.

When $y=5$, $x=\log_e\left(\dfrac{5+\sqrt{5^2+4}}{2}\right)=1.65$.

Q7ai $y=kx^n$, $\dfrac{dy}{dx}=nkx^{n-1}$. $y=\log_e x$, $\dfrac{dy}{dx}=\dfrac{1}{x}$.

At $x=a$, $nka^{n-1}=\dfrac{1}{a}$, $\therefore a^n=\dfrac{1}{nk}$.

Q7aii At $x=a$, $ka^n=\log_e a$.

From Q7ai, $a^n=\dfrac{1}{nk}$, $\therefore k\left(\dfrac{1}{nk}\right)=\log_e a$, $\log_e a=\dfrac{1}{n}$, $a=e^{\frac{1}{n}}$.

$\therefore a^n=e=\dfrac{1}{nk}$. Hence $k=\dfrac{1}{en}$.

Q7bi $x=14t\cos\theta$, $\therefore t=\dfrac{x}{14\cos\theta}$.

$y=14t\sin\theta-4.9t^2$, $y=14\left(\dfrac{x}{14\cos\theta}\right)\sin\theta-4.9\left(\dfrac{x}{14\cos\theta}\right)^2$

$=x\tan\theta-\dfrac{x^2}{40\cos^2\theta}=x\tan\theta-\dfrac{x^2}{40}\sec^2\theta$

$=x\tan\theta-\dfrac{x^2}{40}(1+\tan^2\theta)=mx-\left(\dfrac{1+m^2}{40}\right)x^2$, where $m=\tan\theta$.

Q7bii At $x=10$ and when $m=2\pm\sqrt{3-0.4h}$,

$y=mx-\left(\dfrac{1+m^2}{40}\right)x^2$

$=10\left(2\pm\sqrt{3-0.4h}\right)-2.5\left(1+\left(2\pm\sqrt{3-0.4h}\right)^2\right)=h$.

Since $3-0.4h\geq 0$, $h\leq\dfrac{3}{0.4}=7.5$, $\therefore$ max h is 7.5 m.

Q7biii Given $m=2\pm\sqrt{3-0.4h}$.
When $h=3.9$, $m=0.8$ or 3.2.
When $h=5.9$, $m=1.2$ or 2.8.
$\therefore$ The other interval is $0.8\leq m\leq 1.2$.

Q7biv Let $y=0$ to find the range.


$mx-\left(\dfrac{1+m^2}{40}\right)x^2=0$, $x\left[m-\left(\dfrac{1+m^2}{40}\right)x\right]=0$,

$\therefore x=0$ or $x=\dfrac{40m}{1+m^2}$. $\therefore$ the range is $\dfrac{40m}{1+m^2}$ metres.

For $2.8\leq m\leq 3.2$, $\dfrac{40(3.2)}{1+3.2^2}\leq x\leq\dfrac{40(2.8)}{1+2.8^2}$,

$11.388\leq x\leq 12.670$.
Width of interval $=12.670-11.388\approx 1.282$ metres.

The range is maximum when $\theta°=45°$, i.e. $m=\tan 45°=1$,
which is within the interval $0.8\leq m\leq 1.2$.



When $m=0.8$, $x=\dfrac{40(0.8)}{1+0.8^2}=19.512$; when $m=1$, $x=20$;
when $m=1.2$, $x=19.672$.
Width of interval $=20-19.512\approx 0.488$ metres.

Please inform mathline@itute.com re conceptual, mathematical and/or typing errors.

© Copyright 2007 itute.com