itute.com

2009 NSW BOS Mathematics Extension 2 Solutions
Free download and print from www.itute.com
© Copyright 2009 itute.com

Q1a Let $u = \ln x$, $\dfrac{du}{dx} = \dfrac{1}{x}$.

$$\int \frac{\ln x}{x} dx = \int u \frac{du}{dx} dx = \int u du = \frac{1}{2}u^2 + c = \frac{1}{2}(\ln x)^2 + c .$$

Q1b Let $u = x$ and $v = \dfrac{1}{2}e^{2x}$, $\dfrac{du}{dx} = 1$, $\dfrac{dv}{dx} = e^{2x}$.

$$\int xe^{2x} dx = \int u \frac{dv}{dx} dx = \int u dv = uv - \int v du$$

$$= \frac{1}{2}xe^{2x} - \int \frac{1}{2}e^{2x} \frac{du}{dx} dx = \frac{1}{2}xe^{2x} - \int \frac{1}{2}e^{2x} dx = \frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} + c .$$

Q1c $$\int \frac{x^2}{1+4x^2} dx = \frac{1}{4}\int \frac{4x^2}{1+4x^2} dx = \frac{1}{4}\int \left(1 - \frac{1}{1+4x^2}\right) dx$$

$$= \frac{1}{4}\left(x - \frac{1}{2}\tan^{-1}(2x)\right) + c = \frac{1}{4}x - \frac{1}{8}\tan^{-1}(2x) + c .$$

Q1d $$\int_2^5 \frac{x-6}{x^2+3x-4} dx = \int_2^5 \left(\frac{2}{x+4} - \frac{1}{x-1}\right) dx$$ [Partial fractions]

$$= \left[2\ln(x+4) - \ln(x-1)\right]_2^5 = \left[\ln \frac{(x+4)^2}{x-1}\right]_2^5 = \ln \frac{9}{16} .$$

Q1e Let $x = \tan\theta$, $\dfrac{dx}{d\theta} = \sec^2\theta$, $\dfrac{d\theta}{dx} = \dfrac{1}{\sec^2\theta}$.

$$\int_1^{\sqrt{3}} \frac{1}{x^2\sqrt{1+x^2}} dx = \int_1^{\sqrt{3}} \frac{1}{\tan^2\theta\sqrt{1+\tan^2\theta}} dx = \int_1^{\sqrt{3}} \frac{1}{\tan^2\theta \sec\theta} dx$$

$$\int_1^{\sqrt{3}} \frac{\sec\theta}{\tan^2\theta} \frac{d\theta}{dx} dx = \int_{\frac{\pi}{4}}^{\frac{\pi}{3}} \frac{\sec\theta}{\tan^2\theta} d\theta = \int_{\frac{\pi}{4}}^{\frac{\pi}{3}} \frac{\cos\theta}{\sin^2\theta} d\theta$$

$$= \int_{\frac{\pi}{4}}^{\frac{\pi}{3}} \frac{1}{u^2} \frac{du}{d\theta} d\theta = \int_{\frac{1}{\sqrt{2}}}^{\frac{\sqrt{3}}{2}} \frac{1}{u^2} du$$


Let $u = \sin\theta$, $\dfrac{du}{d\theta} = \cos\theta$

$$= \left[-\frac{1}{u}\right]_{\frac{1}{\sqrt{2}}}^{\frac{\sqrt{3}}{2}} = \frac{3\sqrt{2} - 2\sqrt{3}}{3} .$$


Q2a $i^9 = (i^4)^2 i = i$.

Q2b $$\frac{-2+3i}{2+i} \times \frac{2-i}{2-i} = \frac{-1+8i}{5} = -\frac{1}{5} + \frac{8}{5}i .$$

Q2c



Q2d




Q2ei $z^5 = -1 = cis((2n+1)\pi)$, $\therefore z = cis\left(\dfrac{2n+1}{5}\pi\right)$.

Let $n = 0, \pm 1, \pm 2$.

$z = cis\left(-\dfrac{3\pi}{5}\right)$, $cis\left(-\dfrac{\pi}{5}\right)$, $cis\left(\dfrac{\pi}{5}\right)$, $cis\left(\dfrac{3\pi}{5}\right)$, $cis(\pi)$.

Q2eii



Q2fi Let $a + bi = \sqrt{3+4i}$.

$\therefore a^2 - b^2 = 3$ and $ab = 2$.


Hence $a = \pm 2$ and $b = \pm 1$.

The two square roots are $\pm(2+i)$.


Q2fii $$z = \frac{-i \pm \sqrt{i^2 - 4(-1-i)}}{2} = \frac{-i \pm \sqrt{3+4i}}{2} = \frac{-i \pm (2+i)}{2} .$$

$\therefore z = 1$ or $-1-i$.

Q3ai $y = \dfrac{1}{f(x)}$, undefined at $x = 0$ and $x = 4$.




Q3aii $y = [f(x)]^2$, x-intercepts become turning points.



itute.com

Q3aiii $y = f(x^2)$, $f((-x)^2) = f(x^2)$, $\therefore$ symmetrical about the y-axis. $f(2)$ is a local maximum value approximately, $\therefore x^2 \approx 2$,

$\therefore x \approx \pm\sqrt{2}$. $\dfrac{d}{dx} f(x^2) = 2x \dfrac{d}{d(x^2)} f(x^2)$, at $x = 0$, $\dfrac{d}{dx} f(x^2) = 0$.



Q3bi $x^2 + 2xy + 3y^2 = 18$,

$2x + 2y + 2x\dfrac{dy}{dx} + 6y\dfrac{dy}{dx} = 0$ [implicit differentiation],

$\dfrac{dy}{dx} = -\dfrac{x+y}{x+3y}$.

For horizontal tangent, $\dfrac{dy}{dx} = -\dfrac{x+y}{x+3y} = 0$,

$\therefore x + y = 0$, i.e. $y = -x$.

$\therefore x^2 + 2x(-x) + 3(-x)^2 = 18$, $2x^2 = 18$, $\therefore x = \pm 3$ and $y = \mp 3$.

The points are $(3,-3)$ and $(-3,3)$.

Q3c $P(x) = x^3 + ax^2 + bx + 5 = (x-1)^2(x+5) = x^3 + 3x^2 - 9x + 5$.

$\therefore a = 3$ and $b = -9$.

Q3d Solve $y = x + 1$ and $y = (x-1)^2$ simultaneously.

$x + 1 = (x-1)^2$, $3x - x^2 = 0$, $\therefore x = 0$ or 3.

$$V = \int_0^3 2\pi x\left(x + 1 - (x-1)^2\right)dx = 2\pi\int_0^3 \left(3x^2 - x^3\right)dx$$

$$= 2\pi\left[x^3 - \frac{x^4}{4}\right]_0^3 = \frac{27\pi}{2} \text{ cubic units.}$$

Q4ai $\dfrac{x^2}{a^2} + \dfrac{y^2}{b^2} = 1$, implicit differentiation, $\dfrac{2x}{a^2} + \dfrac{2y}{b^2}\dfrac{dy}{dx} = 0$.

Gradient of tangent $\dfrac{dy}{dx} = -\dfrac{b^2 x}{a^2 y}$,

and hence gradient of the normal at (x_0, y_0) is $\dfrac{a^2 y_0}{b^2 x_0}$.

Equation of the normal is $y - y_0 = \dfrac{a^2 y_0}{b^2 x_0}(x - x_0)$.

Q4aii x-intercept of the normal: Let $y = 0$, $-y_0 = \dfrac{a^2 y_0}{b^2 x_0}(x - x_0)$,

$$\therefore x = x_0 - \frac{b^2 x_0}{a^2} = \left(1 - \frac{b^2}{a^2}\right)x_0 = e^2 x_0.$$

Hence N is $(e^2 x_0, 0)$.

Q4aiii $PS = ePM$, $PS' = ePM'$.

$$\therefore \frac{PS}{PS'} = \frac{ePM}{ePM'} = \frac{e\left(\frac{a}{e} - x_0\right)}{e\left(\frac{a}{e} + x_0\right)} = \frac{a - ex_0}{a + ex_0} = \frac{ae - e^2 x_0}{ae + e^2 x_0} = \frac{NS}{NS'}.$$

Q4aiv $\dfrac{\sin\alpha}{NS'} = \dfrac{\sin\angle PNS'}{PS'}$, $\therefore \sin\alpha = \dfrac{NS'\sin\angle PNS'}{PS'}$.

$\dfrac{\sin\beta}{NS} = \dfrac{\sin\angle PNS}{PS}$, $\therefore \sin\beta = \dfrac{NS\sin\angle PNS}{PS}$.

$\therefore \dfrac{\sin\alpha}{\sin\beta} = \dfrac{PS.NS'\sin\angle PNS'}{PS'.NS\sin\angle PNS}$.

$\angle PNS'$ and $\angle PNS$ are supplementary angles,

$\therefore \sin\angle PNS' = \sin\angle PNS$.

From Q4aiii, $\dfrac{PS.NS'}{PS'.NS} = 1$.

$\therefore \dfrac{\sin\alpha}{\sin\beta} = 1$, i.e. $\sin\alpha = \sin\beta$.

$\therefore \alpha = \beta$ since $\alpha + \beta < \pi$.

Q4bi $0 < \alpha < \dfrac{\pi}{2}$.

Vertical: $N\sin\alpha + T\cos\alpha - mg = 0$ .....(1)

Horizontal: $T\sin\alpha - N\cos\alpha = mr\omega^2$ .....(2)

Q4bii From equation (1), $N = \dfrac{mg - T\cos\alpha}{\sin\alpha}$ .....(3)

Substitute in equation (2), $T\sin\alpha - \dfrac{(mg - T\cos\alpha)\cos\alpha}{\sin\alpha} = mr\omega^2$.

Simplify and write T as the subject, $T = m(g\cos\alpha + r\omega^2\sin\alpha)$.

Substitute in equation (3) and simplify,

$$N = \frac{mg - m(g\cos\alpha + r\omega^2\sin\alpha)\cos\alpha}{\sin\alpha} = m(g\sin\alpha - r\omega^2\cos\alpha).$$

Q4biii If $T = N$,

$m(g\cos\alpha + r\omega^2\sin\alpha) = m(g\sin\alpha - r\omega^2\cos\alpha)$,

$r\omega^2(\sin\alpha + \cos\alpha) = g(\sin\alpha - \cos\alpha)$,

$$\omega^2 = \frac{g(\sin\alpha - \cos\alpha)}{r(\sin\alpha + \cos\alpha)} = \frac{g}{r}\left(\frac{\frac{\sin\alpha - \cos\alpha}{\cos\alpha}}{\frac{\sin\alpha + \cos\alpha}{\cos\alpha}}\right) = \frac{g}{r}\left(\frac{\tan\alpha - 1}{\tan\alpha + 1}\right) \text{.....(4)}$$

Q4biv Solve $\omega^2 = \dfrac{g}{r}\left(\dfrac{\tan\alpha - 1}{\tan\alpha + 1}\right)$ for α in terms of r and ω,

$$\alpha = \tan^{-1}\left(\frac{g + r\omega^2}{g - r\omega^2}\right).$$

Since $\dfrac{g + r\omega^2}{g - r\omega^2} > 1$, $\therefore \alpha > \dfrac{\pi}{4}$.

Hence $\dfrac{\pi}{4} < \alpha < \dfrac{\pi}{2}$.

 © Copyright 2009 itute.com

itute.com

Q5ai Since AB is a diameter, $\therefore \angle ADB = 90°$.
ΔAKY and ΔABD are similar because $\angle A$ is common and $\angle AYK = \angle ADB = 90°$. Hence $\angle AKY = \angle ABD$.

Q5aii $\angle ABD = \angle ACD$ [Subtended by the same arc AD].
$\angle DKX = \angle ABD$ [From Q5ai].
$\therefore \angle DKX = \angle DCX$.
$\therefore CKDX$ is a cyclic quadrilateral because both angles are subtended by the same arc DX.

Q5aiii KX is a diameter since $\angle XDK = 90°$. $\therefore \angle XCK = 90°$.
$\angle ACB = \angle ADB = 90°$ [Subtended by diameter AB].
$\therefore \angle XCK$ and $\angle ACB$ form a straight angle.
Hence B, C and K are collinear.

Q5bi $I_n = \int_0^1 x^{2n+1}e^{x^2}dx$, $\therefore I_{n-1} = \int_0^1 x^{2(n-1)+1}e^{x^2}dx = \int_0^1 x^{2n-1}e^{x^2}dx$.

Let $u = x^{2n}$, $\frac{du}{dx} = 2nx^{2n-1}$, and $\frac{dv}{dx} = xe^{x^2}$, $v = \frac{1}{2}e^{x^2}$.

$$I_n = \left[u(x)v(x)\right]_0^1 - \int_0^1 v\frac{du}{dx}dx = \left[\frac{1}{2}x^{2n}e^{x^2}\right]_0^1 - n\int_0^1 x^{2n-1}e^{x^2}dx$$

$$= \frac{e}{2} - nI_{n-1}.$$

Q5bii $I_0 = \int_0^1 xe^{x^2}dx = \left[\frac{1}{2}e^{x^2}\right]_0^1 = \frac{1}{2}(e-1)$.

$I_1 = \frac{e}{2} - I_0 = \frac{1}{2}$, $I_2 = \frac{e}{2} - 2I_1 = \frac{e}{2} - 1$.

Q5ci

$f(x) = \frac{e^x - e^{-x}}{2} - x$, $f'(x) = \frac{e^x + e^{-x}}{2} - 1$, $f''(x) = \frac{e^x - e^{-x}}{2}$.

For $x > 0$, $e^x > e^{-x}$, $\therefore f''(x) > 0$.

Q5cii
Since $f''(x) > 0$ for $x > 0$, $\therefore f'(x)$ is increasing for $x > 0$.

When $x = 0$, $f'(0) = \frac{e^0 + e^0}{2} - 1 = 0$.

$\therefore f'(x) > 0$ for $x > 0$.

Q5ciii
Since $f'(x) > 0$ for $x > 0$, $\therefore f(x)$ is increasing for $x > 0$.

When $x = 0$, $f(0) = \frac{e^0 - e^0}{2} - 0 = 0$.

$\therefore f(x) = \frac{e^x - e^{-x}}{2} - x > 0$ for $x > 0$.

Hence $\frac{e^x - e^{-x}}{2} > x$ for $x > 0$.

Q6a Translate the solid 4 units left, and reflect in the y-axis.
The base equation becomes $x = y^2$, i.e. $y = \pm\sqrt{x}$, $x \in [0,4]$.
Width of rectangle $= 2\sqrt{x}$, height of rectangle $= x$.

$$V = \int_0^4 2\sqrt{x}.x\,dx = \int_0^4 2x^{\frac{3}{2}}dx = \left[\frac{4x^{\frac{5}{2}}}{5}\right]_0^4 = \frac{128}{5} \text{ cubic units.}$$

Q6bi Let β be the remaining zero.

Product of zeros $= -1$, $\therefore -1 \times \alpha \times \beta = -1$, $\therefore \beta = \frac{1}{\alpha}$.

Q6bii(1) $P(x)$ has real coefficients, $\therefore \beta = \overline{\alpha}$.

$\therefore \overline{\alpha} = \frac{1}{\alpha}$, i.e. $\alpha\overline{\alpha} = 1$, $|\alpha|^2 = 1$. Hence $|\alpha| = 1$.

Q6bii(2) Sum of zeros $= -q$, $\therefore -1 + \alpha + \overline{\alpha} = -q$,
$\alpha + \overline{\alpha} = 1 - q$.

$\therefore 2\,\mathrm{Re}(\alpha) = 1 - q$, $\mathrm{Re}(\alpha) = \frac{1-q}{2}$.

Q6ci $PQ = \sqrt{OP^2 - OQ^2} = \sqrt{x^2 + y^2 - r^2}$.

Q6cii $PQ = PR$, i.e. $\sqrt{x^2 + y^2 - r^2} = c - x$,
$x^2 + y^2 - r^2 = (c-x)^2$.
Expand and simplify: $y^2 = r^2 + c^2 - 2cx$.


Q6ciii $y^2 = -2c\left(x - \frac{r^2 + c^2}{2c}\right)$ and compare with $Y^2 = -4aX$,

$a = \frac{c}{2}$. Distance between vertex and focus $= a = \frac{c}{2}$.

x-coordinate of $S = \frac{r^2 + c^2}{2c} - \frac{c}{2} = \frac{r^2}{2c}$. See diagram below.

$\therefore S$ is $\left(\frac{r^2}{2c}, 0\right)$.

Q6civ



$PS = PM = \frac{r^2 + 2c^2}{2c} - x$, $PQ = PR = c - x$.

$\therefore PS - PQ = \left(\frac{r^2 + 2c^2}{2c} - x\right) - (c - x) = \frac{r^2}{2c}$, independent of x.

 © Copyright 2009 itute.com

itute.com

Q7ai(1) $\ddot{x} = g - rv$, where $g - rv > 0$. Acceleration due to gravity is greater than that due to air resistance.

$$v\frac{dv}{dx} = g - rv,\ \frac{dx}{dv} = \frac{v}{g-rv} = -\frac{1}{r}\left(\frac{g-rv-g}{g-rv}\right) = -\frac{1}{r}\left(1 - \frac{g}{g-rv}\right).$$

$$\therefore x = -\frac{1}{r}\int\left(1 - \frac{g}{g-rv}\right)dv,\ rx = -v - \frac{g}{r}\ln(g-rv) + c.$$

Given $x = 0$ and $v = 0$ initially, $0 = -\frac{g}{r}\ln(g) + c$,

$$\therefore c = \frac{g}{r}\ln(g),\ rx = \frac{g}{r}\ln\left(\frac{g}{g-rv}\right) - v,\ x = \frac{g}{r^2}\ln\left(\frac{g}{g-rv}\right) - \frac{v}{r}.$$

Q7ai(2) Given $g = 9.8$, $r = 0.2$, and $v = 30$ when $x = L$,

$$L = \frac{9.8}{0.2^2}\ln\left(\frac{9.8}{9.8 - 0.2\times 30}\right) - \frac{30}{0.2} \approx 82 \text{ metres.}$$

Q7aii When $x > L$, $x = e^{\frac{-t}{10}}(29\sin t - 10\cos t) + 92$.

$$v = \frac{dx}{dt} = -\frac{1}{10}e^{\frac{-t}{10}}(29\sin t - 10\cos t) + e^{\frac{-t}{10}}(29\cos t + 10\sin t)$$

$$= e^{\frac{-t}{10}}(-2.9\sin t + \cos t + 29\cos t + 10\sin t)$$

$$= e^{\frac{-t}{10}}(7.1\sin t + 30\cos t).$$

Find $x_{\max}$ by letting $v = 0$.

Since $e^{\frac{-t}{10}} \neq 0$, $7.1\sin t + 30\cos t = 0$, $\tan t = -\frac{30}{7.1}$,

$$t = \tan^{-1}\left(-\frac{30}{7.1}\right) \approx 1.8032 \text{ and}$$

$$x_{\max} = e^{\frac{-1.8032}{10}}(29\sin 1.8032 - 10\cos 1.8032) + 92 \approx 117.49.$$

Add the height of the jumper to $x_{\max}$, 119.49 metres, which is less than 125 metres. The jumper's head stays out of the water.

Q7bi $z = cis\theta$, $z^n = cisn\theta = \cos n\theta + i\sin n\theta$,

$z^{-n} = cis(-n\theta) = \cos(-n\theta) + i\sin(-n\theta) = \cos n\theta - i\sin n\theta$.

$\therefore z^n + z^{-n} = 2\cos n\theta$.

Q7bii $(2\cos\theta)^{2m} = (z + z^{-1})^{2m}$

$$= z^{2m} + \binom{2m}{1}z^{2m-1}z^{-1} + \binom{2m}{2}z^{2m-2}z^{-2} + \ldots + \binom{2m}{m-1}z^{m+1}z^{-m+1}$$

$$+ \binom{2m}{m}z^{m}z^{-m} + \binom{2m}{m+1}z^{m-1}z^{-m-1} + \ldots + \binom{2m}{2m-1}z^{1}z^{-2m+1} + z^{-2m}$$

$$= z^{2m} + \binom{2m}{1}z^{2m-2} + \binom{2m}{2}z^{2m-4} + \ldots + \binom{2m}{m-1}z^{2}$$

$$+ \binom{2m}{m} + \binom{2m}{m+1}z^{-2} + \ldots + \binom{2m}{2m-1}z^{-2m+2} + z^{-2m}$$

$$= z^{2m} + z^{-2m} + \binom{2m}{1}z^{2m-2} + \binom{2m}{2m-1}z^{-2m+2}$$

$$+ \binom{2m}{2}z^{2m-4} + \binom{2m}{2m-2}z^{-2m+4} + \ldots$$

$$+ \binom{2m}{m-1}z^{2} + \binom{2m}{m+1}z^{-2} + \binom{2m}{m}$$

$$= (z^{2m} + z^{-2m}) + \binom{2m}{1}(z^{2m-2} + z^{-2m+2})$$

$$+ \binom{2m}{2}(z^{2m-4} + z^{-2m+4}) + \ldots + \binom{2m}{m-1}(z^{2} + z^{-2}) + \binom{2m}{m}$$

$$= 2\cos 2m\theta + \binom{2m}{1}2\cos(2m-2)\theta + \binom{2m}{2}2\cos(2m-4)\theta + \ldots$$

$$+ \binom{2m}{m-1}2\cos 2\theta + \binom{2m}{m}$$

$$= 2\left[\cos 2m\theta + \binom{2m}{1}\cos(2m-2)\theta + \binom{2m}{2}\cos(2m-4)\theta + \ldots\right.$$

$$\left. + \binom{2m}{m-1}\cos 2\theta\right] + \binom{2m}{m}$$

Note:
$\binom{2m}{1} = \binom{2m}{2m-1}$
etc.

Q7biii $(2\cos\theta)^{2m} = 2^{2m}\cos^{2m}\theta$,

$$\int_0^{\frac{\pi}{2}}\cos^{2m}\theta d\theta = \frac{1}{2^{2m}}\int_0^{\frac{\pi}{2}}(2\cos\theta)^{2m}d\theta$$

$$= \frac{2}{2^{2m}}\int_0^{\frac{\pi}{2}}\left[\cos 2m\theta + \binom{2m}{1}\cos(2m-2)\theta + \binom{2m}{2}\cos(2m-4)\theta + \ldots\right.$$

$$\left. + \binom{2m}{m-1}\cos 2\theta + \frac{1}{2}\binom{2m}{m}\right]d\theta$$

$$= \frac{2}{2^{2m}}\left[\frac{\sin 2m\theta}{2m} + \binom{2m}{1}\frac{\sin(2m-2)\theta}{2m-2} + \binom{2m}{2}\frac{\sin(2m-4)\theta}{2m-4} + \ldots\right.$$

$$\left. + \binom{2m}{m-1}\frac{\sin 2\theta}{2} + \frac{1}{2}\binom{2m}{m}\theta\right]_0^{\frac{\pi}{2}}$$

$$= \frac{2}{2^{2m}}\left[\frac{1}{2}\binom{2m}{m}\frac{\pi}{2}\right]$$

Sine of 0 or multiple of π equals 0.

$$= \frac{\pi}{2^{2m+1}}\binom{2m}{m}$$

Q8ai $\cot\theta = \frac{1}{\tan\theta} = \frac{1}{\frac{2\tan\frac{\theta}{2}}{1-\tan^2\frac{\theta}{2}}} = \frac{1-\tan^2\frac{\theta}{2}}{2\tan\frac{\theta}{2}} = \frac{1}{2\tan\frac{\theta}{2}} - \frac{\tan\frac{\theta}{2}}{2}$

$$= \frac{1}{2}\cot\frac{\theta}{2} - \frac{1}{2}\tan\frac{\theta}{2}.$$

$$\therefore \cot\theta + \frac{1}{2}\tan\frac{\theta}{2} = \frac{1}{2}\cot\frac{\theta}{2}.$$

 © Copyright 2009 itute.com

itute.com

Q8aii From Q8ai, $\cot\theta+\frac{1}{2}\tan\frac{\theta}{2}=\frac{1}{2}\cot\frac{\theta}{2}$.

$\therefore \tan\frac{x}{2}=\cot\frac{x}{2}-2\cot x$.

$\therefore \sum_{r=1}^{n}\frac{1}{2^{r-1}}\tan\frac{x}{2^r}=\frac{1}{2^{n-1}}\cot\frac{x}{2^n}-2\cot x$ is true for $n=1$.

Assume it is true for $n=k$,

i.e. $\sum_{r=1}^{k}\frac{1}{2^{r-1}}\tan\frac{x}{2^r}=\frac{1}{2^{k-1}}\cot\frac{x}{2^k}-2\cot x$.

When $n=k+1$,

$\sum_{r=1}^{k+1}\frac{1}{2^{r-1}}\tan\frac{x}{2^r}=\sum_{r=1}^{k}\frac{1}{2^{r-1}}\tan\frac{x}{2^r}+\frac{1}{2^{(k+1)-1}}\tan\frac{x}{2^{k+1}}$

$=\frac{1}{2^{k-1}}\cot\frac{x}{2^k}-2\cot x+\frac{1}{2^k}\tan\frac{x}{2^{k+1}}$

$=\frac{1}{2^{k-1}}\left(\cot\frac{x}{2^k}+\frac{1}{2}\tan\frac{1}{2}.\frac{x}{2^k}\right)-2\cot x$

$=\frac{1}{2^{k-1}}\left(\frac{1}{2}\cot\frac{1}{2}.\frac{x}{2^k}\right)-2\cot x$

$=\frac{1}{2^k}\cot\frac{x}{2^{k+1}}-2\cot x$.

$\therefore \sum_{r=1}^{n}\frac{1}{2^{r-1}}\tan\frac{x}{2^r}=\frac{1}{2^{n-1}}\cot\frac{x}{2^n}-2\cot x$ is true for $n=k+1$.

Hence, it is true for integers $n\geq 1$.

Q8aiii

$\sum_{r=1}^{n}\frac{1}{2^{r-1}}\tan\frac{x}{2^r}=\frac{1}{2^{n-1}}\cot\frac{x}{2^n}-2\cot x=\frac{1}{2^{n-1}\tan\frac{x}{2^n}}-2\cot x$,

As $n\to\infty$, $\tan\frac{x}{2^n}\to\frac{x}{2^n}$.

Hence, $\lim_{n\to\infty}\sum_{r=1}^{n}\frac{1}{2^{r-1}}\tan\frac{x}{2^r}\to\frac{1}{2^{n-1}.\frac{x}{2^n}}-2\cot x=\frac{2}{x}-2\cot x$.

Q8aiv Let $x=\frac{\pi}{2}$, $\tan\frac{\pi}{4}+\frac{1}{2}\tan\frac{\pi}{8}+\frac{1}{4}\tan\frac{\pi}{16}+.....$

$=\lim_{n\to\infty}\sum_{r=1}^{n}\frac{1}{2^{r-1}}\tan\frac{\frac{\pi}{2}}{2^r}$

$=\frac{2}{\frac{\pi}{2}}-2\cot\frac{\pi}{2}=\frac{4}{\pi}$.

Q8b $A_{small}<\int_{n-1}^{n}\frac{1}{x}dx<A_{large}$, $\frac{1}{n}<\ln\left(\frac{n}{n-1}\right)<\frac{1}{n-1}$,

$e^{\frac{1}{n}}<\frac{n}{n-1}<e^{\frac{1}{n-1}}$, $e^{-\frac{1}{n}}>\frac{n-1}{n}>e^{-\frac{1}{n-1}}$, $e^{-\frac{1}{n-1}}<\frac{n-1}{n}<e^{-\frac{1}{n}}$,

$\left(e^{-\frac{1}{n-1}}\right)^n<\left(1-\frac{1}{n}\right)^n<\left(e^{-\frac{1}{n}}\right)^n$, $\therefore e^{-\frac{n}{n-1}}<\left(1-\frac{1}{n}\right)^n<e^{-1}$.

Q8ci Let A_1 represent A_1 wins, and A_1i represent A_1 wins in the ith draw.

For any one, in a single draw, probability of winning $=p=\frac{1}{n}$,

and probability of not winning $=q=1-\frac{1}{n}$.

$\Pr(A_1)=\Pr(A_1 1)+\Pr(A_1 2)+\Pr(A_1 3)+.....$

$\therefore W=p+q^n p+q^{2n}p+q^{3n}p+.....$

$\therefore W=p+q^n\left(p+q^n p+q^{2n}p+q^{3n}p+.....\right)$

$\therefore W=p+q^n W$.

Q8cii $W_m=p+q^n p+q^{2n}p+.....+q^{(m-1)n}p=\frac{p\left(1-q^{mn}\right)}{1-q^n}$.

[Sum of a GP with $a=p$ and $r=q^n$]

From Q8ci, $W=p+q^n W$, $\therefore W=\frac{p}{1-q^n}$.

$\therefore \frac{W_m}{W}=1-q^{mn}=1-\left(1-\frac{1}{n}\right)^{mn}$.

From Q8b, $e^{-\frac{n}{n-1}}<\left(1-\frac{1}{n}\right)^n<e^{-1}$, and $e^{-\frac{n}{n-1}}\to e^{-1}$ for large n.

$\therefore \left(1-\frac{1}{n}\right)^n\approx e^{-1}$ for large n.

$\therefore \frac{W_m}{W}=1-\left(1-\frac{1}{n}\right)^{mn}\approx 1-\left(e^{-1}\right)^m=1-e^{-m}$.

Please inform mathline@itute.com re conceptual, mathematical and/or typing errors.

 © Copyright 2009 itute.com